

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Repository Corticotropin Medications (Dermatomyositis and Polymyositis)

<u>PREFERRED</u>	<u>NON-PREFERRED</u>
<input type="checkbox"/> Purified Cortrophin™ Gel (repository corticotropin)	<input type="checkbox"/> HP Acthar® Gel (repository corticotropin) *Member must have tried and failed preferred Purified Cortrophin™ Gel and meet all applicable PA criteria below

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Member has diagnosis of **DERMATOMYOSITIS** OR **POLYMYOSITIS** with one of the following:

Idiopathic Inflammatory Myopathy

Refractory to conventional therapy or with severe organ-threatening manifestations

(Continued on next page)

1. **Diagnosis of Idiopathic Inflammatory Myopathy**, member **MUST** have tried and failed the therapies below **WITHIN THE PAST 6 MONTHS:**

- Prednisone 0.5-1mg/kg/day for 2-4 weeks, then taper for 2 weeks,

AND

- Prednisone **MUST** have been taken CONCURRENTLY WITH AN IMMUNOSUPPRESSIVE DRUG FOR **AT LEAST 90 DAYS** within the past 6 months (must note therapy tried):

<input type="checkbox"/> Methotrexate target dose 25mg/wk	<input type="checkbox"/> Azathioprine 2mg/kg IBW twice daily
<input type="checkbox"/> Mycophenolate mofetil, 500mg twice daily, increased by 500mg/wk until 1000mg twice daily	<input type="checkbox"/> Cyclophosphamide, 0.6-1g/m ² IV every 4 weeks or 1-2mg/kg/day orally, >3 months

2. For diagnosis that is refractory to conventional therapy or with severe organ-threatening manifestations, member must have tried and failed the therapies below **WITHIN THE PAST 6 MONTHS:**

- Methylprednisolone, 500-1000mg/day IV for 1-3 days for 3 months,

AND

- Member **MUST** have had trial and failure of **ONE** of the following therapies for at least 90 days **WITHIN THE PAST 6 MONTHS** (MUST note therapy tried):

<input type="checkbox"/> IVIG, 1g once a month for 1-6 months	<input type="checkbox"/> Cyclophosphamide, 0.6-1g/m ² IV every 4 weeks or 1-2 mg/kg/day orally, >3 months
<input type="checkbox"/> Rituximab, 1000 mg repeat on day 15, or 375 mg/m ² once weekly for 4 weeks	<input type="checkbox"/> Cyclosporine A, 3.0-3.5 mg/kg per day

Medication being provided by a Specialty Pharmacy - PropriumRx

****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.****

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.