SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

NON-PREFERRED

□ Acthar® Gel (repository corticotropin) 80 USP

<u>Drug Requested</u>: Repository Corticotropin Medications Dermatomyositis and Polymyositis

PREFERRED

□ Purified Cortrophin[™] Gel

adults only.

(repository corticotropin)	Units/mL 5 mL multi-dose vial	
(repository corticotropin)	□ Acthar® Gel (repository corticotropin) 40 USP	
	Units/0.5 mL single-dose prefilled SelfJect	
	injector	
	☐ Acthar® Gel (repository corticotropin) 80 USP Units/mL single-dose prefilled SelfJect injector	
	*Member must have tried and failed preferred	
	Purified Cortrophin [™] Gel and meet all applicable PA	
	criteria below	
MEMBER & PRESCRIBER INF	TORMATION: Authorization may be delayed if incomplete.	
Member Name:		
Member Sentara #:	Date of Birth:	
Prescriber Name:		
Prescriber Signature:		
Office Contact Name:		
Phone Number:	Fax Number:	
NPI #:		
DRUG INFORMATION: Authoriz	zation may be delayed if incomplete.	
Drug Name/Form/Strength:		
Dosing Schedule:	Length of Therapy:	
Diagnosis:	ICD Code, if applicable:	
Weight (if applicable):	Date weight obtained:	
• Acthar Gel single-dose pre-fille	d SelfJect injector is for subcutaneous administration by	

(Continued on next page)

(Continued from previous page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval.	To
support each line checked, all documentation, including lab results, diagnostics, and/or chart notes,	must be
provided or request may be denied. Check box below for the Diagnosis that applies.	

r with severe			
1 the therapies			
PRESSIVE tried):			
twice daily			
m^2 IV every 4 lly, > 3 months			
ntening ΓΗΕ PAST 6			
■ Member MUST have had trial and failure of ONE of the following therapies for at least 90 day WITHIN THE PAST 6 MONTHS (MUST note therapy tried):			
m^2 IV every 4 ally, > 3months			
g/kg per day			
tt/r			

Medication being provided by a Specialty Pharmacy – Proprium Rx

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.