SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

<u>Drug Requeste</u>d: Symdeko[®] (tezacaftor/ivacaftor)

MEMBER & PRESCRIBER INF	ORMATION: Authorization may be delayed if incomplete.			
Member Name:				
Member Sentara #:				
Prescriber Name:				
Prescriber Signature:				
Office Contact Name:				
hone Number: Fax Number:				
NPI #:				
DRUG INFORMATION: Authoriz	zation may be delayed if incomplete.			
Drug Form/Strength:				
Dosing Schedule:	Length of Therapy:			
Diagnosis:	ICD Code, if applicable:			
Weight (if applicable):	Date weight obtained:			
Recommended Dosing:				

- Pediatric patients aged 6 to less than 12 years weighing less than 30 kg: one tablet (containing tezacaftor 50 mg/ivacaftor 75 mg) in the morning and one tablet (containing ivacaftor 75 mg) in the evening, approximately 12 hours apart. SYMDEKO should be taken with fat-containing food.
- Adults and pediatric patients aged 12 years and older or pediatric patients aged 6 to less than 12 years weighing 30 kg or more: one tablet (containing tezacaftor 100 mg/ivacaftor 150 mg) in the morning and one tablet (containing ivacaftor 150 mg) in the evening, approximately 12 hours apart. SYMDEKO should be taken with fat-containing food.

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

☐ Member is <u>6 years of age or older</u> with a diagnosis of Cystic Fibrosis

(Continued on next page)

	Member must have <u>ONE</u> of the following mutation types in the cystic fibrosis transmembrane conductance regulator (CFTR) gene:				
	☐ Member is homozygous for the F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene (test result must be attached)				
	Member has <u>at least one mutation</u> in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to Symdeko [®] detected by an FDA-cleared test (test result must be attached)				
	Prescribing physician is a pulmonologist or has consulted with a pulmonologist who specializes in the treatment of Cystic Fibrosis				
	Baseline FEV ₁ within the last 30 days must be submitted (test results must be attached), unless the member is unable to perform a pulmonary function test (documentation required)				
	Number of pulmonary exacerbations or hospitalizations in the preceding 6 months must be noted:				
	Baseline body mass index must be noted:				
	Baseline liver function tests have been completed prior to initiating therapy and will be completed annually (labs must be attached)				
	Provider attests a baseline ophthalmic examination to monitor lens opacities/cataracts has been completed for pediatric members				
	Member will <u>NOT</u> be taking Symdeko [®] , in combination with any other CFTR modulator therapy (i.e., Orkambi [®] , Kalydeco [®] , Trikafta [™] , Alyftrek [™]); <u>NOTE</u> : concurrent therapy with these agents will <u>NOT</u> be approved				
	Member will avoid concomitant use of strong CYP3A inducers (e.g., rifampin, carbamazepine, phenytoin, phenobarbital, St. John's Wort) and strong or moderate CYP3A inhibitors (i.e. fluconazole, itraconazole)				
suppo	athorization: 12 months. Check below all that apply. All criteria must be met for approval. To ort each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be ded or request may be denied.				
	Member continues to meet all initial authorization criteria				
	Member has demonstrated disease response as indicated by <u>one or more</u> of the following (must submit current labs and chart notes):				
	 Decreased pulmonary exacerbations or hospitalizations compared to pretreatment baseline 				
	Stabilization of lung function as measured by FEV ₁ within the last year compared to baseline				
	☐ Improvement in quality of life, weight gain, or growth Member has NOT received a lung transplant				
	Member has experienced an absence of unacceptable toxicity from therapy (i.e., elevated transaminases				
_	(ALT or AST), development of cataracts or lens opacities)				

(Continued on next page)

2

Date of initiation of Symdeko® therapy:	Re-Authorization Date:			
Baseline FEV ₁ (last FEV ₁ prior to starting Symdeko®):	Current FEV ₁ (FEV ₁ AFTER last dose of Symdeko®):			
Baseline Weight:	Current weight:			
Baseline BMI:	Current BMI:			
Number of hospitalizations since last approval of Symdeko® must be noted				

Medication b	oeing provided b	y Specialty	Pharmacy –	Proprium	Rx
--------------	------------------	-------------	------------	----------	----

^{**}Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

^{*}Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *