SENTARA COMMUNITY PLAN (MEDICAID)

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-844-305-2331</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization can be delayed</u>.

Infliximab Category (Medical)

PREFERRED

Drug Requested: (Select drug below)

□ **Infliximab (JI745)** NDC (57894-0160-01)

100mg/ml; 1 vial=10 billable units							
NON-PREFERRED							
(Q	vsola™(infliximab- axxq) 25121) 0mg/ml; 1 vial=10 billable its		Inflectra® (infliximab- dyyb) (Q5103) 100mg/ml; 1 vial=10 billable units		Remicade® (infliximab) (J1745) NDC (57894-0030-01) 100mg/ml; 1 vial=10 billable units		
abd	enflexis® (infliximabda) (Q5104) Omg/ml; 1 vial=10 billable its		Zymfentra™ (infliximab- dyyb)* (J1748) *(Refer to Zymfentra Pharmacy PA form)				
MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.							
Membe	er Name:						
Member Sentara #: Date of Birth:							
Prescri	ber Name:						
Prescriber Signature:							
Office (Contact Name:						
Phone Number: Fax Number:							
NPI #:							
DRUG INFORMATION: Authorization may be delayed if incomplete.							
Drug N	Name/Form/Strength:						
Dosing Schedule:			Length of	f Th	erapy:		
Diagnosis:			ICD Code	ICD Code, if applicable:			
Weight (if applicable):			Date	e we	eight obtained:		

(Continued on next page)

- □ Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.
 - Effective July 1, 2023 per DMAS Infliximab is the preferred infliximab product. Remicade®, Inflectra®, Avsola®, Renflexis®, Zymfentra™ are non-preferred.
 - If requesting preferred drug, Infliximab: please check diagnosis and enter the dosing below that applies. No additional prior authorization criteria is required.
 - If requesting a non-preferred drug, Renflexis®, Remicade®, Inflectra®, Avsola® or Zymfentra™ please complete all of the required prior authorization criteria.

DIAGNOSIS	Recommended Dose			
☐ Ankylosing Spondylitis (AS) Dosing:	• 5mg/kg at week 0, 2 and 6, then every 6 weeks thereafter			
☐ Crohn's Disease (CD) Dosing:	• 5mg/kg at week 0, 2 and 6, then every 8 weeks thereafter. Max dose 10mg/kg every 8 weeks			
 □ Pediatric Crohn's Disease (CD) Age ≥ 6 years Dosing:	• 5mg/kg at week 0, 2 and 6 weeks, then every 8 weeks thereafter			
☐ Plaque Psoriasis (Ps) Dosing:	• 5mg/kg at week 0, 2 and 6, then every 8 weeks thereafter			
☐ Psoriatic Arthritis (PsA) Dosing:	• 5mg/kg at week 0, 2 and 6, then every 8 weeks thereafter			
☐ Rheumatoid Arthritis (RA) in combination with methotrexate Dosing:	• 3mg/kg at week 0, 2 and 6, then every 8 weeks thereafter. Max dose 10mg/kg every 8 weeks or 3mg/kg every 4 weeks			
☐ Ulcerative Colitis (UC) Dosing:	• 5mg/kg at week 0, 2 and 6, then every 8 weeks thereafter			
☐ Pediatric Ulcerative Colitis Age ≥ 6 years Dosing:	• 5mg/kg at week 0, 2 and 6, then every 8weeks thereafter			

CLINICAL CRITERIA: Check below all that apply. All criteria/diagnosis must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied. If requesting an increase in dose, recent lab values and symptoms documenting active disease must be submitted with request.

(Continued on next page)

□ Diagnosis: Rheumatoid Arthritis or Psoriatic Arthritis							
	Check diagnosis:						
	□ Rheumatoid Arthritis <u>OR</u>	<u>.</u>		Psoriatic Arthritis			
	Trial and failure to ONE of the preferred drugs below:						
	☐ Humira [®]	□ En	□ Enbrel [®]				
	☐ Trial and failure to infliximab therapy						
□ D	iagnosis: Ankylosing Spondylitis						
	Diagnosed for active ankylosing spondylitis						
	Trial and failure of ONE of the preferred drugs below:						
	☐ Humira [®]	□ Er	nbrel	®			
	☐ Trial and failure to infliximab therapy						
□ D	iagnosis: Plaque Psoriasis						
	□ Diagnosed for Plaque Psoriasis						
	Trial and failure of ONE of the preferred	drugs below:					
	□ Humira [®] □ Enbrel [®]						
	Trial and failure to infliximab therapy						
□ D	iagnosis: Crohn's Disease <u>OR</u> Ocu	ılar Sarcoid	losis	- moderate to severe			
	Diagnosed for:						
	☐ Crohn's Disease OR	<u>.</u>	Į	□ Ocular Sarcoidosis			
	Member is 6 years of age or older for diagnosis of Crohn's disease						
	Trial and failure to Humira® AND Inflixing	mab therapy for	or Cr	rohn's disease indication			
□ D	iagnosis: Moderate-to-severe Ulce	rative Colit	is di	isease			
	☐ Diagnosed for moderate-to-severe Ulcerative Colitis						
	Member is 6 years of age or older						
	Trial and failure of BOTH of the preferred biologics below:						
	□ Humira [®]	□ In:	flixir	nab			

(Continued on next page)

Medication being provided by: Please check applicable box below.					
□ Location/site of drug administration:					
NPI or DEA # of administering location:					
<u>OR</u>					
□ Specialty Pharmacy – PropriumRx					
For urgent reviews: Practitioner should call Sentara Health Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.					
Use of samples to initiate therapy <u>does not</u> meet step-edit/preauthorization criteria.					
*Previous therapies will be verified through pharmacy paid claims or submitted chart notes					