

SENTARA COMMUNITY PLAN (MEDICAID)

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-844-305-2331.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization will be delayed.**

Infliximab Category (MEDICAL)

DRUG REQUESTED: (Select drug below)

PREFERRED			
<input type="checkbox"/> Infliximab (J1745) NDC (57894-0160-01)			
NON-PREFERRED			
<input type="checkbox"/> Avsola™ (infliximab-axxq) (Q5121)	<input type="checkbox"/> Inflectra® (infliximab-dyyb) (Q5103)	<input type="checkbox"/> Remicade® (infliximab) (J1745) NDC (57894-0030-01)	<input type="checkbox"/> Renflexis® (infliximab-abda) (Q5104)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

- ☐ Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.
- **Effective July 1, 2023 per DMAS Infliximab is the preferred infliximab product. Remicade®, Inflectra®, Avsola®, Renflexis® are non-preferred.**
- If requesting preferred drug, Infliximab: please check diagnosis and enter the dosing below that applies. No additional prior authorization criteria is required.
- **If requesting a non-preferred drug, Renflexis®, Remicade®, Inflectra® or Avsola® please complete all of the required prior authorization criteria.**

(Continued on next page)

DIAGNOSIS	Recommended Dose
<input type="checkbox"/> Ankylosing Spondylitis (AS) Dosing: _____	<ul style="list-style-type: none"> 5mg/kg at week 0, 2 and 6, then every 6 weeks thereafter
<input type="checkbox"/> Crohn's Disease (CD) Dosing: _____	<ul style="list-style-type: none"> 5mg/kg at week 0, 2 and 6, then every 8 weeks thereafter. Max dose 10mg/kg every 8 weeks
<input type="checkbox"/> Pediatric Crohn's Disease (CD) <input type="checkbox"/> Age \geq 6 years Dosing: _____	<ul style="list-style-type: none"> 5mg/kg at week 0, 2 and 6 weeks, then every 8 weeks thereafter
<input type="checkbox"/> Plaque Psoriasis (Ps) Dosing: _____	<ul style="list-style-type: none"> 5mg/kg at week 0, 2 and 6, then every 8 weeks thereafter
<input type="checkbox"/> Psoriatic Arthritis (PsA) Dosing: _____	<ul style="list-style-type: none"> 5mg/kg at week 0, 2 and 6, then every 8 weeks thereafter
<input type="checkbox"/> Rheumatoid Arthritis (RA) in combination with methotrexate Dosing: _____	<ul style="list-style-type: none"> 3mg/kg at week 0, 2 and 6, then every 8 weeks thereafter. Max dose 10mg/kg every 8 weeks or 3mg/kg every 4 weeks
<input type="checkbox"/> Ulcerative Colitis (UC) Dosing: _____	<ul style="list-style-type: none"> 5mg/kg at week 0, 2 and 6, then every 8 weeks thereafter
<input type="checkbox"/> Pediatric Ulcerative Colitis <input type="checkbox"/> Age \geq 6 years Dosing: _____	<ul style="list-style-type: none"> 5mg/kg at week 0, 2 and 6, then every 8 weeks thereafter

CLINICAL CRITERIA: Check below all that apply. All criteria/diagnosis must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied. If requesting an increase in dose, recent lab values and symptoms documenting active disease must be submitted with request.

☐ **Diagnosis: Rheumatoid Arthritis or Psoriatic Arthritis**

☐ Check diagnosis:

☐ **Rheumatoid Arthritis**

OR

☐ **Psoriatic Arthritis**

AND

(Continued on next page)

- ☐ Prescriber is a **Rheumatologist**

AND

- ☐ Trial and failure of **ONE** of the **PREFERRED** drugs below:

<input type="checkbox"/> azathioprine	<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> 6-mercaptopurine	<input type="checkbox"/> methotrexate
<input type="checkbox"/> leflunomide	<input type="checkbox"/> aminosalicylates	<input type="checkbox"/> auranofin	<input type="checkbox"/> sulfasalazine
<input type="checkbox"/> Other			

AND

- ☐ Trial and failure to Humira® or Enbrel® **AND** Infliximab therapy

☐ **Diagnosis: Ankylosing Spondylitis**

- ☐ Diagnosed for **active ankylosing spondylitis**

AND

- ☐ Prescribed by or in consultation with a **Rheumatologist**

AND

- ☐ Trial and failure, contraindication, or intolerance to **TWO** NSAIDs

AND

- ☐ Trial and failure of **ONE** of the **PREFERRED** drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®
----------------------------------	----------------------------------

AND

- ☐ Trial and failure to Infliximab therapy

☐ **Diagnosis: Plaque Psoriasis**

- ☐ Diagnosed for **Plaque Psoriasis**

AND

- ☐ Prescribed by or in consultation with a Dermatologist

AND

- ☐ Trial and failure of **ONE** of the **PREFERRED** drugs below:

<input type="checkbox"/> acitretin	<input type="checkbox"/> cyclosporine	<input type="checkbox"/> methotrexate
------------------------------------	---------------------------------------	---------------------------------------

AND

(Continued on next page)

- ☐ Trial and failure to Humira® or Enbrel® AND Infliximab therapy

☐ **Diagnosis: Crohn's Disease OR Ocular Sarcoidosis - moderate to severe with inadequate response to:**

- ☐ Diagnosed for:

- ☐ Crohn's Disease OR ☐ Ocular Sarcoidosis

AND

- ☐ Prescribed by or in consultation with a Gastroenterologist

OR

- ☐ Prescribed by or in consultation with an Ophthalmologist

AND

- ☐ Inadequate response to high dose steroids (e.g., 40-60 mg prednisone)

AND

- ☐ Trial and failure of ONE of the PREFERRED drugs below:

<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> 6-mercaptopurine	<input type="checkbox"/> methotrexate	<input type="checkbox"/> azathioprine
<input type="checkbox"/> aminosalicylates	<input type="checkbox"/> auranofin	<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> Other: _____			

AND

- ☐ Trial and failure to Humira® AND Infliximab therapy for Crohn's disease indication

☐ **Diagnosis: Moderate-to-severe Ulcerative Colitis disease**

- ☐ Diagnosed for moderate-to-severe Ulcerative Colitis

AND

- ☐ Prescribed by or in consultation with a Gastroenterologist

AND

- ☐ Inadequate response to high dose steroids (e.g. 40-60 mg prednisone)

AND

(Continued on next page)

- ☐ Trial and failure of **ONE** of the **PREFERRED** drugs below:

<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> 6-mercaptopurine	<input type="checkbox"/> methotrexate	<input type="checkbox"/> azathioprine
<input type="checkbox"/> aminosaliclates	<input type="checkbox"/> auranofin	<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> Other: _____			

AND

- ☐ Trial and failure to Humira® **AND** Infliximab therapy

Medication being provided by (check below that applies):

- ☐ Location/site of drug administration: _____
NPI or DEA # of administering location: _____

OR

- ☐ Specialty Pharmacy - PropriumRx

For urgent reviews: Practitioner should call Sentara Health Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****