SENTARA HEALTH PLANS

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-844-668-1550</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If information provided is not complete, correct, or legible, authorization can be delayed.

For Medicare Members: Medicare Coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <u>https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx</u>. Additional indications may be covered at the discretion of the health plan.

Intravitreal Complement Inhibitors (Medical)

Drug Requested: (Select drug below)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name:	
Member Sentara #:	
Prescriber Name:	
	Date:
Office Contact Name:	
Phone Number:	Fax Number:
DEA OR NPI #:	
DRUG INFORMATION: Authoriza	ation may be delayed if incomplete.
Drug Form/Strength:	
Dosing Schedule	Length of Therany

	Eengen of Therapy:
Diagnosis:	ICD Code, if applicable:
Weight:	Date:

□ Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

Quantity Limit:

- Izervay Maximum 1 vial per eye per 28 days
- Syfovre Maximum 1 vial per eye per 25 days; 1 vial = 15 billable units

(Continued on next page)

Recommended Dosage:

- Izervay 2 mg (0.1 mL of 20 mg/mL solution) administered by intravitreal injection to each affected eye once monthly (approximately every 28 ± 7 days) for up to 12 months.
- Syfovre 15 mg (0.1 mL of 150 mg/mL solution) administered by intravitreal injection to each affected eye once every 25 to 60 days

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 12 months

- **D** Provider is an Optometry Specialist or an Ophthalmologist
- □ Member has a diagnosis of geographic atrophy (GA) confirmed by <u>ALL</u> the following:
 - □ Defined by a phenotype of central geographic atrophy having 1 or more zones of well demarcated retinal pigmented epithelium (RPE) and/or choriocapillaris atrophy
 - Disease is secondary to age-related macular degeneration (AMD)
 - □ Conditions other than AMD have been ruled out (e.g., Stargardt disease, cone rod dystrophy, toxic maculopathies such as plaquenil maculopathy in either eye)
- □ Provider has submitted member's baseline for at least <u>ONE</u> of the following:
 - □ Best Corrected visual acuity (BCVA) score: _____
 - □ Fundus autofluorescence (FAF) imaging: _____
 - □ Optical coherence tomography (OCT):
- □ Requested medication will <u>NOT</u> be used in combination with other intravitreal complement inhibitor therapies
- □ Member does <u>NOT</u> have category 6, or higher, visual impairment or blindness (i.e., no light perceptiontotal blindness)
- □ Provider is requesting <u>ONE</u> of the following dosing frequencies:
 - □ Monthly
 - Every other month

<u>Reauthorization</u>: 12 months. All criteria that apply must be checked for approval. To support each line checked all documentation (lab results, diagnostics, and/or chart notes) must be provided or request may be denied.

- □ Member has had disease stabilization or slowing rate of disease progression while on therapy compared to previous baseline as measured by at least <u>ONE</u> the following:
 - □ Member's best corrected visual acuity (BCVA) has improved or stabilized from baseline
 - □ Member's fundus autofluorescence (FAF) imaging has improved or stabilized from baseline
 - □ Member's optical coherence tomography (OCT) has improved or stabilized from baseline

- Provider attests continued administration is necessary for the maintenance treatment of the condition and both member and provider have discussed potential decrease in frequency of administration if receiving monthly
- Member has experienced an absence of unacceptable toxicity from the drug including but not limited to endophthalmitis, retinal detachment, neovascular (wet) AMD or choroidal neovascularization, intraocular inflammation (e.g., vitritis, vitreal cells, iridocyclitis, uveitis, anterior chamber cells, iritis, and anterior chamber flare), increased intraocular pressure, that cannot be adequately treated

Medication being provided by: Please check applicable box below.

Location/site of drug administration:

NPI or DEA # of administering location:

<u>OR</u>

D Specialty Pharmacy – Proprium Rx

For urgent reviews: Practitioner should call Sentara Health Plans Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health Plan's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

** Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *