

SENTARA COMMUNITY PLAN (MEDICAID)

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-844-305-2331. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If information provided is not complete, correct, or legible, authorization can be delayed.

Drug Requested: Tepezza™ (teprotumumab-trbw) Injection (J3241) (Medical)
NDC: 75987-0130-15

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

- Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

Dosing Limits:

- IV: 10 mg/kg as a single dose, followed by 20 mg/kg every 3 weeks for 7 additional doses
- Maximum 6-month authorization and maximum of 8 infusions
- NDC: 75987-0130-15; 500 mg vial (1 each) = 50 billable units

Warnings and Precautions:

- Tepezza may cause severe hearing problems including hearing loss, which in some cases may be permanent. Tell your doctor if you have any signs or symptoms of hearing problems or changes in hearing.
- Inflammatory bowel disease (IBD), such as Crohn's disease or ulcerative colitis, Tepezza may make your IBD symptoms worse.

(Continued on next page)

Renewal: None

Part A:

- Globe protrusion: 13.9mm in Asian males, 16.5mm in white males, 18.5mm in African American males. Adult females have lower exophthalmometry readings than adult males with average of 15.4mm in white women and 17.8mm in African American women

Provider please note: The use of teprotumumab (Tepezza) does **NOT** meet the definition of medical necessity for the treatment of inactive TED (CAS \leq 2) due to insufficient evidence in peer-reviewed medical literature and guidelines to support safety, efficacy and net health outcomes

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Authorization Criteria: Length of Approval – 6 months (Maximum of 8 doses per lifetime)

- Member is \geq 18 years of age

AND

- Prescriber is a specialist in ophthalmology, endocrinology, oculoplastic surgery or neuro-ophthalmology

AND

- Member has a clinical diagnosis of Thyroid Eye disease that is related to Graves’ Orbitopathy

AND

- Documentation of one (1) of the following:

- Lid retraction of \geq 2mm
- Proptosis of \geq 3mm above the normal values for race and sex (see Part A)
- exophthalmometer \geq 20mm

AND

- Symptoms began within 12 months of the date of prior authorization form submission

AND

- Member has a Clinical Activity Score of at least \geq 4 (please complete table below):

Parameters Assessed	Spontaneous retrobulbar pain	Pain on attempted upward or downward gaze	Eyelid erythema	Eyelid edema	Conjunctival hyperaemia	Conjunctival chemosis	Inflammation of caruncle or plica
Score: Present=1 or Absent=0							
Total:							

(Continued on next page)

AND

- Member is **NOT** currently smoking and has not smoked within the last 30 days

AND

- If member has diabetes, disease must be adequately controlled (HbA1c <9%) and prescriber must attest to monitoring glycemic levels prior to starting Tepezza[®]

AND

- Member must have tried and failed 6 weeks of intravenous methylprednisolone at dose of ≥ 500 mg/week
Date started: _____

AND

- Member must have been compliantly taking thyroid medication for the last 3 months and must be euthyroid **OR** has lab levels within the following ranges (**must submit labs completed within the last 30 days**):
 - Free triiodothyronine (FT3): 3.5-6.5 pmol/liter OR 230-619 pg/d
 - Free Thyroxine (FT4): 11.5-22.7 pmol/liter OR 0.7-1.9 ng/dl
 - Thyrotropin (TSH): 0.55-4.78mIU/liter OR 0.5-6 uU/ml

AND

- Requested dosing is in accordance with the United States Food and Drug Administration and medication will be prescribed for a maximum of 8 doses per lifetime

Reauthorization: NOT COVERED. The clinical benefit of Tepezza has not been demonstrated beyond 8 infusions in phase 3 clinical trials. The continued use of Tepezza beyond 8 infusions in the patient's lifetime is unproven and not medically necessary

Medication being provided by (check applicable box(es) below):

- Physician's office **OR** Specialty Pharmacy – PropriumRx

For urgent reviews: Practitioner should call Sentara Health Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****