

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Taltz[®] SQ (ixekizumab) (self-administered, Pharmacy) (Non-Preferred)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

Recommended Dose:

Indication	Dosage
<input type="checkbox"/> Moderate-to-Severe Plaque Psoriasis (Adults) – who are candidates for systemic therapy or phototherapy	<ul style="list-style-type: none"> • Two 80mg injections initially for 14 days, • Then, two 80mg/28days until week 12 • Then, one 80mg injection every 28 days
<input type="checkbox"/> Moderate-to-Severe Plaque Psoriasis (Children ≥ 6 years) - who are candidates for systemic therapy or phototherapy	In pediatrics weight based: <ul style="list-style-type: none"> • >50 kg: Two 80 mg injections/28 days initially, then one 80 mg injection/28 days • 25-50 kg: Two 40 mg injections/28 days initially, then one 40 mg injection/28 days • <25kg: Two 20 mg injections/28 days initially, then one 20 mg injection/28 days
<input type="checkbox"/> Active Psoriatic Arthritis (PsA)	<ul style="list-style-type: none"> • Two 80mg injections/28days, then one 80mg injection every 4 weeks

(Continued on next page)

<input type="checkbox"/> Active Ankylosing Spondylitis	<ul style="list-style-type: none"> Two 80mg injections/28days, then one 80mg injection every 4 weeks
<input type="checkbox"/> Active non-radiographic axial spondyloarthritis (nr-axSpA) with objective signs of inflammation (Adults)	<ul style="list-style-type: none"> One 80mg injection subcutaneously every 4 weeks

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

DIAGNOSIS: (Check below the applicable diagnosis or authorization will be delayed or denied)

Moderate to Severe Plaque Psoriasis

- Moderate-to-severe active **Chronic Plaque Psoriasis** who are candidates for systemic therapy or phototherapy
- Member is at least 6 years of age or older
- Trial and failure of at least **TWO (2)** topical treatments, such as corticosteroids, calcipotriene, coal tar, tazarotene, or anthralin. List drugs below:
 1. _____ 2. _____
- Trial and failure of, contraindication, or adverse reaction to methotrexate
- Trial and failure of **TWO (2)** of the **PREFERRED** drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

Active Psoriatic Arthritis (PsA)

- Trial and failure of, contraindication, or adverse reaction to methotrexate
- Trial and failure of **TWO (2)** of the **PREFERRED** drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

Ankylosing Spondylitis

- Trial and failure of, contraindication, or adverse reaction to methotrexate
- Trial and failure of **TWO (2)** of the **PREFERRED** drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

(Continued on next page)

Non-Radiographic Axial Spondyloarthritis

- Member has objective signs of inflammation
- Trial and failure of, contraindication, or adverse reaction to methotrexate
- Trial and failure of **TWO (2)** of the **PREFERRED** drugs below:

Humira®

Enbrel®

Medication being provided by Specialty Pharmacy - PropriumRx

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****