

# SENTARA HEALTH PLANS

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

**Drug Requested:** PiaSky<sup>®</sup> (crovalimab-akkz) SQ (Pharmacy)  
Paroxysmal Nocturnal Hemoglobinuria (PNH)

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Name/Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight (if applicable): \_\_\_\_\_ Date weight obtained: \_\_\_\_\_

**Recommended Dosing:** Piasky<sup>®</sup> 340 mg/2 mL solution in single-dose vials for infusion

- **Weight  $\geq$  40 kg to <100kg:**
  - **Loading Dose:** 1,000 mg IV on day 1 followed by 340 mg SQ on days 2, 8, 15, 22
  - **Maintenance Dose:** 680 mg SQ on day 29 and every 4 weeks thereafter
- **Weight  $\geq$  100 kg:**
  - **Loading Dose:** 1,500 mg IV on day 1 followed by 340 mg SQ on days 2, 8, 15, 22
  - **Maintenance Dose:** 1020 mg SQ on day 29 and every 4 weeks thereafter

**Quantity Limit:** 6 mL (3 vials) per 28 days

(Continued on next page)

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Initial Authorization: 6 months**

- ☐ Prescribed by or in consultation with a hematologist or nephrologist
- ☐ Prescriber is enrolled in the PiaSky Risk Evaluation and Mitigation Strategy (PIASKY REMS) program
- ☐ Member must be 13 years of age or older
- ☐ Member body weight is at least 40 kg
- ☐ Member must have a diagnosis of Paroxysmal Nocturnal Hemoglobinuria (PNH) confirmed by detection of PNH clones of at least 10% by flow cytometry testing (**must submit labs**)
- ☐ Flow cytometry pathology report must demonstrate at least 2 different glycosylphosphatidylinositol (GPI) protein deficiencies (e.g., CD55, CD59, etc.) within 2 different cell lines from granulocytes, monocytes, erythrocytes (**must submit labs**)
- ☐ Member must have **ONE** of the following indications for therapy (**must submit chart notes and labs**):
  - ☐ Member is transfusion dependent as defined by having a transfusion within the last 12 months and **ONE** of the following:
    - ☐ Member's hemoglobin is less than or equal to 7 g/dL
    - ☐ Member has symptoms of anemia, and the hemoglobin is less than or equal to 9 g/dL
  - ☐ Member has high lactate dehydrogenase (LDH) level (defined as  $\geq 1.5$  times the upper limit of the normal range with clinical symptoms)
  - ☐ Presence of a thrombotic event (e.g., DVT, PE)
  - ☐ Presence of organ damage secondary to chronic hemolysis
  - ☐ Member is pregnant and potential benefit outweighs potential fetal risk
- ☐ Member does **NOT** have evidence of an active infection caused by encapsulated bacteria (e.g., *Streptococcus pneumoniae*, *Neisseria meningitidis* or *Haemophilus influenzae*)
- ☐ Member must be vaccinated against encapsulated bacteria (*Streptococcus pneumoniae*, *Neisseria meningitidis*, and *Haemophilus influenzae type B*) **at least two weeks prior** to initiation of PiaSky<sup>®</sup> therapy and revaccinated according to current medical guidelines for vaccine use
- ☐ Member has **NOT** received a vaccination **at least two weeks prior** to the initiation of therapy with PiaSky<sup>®</sup> and documented the risks of delaying PiaSky<sup>®</sup> therapy outweigh the risks of developing an infection
- ☐ Medication will **NOT** be prescribed concurrently with another FDA approved product prescribed for treatment of PNH (e.g., Bkempv<sup>™</sup>, Epysqli<sup>™</sup>, Soliris<sup>®</sup>, Ultomiris<sup>®</sup>, Empaveli<sup>®</sup>, Fabhalta<sup>®</sup>, Voydeya<sup>™</sup>)

(Continued on next page)

- ☐ PiaSky® will be used as switch therapy **AND** member meets **ALL** the following:
  - ☐ Member is currently receiving treatment with eculizumab or ravulizumab and has shown a beneficial disease response and absence of unacceptable toxicity while on therapy
  - ☐ Provider attests administration of the IV loading dose will occur at the time of the next scheduled C5 inhibitor dose

**Reauthorization: 6 months.** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied

- ☐ Member continues to meet all initial authorization criteria
- ☐ Provider attests to an absence of unacceptable toxicity from the drug (e.g., serious meningococcal infections (septicemia and/or meningitis), infusion reactions, serious infections)
- ☐ Member has experienced positive disease response indicated by at least **ONE** of the following (**check all that apply; results must be submitted to document improvement**):
  - ☐ Documentation of a recent (within 3 months) LDH level that shows a reduction from baseline
  - ☐ Documentation that the member has stabilized hemoglobin levels as supported by **ONE** of the following:
    - ☐ Member had a reduction in number of transfusions **OR** units of packed red cells transfused from baseline
    - ☐ Member maintained a hemoglobin concentration above 7 g/dL **OR** maintained a hemoglobin concentration above 9 g/dL if member had a baseline hemoglobin level above 7 g/dL but below 9 g/dL
    - ☐ Member had a reduction in thrombotic events (e.g., DVT, PE)

**EXCLUSIONS.** Therapy will **NOT** be approved if member has history of any of the following:

- Unresolved meningococcal disease
- Any systemic bacterial or significant infections that have not been treated with appropriate antibiotics

**Medication being provided by Specialty Pharmacy – Proprium Rx**

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****