

SENTARA HEALTH PLAN

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-844-668-1550**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization can be delayed.**

For Medicare Members: Medicare Coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx>. Additional indications may be covered at the discretion of the health plan.

Botulinum Toxin Injections[®], Type A (Medical)

Drug Requested: (check applicable drug below)

☐ **Botox[®]** (onabotulinumtoxinA) (J0585)

☐ **Xeomin[®]** (incobotulinumtoxinA) (J0588)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code: _____

Weight: _____ Date: _____

☐ Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

- **Max quantity limits:** 400 units in a 3-month period
- **Cosmetic indications are EXCLUDED**

NOTE: In treating adult patients for one or more indications, the maximum cumulative dose should not exceed 400 units, in a 3-month interval. In pediatric patients, the total dose should not exceed the lower of 10 units/kg body weight or 340 units, in a 3-month interval.

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

☐ **Achalasia, Primary idiopathic esophageal**

- ☐ Member failed or had a clinically significant adverse reaction to conventional therapy (nitrates or calcium channel blockers)

OR

- ☐ Member ineligible for surgical treatment due to advance age or multiple co-morbidities (poor surgical risk)

OR

- ☐ Member is at high risk of complications of pneumatic dilation or surgical myotomy

OR

- ☐ Failure of prior myotomy or dilation

OR

- ☐ Member has an epiphrenic diverticulum or hiatal hernia, both of which increase the risk of dilation induced perforation

☐ **Achalasia, Internal anal sphincter (IAS)**

- ☐ Member has not responded to treatment with laxatives

AND

- ☐ Member has not responded to or is not a candidate for anal sphincter myectomy

☐ **Anal Fissure – Chronic**

- ☐ Member failed (at least 60 days) topical nitroglycerin or topical calcium channel blocker

☐ **Blepharospasm**

☐ **Cerebral Palsy – Dynamic Contracture**

☐ **Cerebral Palsy – Spasticity** (including diplegia, hemiplegia, paraplegia, or quadriplegia)

☐ **Cervical Dystonia** (spasmodic torticollis) and **Mixed Cervical Dystonia**

☐ **CVA-related spasticity** within 1 year of onset

☐ **Drooling in Parkinson's disease**

☐ **Essential hand tremor in patients who fail oral agents**

☐ **Hand Dystonia**

☐ **Hemifacial spasm**

☐ **Hirschsprung's Disease**

(Continued on next page)

- ☐ **Laryngeal Dysphonia – Spastic**
- ☐ **Laryngeal Dystonia** (adductor spasmodic dysphonia)
- ☐ **Laryngeal Spasm**
- ☐ **Motor tics**
- ☐ **Neurogenic detrusor overactivity (NDO) and/or detrusor sphincter dyssynergia:**
 - ☐ Member has diagnosis of incontinence due to NDO or detrusor sphincter dyssynergia associated with a neurologic condition (e.g., multiple sclerosis, spinal cord injury, brain injury) that has been confirmed by urodynamic testing (**submit documentation of diagnosis**)
 - ☐ **For members aged 5-17 years:** Member has had a 30-day trial and failure of oxybutynin (oral or intravesical use) and one other oral systemic medication from the following classes: anticholinergics or beta-3 antagonists (**Must submit chart notes documenting therapy failures**)
 - ☐ **For members aged 17 years and older:** Member has had a 30-day trial and failure of two oral systemic medications from the following classes: anticholinergics or beta-3 antagonists (**Must submit chart notes documenting therapy failures**)
- ☐ **Orofacial Dyskinesia**
- ☐ **Overactive Bladder** – Members must have met all the following criteria:
 - ☐ Diagnosis of incontinence
 - ☐ Symptoms of urge incontinence, urgency, and frequency (experienced at least 3 urinary incontinence episodes and at least 24 micturitions in 3 days)
 - ☐ 8-12 week trial and failure of behavioral therapy (e.g. bladder training, control strategies, pelvic floor muscle training, fluid management)
 - ☐ Failed or inadequate response to anticholinergic therapy within the last 9 months (4-8 week trial per agent)
 - ☐ 2 anticholinergic agents and 1 β -3 adenosine receptor agonist (**requires PA**); **OR**
 - ☐ 1 anticholinergic agent and 1 alpha blocker and 1 β -3 adenosine receptor agonist (**requires PA**)
 Please indicate drugs used: _____
- ☐ **Strabismus** (injections done in lieu of coverage for surgery)
- ☐ **Synkinetic Eyelid Closure – VII Cranial Nerve**
- ☐ **Torticollis**

Medication being provided by: Please check applicable box below.

- ☐ **Physician's office** **OR** ☐ **Specialty Pharmacy – PropriumRx**

For urgent reviews: Practitioner should call Sentara Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****