

SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization may be delayed.**

Drug Requested: Kerendia[®] (finerenone)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code: _____

Weight: _____ Date: _____

Quantity Limit: 30 tablets per 30 days

Kerendia[®] Initial Dosing Recommendations:

eGFR (mL/min/1.73m ²)	Starting Dose
≥ 60	20 mg once daily
≥ 25 to < 60	10 mg once daily
< 25	Not Recommended

Dose Adjustment Based on Current Serum Potassium Concentration and Current Kerendia [®] Dose			
		10 mg once daily	20 mg once daily
Current Serum Potassium (mEq/L)	≤ 4.8	Increase the dose to 20 mg once daily*	Maintain 20 mg once daily
	> 4.8 – 5.5	Maintain 10 mg once daily	Maintain 20 mg once daily
	> 5.5	Withhold Kerendia [®] . Consider restarting at 10 mg once daily when serum potassium ≤ 5.0 mEq/L	Withhold Kerendia [®] . Restart at 10 mg once daily when serum potassium ≤ 5.0 mEq/L

(Continued on next page)

*If eGFR has decreased by more than 30% compared to previous measurement, maintain 10 mg dose.

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

- Member is 18 years of age or older and has a diagnosis of chronic kidney disease associated with type 2 diabetes
- Must submit lab test results documenting **BOTH** of the following obtained within the past 60 days
 - Members' current eGFR is $> 25\text{mL/minute}/1.73\text{ m}^2$
 - Member's current Urinary Albumin-to -Creatinine Ratio (UACR) is $\geq 30\text{ mg/g}$
- Member's current serum potassium is $\leq 5\text{ mEq/L}$ along with **BOTH** of the following (**submit current lab documentation obtained within the past 60 days**):
 - Therapy will **NOT** be initiated if serum potassium $> 5\text{ mEq/L}$
 - Initiation with increased serum potassium monitoring during the first 4 weeks will be preformed if serum potassium is $> 4.8\text{ to }5\text{ mEq/L}$
- Member is established on treatment with maximally tolerated dose of angiotensin-converting enzyme inhibitor (ACE) or angiotensin receptor blocker (ARB) medication and will continue to take along with Kerendia[®] (finerenone)
- Member tried and failed, has a contradiction, or intolerance to at least **ONE** sodium glucose transport protein 2 (SGLT2) inhibitor that is indicated for use in patients with chronic kidney disease (e.g., Farxiga[®], Jardiance[®])
- Member does **NOT** have a diagnosis of adrenal insufficiency or a diagnosis of known significant non-diabetic renal disease, including clinically relevant renal artery stenosis
- Member is **NOT** receiving simultaneous treatment with strong CYP3A4 inhibitors
- For initial therapy, member will be dosed as follows:
 - eGFR $\geq 60\text{ mL/minute}/1.73\text{ m}^2$: starting dose will be 20 mg once daily
 - eGFR $\geq 25\text{ to } < 60\text{ mL/minute}/1.73\text{ m}^2$: starting dose will be 10 mg once daily

Reauthorization: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- Member's current eGFR is $> 25\text{ mL/minute}/1.73\text{ m}^2$ (**submit current lab documentation**)
- Member's Urinary Albumin-to-creatinine Ratio (UACR) has decreased by $\geq 30\%$ from baseline level and/or been sustained at $\geq 30\%$ below baseline level since last approval (**submit current lab documentation**)

(Continued on next page)

- Member's current serum potassium level does **NOT** exceed 5.5 mEq/L (**submit current lab documentation**)
- Provider attests Kerendia® will be withheld if serum potassium is >5.5 mEq/L and will consider restarting therapy when serum potassium normalizes (≤ 5.0 mEq/L)

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****