

# SENTARA HEALTH PLANS

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

### Migraine Treatment: Non-injectable drugs

**Drug Requested:** (Select drug below)

PREFERRED	
<input type="checkbox"/> <b>Nurtec<sup>®</sup> ODT</b> (rimegepant)	<input type="checkbox"/> <b>Qulipta<sup>™</sup></b> (atogepant)
NON-PREFERRED	
<input type="checkbox"/> <b>Reyvow<sup>®</sup></b> (lasmiditan) *Member must have tried and failed preferred Nurtec <sup>®</sup> ODT and meet all PA criteria	<input type="checkbox"/> <b>Ubrelvy<sup>™</sup></b> (ubrogepant) *Member must have tried and failed preferred Nurtec <sup>®</sup> ODT and meet all PA criteria
<input type="checkbox"/> <b>Zavzpret<sup>™</sup></b> (zavegepant) 10 mg nasal spray *Member must have tried and failed preferred Nurtec <sup>®</sup> ODT and meet all PA criteria	

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_  
Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_  
Prescriber Name: \_\_\_\_\_  
Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_  
Office Contact Name: \_\_\_\_\_  
Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_  
NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Name/Form/Strength: \_\_\_\_\_  
Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_  
Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_  
Weight (if applicable): \_\_\_\_\_ Date weight obtained: \_\_\_\_\_

(Continued on next page)

**PA Migraine Treatment: Non-injectable drugs (CORE)**  
(Continued from previous page)

- Will the member be discontinuing a previously prescribed non-injectable calcitonin gene-related peptide (CGRP) antagonist medication if approved for requested medication?  
 Yes **OR**  No
- If yes, please list the medication that will be discontinued and the medication that will be initiated upon approval along with the corresponding effective date.

**Medication to be discontinued:** \_\_\_\_\_ **Effective date:** \_\_\_\_\_

**Medication to be initiated:** \_\_\_\_\_ **Effective date:** \_\_\_\_\_

**Recommended Dosing:**

<b>Drug</b>	<b>Dose</b>	<b>Quantity Limit</b>
<b>Nurtec® ODT</b>	<ul style="list-style-type: none"> <li>• Acute Migraine: 75 mg orally as a single dose; Maximum: 75 mg/24 hours</li> <li>• Preventive Migraine (Episodic): 75 mg orally every other day</li> <li>• The safety of treating &gt; 18 doses in a 30-day period has not been established</li> </ul>	<ul style="list-style-type: none"> <li>• Acute Migraine: 8 tablets per 30 days</li> <li>• Preventive Migraine: 16 tablets per 30 days</li> </ul>
<b>Ubrelvy®</b>	<ul style="list-style-type: none"> <li>• Acute Migraine: Initial: 50 to 100 mg as a single dose; May repeat once based on response and tolerability after ≥ 2 hours</li> <li>• Maximum dose: 200 mg per 24 hours</li> <li>• The safety of treating &gt; 8 migraines/month has not been established</li> </ul>	<ul style="list-style-type: none"> <li>• 10 tablets per 30 days</li> </ul>
<b>Reyvow®</b>	<ul style="list-style-type: none"> <li>• Acute Migraine: Initial: 50 to 100 mg as a single dose; maximum of 1 dose in 24 hours</li> <li>• The safety of treating &gt; 4 migraines/month has not been established</li> </ul>	<ul style="list-style-type: none"> <li>• 4 tablets per 30 days</li> </ul>
<b>Qulipta®</b>	<ul style="list-style-type: none"> <li>• Preventive Migraine (Chronic &amp; Episodic): 10 mg, 30 mg or 60 mg orally once daily</li> <li>• Maximum dose: 60 mg/day</li> </ul>	<ul style="list-style-type: none"> <li>• 30 tablets per 30 days</li> </ul>
<b>Zavzpret™</b>	<ul style="list-style-type: none"> <li>• Acute Migraine: 10 mg given as a single spray in one nostril, as needed; Maximum: 10 mg/24 hours</li> <li>• The safety of treating more than 8 migraines in a 30-day period has not been established</li> </ul>	<ul style="list-style-type: none"> <li>• 1 carton (6 sprays) per 30 days</li> </ul>

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Authorization Criteria:**

- Member must be 18 years of age or older
- Provider has attested to all clinical criteria for **ONE** of the applicable diagnoses below

(Continued on next page)

**❑ Diagnosis: Acute Migraine**

- ❑ Member must meet **ONE** of the following:
  - ❑ Member has failed (defined as  $\geq 2$  attacks) at least **TWO** triptans (such as sumatriptan, rizatriptan) supported by the American Headache Society/American Academy of Neurology treatment guidelines, taken at maximum recommended doses
  - ❑ Provider attests member has an intolerance to triptan therapy
  - ❑ Member has at least **ONE** of the following cardiovascular or non-cardiovascular contraindications to triptan therapy:
    - ❑ Ischemic coronary artery disease (CAD) including angina pectoris, history of myocardial infarction, documented silent ischemia, coronary artery vasospasm (including Prinzmetal's angina)
    - ❑ History of stroke or transient ischemic attack (TIA)
    - ❑ Peripheral vascular disease
    - ❑ Ischemic bowel disease
    - ❑ Uncontrolled hypertension
- ❑ For Reyvow<sup>®</sup>, Ubrelvy<sup>®</sup> and Zavzpret<sup>™</sup> requests: Member must have trial and failure of Nurtec<sup>®</sup> ODT (**verified through pharmacy paid claims or chart notes**)
- ❑ For Nurtec<sup>®</sup> ODT or Ubrelvy<sup>®</sup> provider must attest to **ALL** the following:
  - ❑ Member does **NOT** have a CrCl < 15 mL/minute
  - ❑ Member is **NOT** currently using a strong CYP3A4 inhibitor (such as ketoconazole, itraconazole, or clarithromycin) or a strong CYP3A inducer (such as phenobarbital, phenytoin, or rifampin)
  - ❑ Member does **NOT** have severe hepatic impairment (Child-Pugh C)
- ❑ For Reyvow<sup>®</sup> requests: provider attests member has agreed to **NOT** drive or operate machinery until at least 8 hours after taking each dose
- ❑ Requested medication will **NOT** be used in combination with another oral CGRP inhibitor

**❑ Diagnosis: Preventive Migraine (Applies to Nurtec<sup>®</sup> ODT and Qulipta<sup>®</sup> only)**

- ❑ Member must have a diagnosis of Chronic or Episodic Migraine Headache defined by **BOTH** of the following:
  - ❑ Member has  $\geq 4$  migraine headaches per month
  - ❑ Member must have failed a **2-month** trial of at least one medication from **TWO** different migraine prophylactic classes supported by the American Headache Society/American Academy of Neurology treatment guidelines 2012/2015/2021/2024, Level A and B evidence; ICSI 2013, high quality evidence:
    - ❑ Anticonvulsants (divalproex, valproate, topiramate)
    - ❑ Beta blockers (atenolol, metoprolol, nadolol, propranolol, timolol)
    - ❑ Antidepressants (amitriptyline, venlafaxine)
    - ❑ Angiotensin II Receptor Blocker (candesartan) **\*requires prior authorization\***
    - ❑ CGRP inhibitors (Aimovig<sup>®</sup>, Emgality<sup>®</sup>, Ajovy<sup>®</sup>, Vyepti<sup>®</sup>)

(Continued on next page)

- ❑ Provider must attest to **ALL** the following:
  - ❑ Member does **NOT** have a CrCl < 15 mL/minute for Nurtec<sup>®</sup> ODT
  - ❑ Member is **NOT** currently using a strong CYP3A4 inhibitor (such as ketoconazole, itraconazole, or clarithromycin) or a strong CYP3A inducer (such as phenobarbital, phenytoin, or rifampin)
  - ❑ Member does **NOT** have severe hepatic impairment (Child-Pugh C)
  - ❑ Requested medication will **NOT** be used in combination with another oral CGRP inhibitor
  - ❑ Nurtec<sup>®</sup> ODT and Qulipta<sup>®</sup> will **NOT** be used in combination with Aimovig<sup>®</sup>, Emgality<sup>®</sup>, Ajoovy<sup>®</sup> or Vyepiti<sup>®</sup>
- ❑ **Requests for concurrent use of Calcitonin Gene-Related Peptide (CGRP) inhibitors with Botox<sup>®</sup> (onabotulinumtoxinA) for migraine headache prevention (if applicable):** Member must meet **ALL** the following criteria (**verified by chart notes and/or pharmacy paid claims**):
  - ❑ Member must have a diagnosis of Chronic or Episodic Migraine Headache and is continuing to experience ≥ 4 migraine headache days per month after receiving therapy with **ALL** the following criteria:
    - ❑ Member must have failed a **2-month** trial of at least one medication from **TWO** different migraine prophylactic classes supported by the American Headache Society/American Academy of Neurology treatment guidelines 2012/2015/2021/2024, Level A and B evidence: ICSI 2013, high quality evidence:
      - ❑ Anticonvulsants (divalproex, valproate, topiramate)
      - ❑ Beta blockers (atenolol, metoprolol, nadolol, propranolol, timolol)
      - ❑ Antidepressants (amitriptyline, venlafaxine)
      - ❑ Angiotensin II receptor blocker (candesartan) **\*requires prior authorization\***
    - ❑ Member must meet **ONE** of the following:
      - ❑ Member has had an inadequate response to a **2-month** trial with an injectable CGRP inhibitor (e.g., Aimovig<sup>®</sup>, Ajoovy<sup>®</sup>, Emgality<sup>®</sup>) or an oral CGRP inhibitor indicated for migraine prevention (e.g., Nurtec<sup>®</sup> ODT, Qulipta<sup>™</sup>) **\*requires prior authorization\***
      - ❑ Member has had an inadequate response to a **6-month** trial (2 injection cycles) of Botox<sup>®</sup> (onabotulinumtoxinA) **\*requires prior authorization\***

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****