

OPTIMA HEALTH MEDICAID

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to **1-804-799-5118**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization will be delayed.**

Drug Requested: Orencia® (abatacept) (J-0129) (IV INFUSION ONLY) (Medical)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Optima #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code: _____

Weight: _____ Date: _____

DIAGNOSIS	Recommended Quantity
<input type="checkbox"/> Moderately to severely active Rheumatoid Arthritis (RA)	INTRAVENOUS <ul style="list-style-type: none">Weight <60kg: Two vials per 28 daysWeight 60-100kg: Three vials per 28 daysWeight >100kg: Four vials per 28 days
<input type="checkbox"/> Psoriatic arthritis (PsA)	INTRAVENOUS <ul style="list-style-type: none">Weight <60kg: Six 250mg vials initial 28 days Two vials per 28 days after inductionWeight 60-100kg: Nine vials in initial 28 days Three vials per 28 days after inductionWeight >100kg: Twelve vials initial 28 days Four vials per 28 days after induction

(Continued on next page)

DIAGNOSIS	Recommended Quantity
<input type="checkbox"/> Juvenile Idiopathic Arthritis (JIA) in members 2 years and older	INTRAVENOUS <ul style="list-style-type: none"> • ≥ 6 y/o (subcutaneous formulation indicated for 2 years and older) • Weight <75kg: 10mg/kg every 28 days (3 vial) • Weight >75kg: Follow adult Rheumatoid Arthritis dosing above (not to exceed max dose 1000mg)
<input type="checkbox"/> Prophylaxis of acute graft versus host disease (aGVHD)	INTRAVENOUS <ul style="list-style-type: none"> • Patients 2 to less than 6 years old, administer a 15 mg/kg dose as a 60-minute infusion on the day before transplantation, followed by a 12 mg/kg dose as a 60-minute infusion on Day 5, 14, and 28 after transplant • Patients 6 years and older, 10 mg/kg dose (maximum dose 1,000 mg) as a 60-minute infusion day before transplantation, followed by a dose on Day 5, 14, and 28 after transplant

- Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Diagnosis: Moderate to severe Active Rheumatoid Arthritis (RA)

- Prescriber is a **Rheumatologist**

AND

- Member has been diagnosed with **moderate to severe rheumatoid arthritis**

AND

- Trial and failure of, contraindication, or adverse reaction to methotrexate

AND

- Trial and failure of at least **ONE (1)** other **DMARD therapy** including, but not limited to: (**check each tried**)

<input type="checkbox"/> auranofin	<input type="checkbox"/> sulfasalazine
<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> other: _____

AND

- Member has tried and failed **TWO (2)** of the following biologics:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

Diagnosis: Juvenile Idiopathic Arthritis (JIA)

- Prescriber is a **Rheumatologist**

AND

- Member has been diagnosed with **moderate to severe active Juvenile Idiopathic Arthritis (JIA)**

AND

- Trial and failure of, contraindication, or adverse reaction to methotrexate

AND

- Trial and failure of at least **ONE (1)** other **DMARD therapy** including, but not limited to: (**check each tried**):

<input type="checkbox"/> auranofin	<input type="checkbox"/> sulfasalazine
<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> other: _____

AND

- Patient has tried and failed **TWO (2)** of the biologics below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®
----------------------------------	----------------------------------

Diagnosis: Active Psoriatic Arthritis (PsA)

- Prescriber is a **Rheumatologist**

AND

- Member has been diagnosed with **moderate to severe active Psoriatic Arthritis (PsA)**

AND

- Trial and failure of, contraindication, or adverse reaction to methotrexate

AND

(Continued on next page)

- Trial and failure of at least **ONE (1)** other **DMARD therapy** including, but not limited to: (**check each tried**):

<input type="checkbox"/> auranofin	<input type="checkbox"/> sulfasalazine
<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> other: _____

AND

- Member has tried and failed **TWO (2)** of the following biologics:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

Diagnosis: Prophylaxis of acute graft versus host disease (aGVHD)

- Member has a diagnosis of prophylaxis of acute graft versus host disease (aGVHD)

AND

- Member is 2 years of age and older undergoing hematopoietic stem cell transplantation (HSCT) from a matched or 1 allele-mismatched unrelated donor

AND

- The medication is used in combination with a calcineurin inhibitor and methotrexate

Medication being provided by (check applicable box(es) below):

- Location/site of drug administration: _____
NPI or DEA # of administering location: _____

OR

- Physician's office **OR** Specialty Pharmacy – PropriumRx

For urgent reviews: Practitioner should call Optima Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Optima’s definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member’s ability to regain maximum function.

****Use of samples to initiate therapy does not meet step-edit/preauthorization criteria.****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****