## SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

NON-PREFERRED

☐ Acthar® Gel (repository corticotropin) 80 USP

☐ Acthar® Gel (repository corticotropin) 40 USP Units/0.5 mL single-dose prefilled SelfJect

Units/mL 5 mL multi-dose vial

**<u>Drug Requested</u>**: Repository Corticotropin Medications - Nephrotic Syndrome (NS)

**PREFERRED** 

□ Purified Cortrophin<sup>™</sup> Gel

(repository corticotropin)

adults only.

	injector			
	☐ Acthar® Gel (repository corticotropin) 80 USP			
	Units/mL single-dose prefilled SelfJect injector			
	*Member must have tried and failed preferred			
	Purified Cortrophin <sup>™</sup> Gel and meet all applicable PA criteria below			
	circin below			
MEMBER & PRESCRIBER INFOR	RMATION: Authorization may be delayed if incomplete.			
Member Name:				
Member Sentara #:	ra #: Date of Birth:			
Prescriber Name:				
Prescriber Signature:	er Signature: Date:			
Office Contact Name:				
Phone Number:	ne Number: Fax Number:			
NPI #:				
DRUG INFORMATION: Authorization	on may be delayed if incomplete.			
Drug Name/Form/Strength:				
Dosing Schedule:	Length of Therapy:			
Diagnosis:	ICD Code, if applicable:			
Weight (if applicable):	t (if applicable): Date weight obtained:			

(Continued on next page)

Acthar Gel single-dose pre-filled SelfJect injector is for subcutaneous administration by

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied. Check box below for the Diagnosis that applies.

ш	Member MUSI nave a docum	nentea alagnosis o	i Nepnro	tic S	Syndrome with <u>ONE</u> of the following:		
	☐ Focal Segmental Glomerulo	osclerosis (FSGS)	OR		Membranous Nephropathy (MPGN)		
	☐ Minimal Change Disease _						
	The following MUST be noted	d:					
	1. Baseline current kg:		_				
					rine protein/creatinine ratio with mg/mg nephrotic range proteinuria)		
	Member <u>MUST</u> have tried and failed both a corticosteroid <u>AND</u> a calcineurin inhibitor (CNI) taken concurrently within the year of request. Failure is defined as no change or an increase from baseline proteinuria levels after 90 consecutive days of concomitant corticosteroid and calcineurin therapy trial. Approval will be based on proteinuria increase from baseline after 90 consecutive days of concomitant corticosteroids and calcineurin inhibitor therapy.						
	3. 90 days post concurrent cor	ticosteroid and calc	ineurin ir	nhibi	itor trial, urine protein/creatinine ratio;		
	Date:	;			(mg/mg nephrotic range proteinuria)		
	Member MUST have had trial and failure of high dose corticosteroid for a minimum of 90 consecutive days within last 12 months. Note name of therapy tried and dose (must note therapy tried and trial MUST be noted in pharmacy or medical claims):						
	□ 1 mg/kg (max 80 mg)	OR	□ 2mg	g/kg	alternate day (max 120 mg)		
	AND						
	Member <u>MUST</u> have had concurrent trial and failure of calcineurin inhibitor for a minimum of 90 days consecutive days within last 12 months ( <u>must</u> note therapy tried and trial <u>MUST</u> be noted in pharmacy paid claims):						
	□ Cyclosporine	□ Tacrolimus			□ Cyclophosphamide		
	OR						
	If member has a relative <u>cont</u> uncontrolled diabetes BS > 20						
	Member has had trial and failur pharmacy paid claims):	re of calcineurin inh	ibitor on	ly (tl	herapy tried <u>MUST</u> be noted in		
	☐ Cyclosporine: mg mg/m²/day in 2 divided dos	es; adjust doses bas			doses for at least 12 months <b>OR</b> 150 levels {(pediatrics): 80 to 100 ng/mL}		
	☐ Tacrolimus:						
	☐ Cyclophosphamide:	mg					

(Continued on next page)

PA Repository Corticotropin-NS (Medicaid)
(Continued from previous page)

□ Pro	Progress notes MUST be submitted with documentation of ALL THREE (3) of the following labs:						
	Proteinuria	□ Serum Albumin	□ Cyclosporine levels				
Do	Dose Regimen: Anticipated Length of therapy:						
NOTE: Approval will be for a period of 6 weeks with a follow up Proteinuria lab required to be submitted. IF additional therapy is needed; the prescribing physician will need to submit a second request for continuation of therapy.							
Medica	tion being provided by	Specialty Pharmacy – Propriu	m Rx				

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. \*\*

\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\*