

SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Sirturo[®] (bedaquiline)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

Quantity Limit: 68 tablets for the first 28 days of treatment and then 24 tablets per 28 days for the next 20 weeks.

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- Member is \geq 18 years old **AND** enrolled in a DOT (**Directly Observed Therapy**) Program

AND

- Diagnosis of Pulmonary Multi-Drug Resistant Tuberculosis (MDR-TB)

(Please send Sputum culture for mycobacterium. Cultures provide precise species identification, drug sensitivity testing, and genotyping for epidemiologic purposes.)

OR

- Charts/Labs **MUST** be provided to document an M. tuberculosis isolate that is resistant to at least isoniazid, rifampin, and possibly additional agents

(Continued on next page)

AND

- Does the patient have diagnosis of latent or extra-pulmonary tuberculosis? Yes **OR** No
(Not indicated for treatment of latent, extra-pulmonary or drug sensitive TB)

AND

- Sirturo[®] to be used in combination with three other drugs? Yes **OR** No

AND

- Please mark all drugs the member is using in combination with Sirturo[®]: **(at least 3 must be marked)**

<u>Antibiotic Drugs</u> (check each that the member is using in combination with Sirturo [™] ; at least three (3) must be marked.)			
<input type="checkbox"/> Amikacin	<input type="checkbox"/> Capreomycin	<input type="checkbox"/> Clofazimine	<input type="checkbox"/> Cycloserine
<input type="checkbox"/> Dapsone	<input type="checkbox"/> Ethambutol	<input type="checkbox"/> Ethionamide	<input type="checkbox"/> Isoniazid
<input type="checkbox"/> Kanamycin	<input type="checkbox"/> Linezolid	<input type="checkbox"/> Ofloxacin	<input type="checkbox"/> Pyrazinamide
<input type="checkbox"/> Rifampicin	<input type="checkbox"/> Terizidone	<input type="checkbox"/> Streptomycin	<input type="checkbox"/> 4-Aminosalicylic acid

Not all drugs may be covered under every Plan

If a drug is non-formulary on a Plan, documentation of medical necessity will be required.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****