SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information <u>(including phone and fax #s)</u> on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Drug Requested: Stelara® SQ (ustekinumab) For PsA & PsO (Pharmacy) (Preferred)

r Name:	Date of Birth: Date: Fax Number:					
r Sentara #: ber Name: ber Signature: Contact Name: Number: R NPI #:	Date of Birth: Date: Fax Number:					
ber Signature:	Date: Fax Number:					
ber Signature:	Date: Fax Number:					
Number:R NPI #:	Fax Number:					
R NPI #:						
CINFORMATION: Authorization may be dele						
JIM OMMATION. Aumonzanon may be dela	yed if incomplete.					
orm/Strength:						
Schedule:	Length of Therapy:					
Diagnosis: ICD Code, if applicable						
: 1	Date:					
ICAL CRITERIA: Check below all that apply teach line checked, all documentation, including label or request may be denied. Check the diagnosis be	results, diagnostics, and/or chart notes, must be					
agnosis: Active Psoriatic Arthritis						
SubQ: 45 mg at 0 and 4 weeks; then every 12 weeks, may be administered alone or in combination with						
tent psoriatic arthritis and moderate-to-severe pla aintenance: 90 mg at 0 and 4 weeks; then every 12 v	1 1					
\square Member is ≥ 6 years old and has a diagnosis of active psoriatic arthritis						
 Prescribed by or in consultation with a Rheumatologist 						

(Continued on next page)

	Member tried and failed at least one DMARD for at least three (3) months (check each tried below):						
	□ methotrexate		azathioprine		hydroxychloroquine		
	□ sulfasalazine		leflunomide		auranofin		
	Other:						
□ Diagnosis: Moderate-to-Severe Plaque Psoriasis							
Dosing: SubQ: ≤100 kg: 45 mg at 0 and 4 weeks; then every 12 weeks thereafter. >100 kg: 90 mg at 0 and 4 weeks; then every 12 weeks thereafter. NOTE: Doses of 45 mg given to patient's >100 kg were also efficacious; however, 90 mg is the recommended dose in these patients due to greater efficacy							
\square Member is ≥ 6 years old and has a diagnosis of moderate-to-severe plaque psoriasis							
□ F	☐ Prescribed by or in consultation with a Dermatologist						
	☐ Member tried and failed at least ONE (1) of either Phototherapy or Alternative Systemic Therapy for at least three (3) months (check each tried below):						
	□ Phototherapy:		□ Alternative System	mic	Therapy:		
	UV Light Therapy		□ Oral Medicati	ons			
	□ NB UV-B		□ acitretin				
	□ PUVA		□ methotrexa				
			□ cyclosporin	ne			
Medication being provided by a Specialty Pharmacy - PropriumRx							

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *