SENTARA HEALTH PLANS

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-844-668-1550</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not</u> complete, correct, or legible, authorization can be delayed.

For Medicare Members: Medicare Coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: <u>https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx</u>. Additional indications may be covered at the discretion of the health plan.

Gonadotropin-releasing Hormone Agonists (GnRH) (Medical)

Drug Requested: (Select drug below)

Preferred Drugs				
Camcevi [®] (leuprolide mesylate) 42 mg (6-		Eligard Depot [®] (leuprolide acetate)		
month) (J1952)	Su	spension (J9217):		
		7.5 mg (1-month)		
		□ 22.5 mg (3-month)		
		□ 30 mg (4-month)		
 		□ 45 mg (6-month)		
Leuprolide acetate 5 mg/mL SubQ		Lupron Depot [®] (leuprolide acetate)		
Solution (J9218)		Suspension Kit (J1950):		
		□ 3.75 mg (1-month)		
		7.5 mg (Pediatric 1-month)		
		□ 11.25 mg (Pediatric 1-month)		
		□ 11.25 mg (Pediatric 3-month)		
		□ 15 mg (Pediatric 1-month)		
 <u>^</u>		30 mg (Pediatric 3-month)		
Lupron Depot [®] (leuprolide acetate)		Trelstar [®] (triptorelin pamoate) Suspension		
Suspension Kit (J9217):		(J3315):		
□ 7.5 mg (1-month)		□ 3.75 mg (1-month)		
□ 11.25 mg (3-month)		□ 11.25 mg (3-month)		
22.5 mg (3-month) [syringe kit]		□ 22.5 mg (6-month)		
□ 22.5 mg (3-month) [vial]				
□ 30 mg (4-month)				
 □ 45 mg (6-month)				
Zoladex [®] (goserelin) (J9202):				
3.6 mg (1-month)				
 □ 10.8 mg (3-month)				
Non-Prefer	red	Drugs		
Fensolvi [®] (leuprolide acetate) 45mg		Supprelin [®] LA (histrelin acetate) 50 mg		
(6- month) (J1951)		(12- month) (J9226)		
Triptodur [®] (triptorelin) 22.5 mg (6-month)				
(J3316)				

MEMBER & PRESCRIBER IN	FORMATION: Authorization may be delayed if incomplete.		
Member Name:			
Member Sentara #:			
Prescriber Name:			
	Date:		
Office Contact Name:			
Phone Number:	ber: Fax Number:		
DEA OR NPI #:			
DRUG INFORMATION: Author	ization may be delayed if incomplete.		
Drug Form/Strength:			
Dosing Schedule:	Length of Therapy:		
iagnosis: ICD Code, if applicable:			
Weight:	Date:		

□ Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

A. Length of Authorization:

- Endometriosis: Coverage will be provided for 6 months and is eligible for conditional renewal
- Uterine leiomyomata (fibroids): Coverage will be provided for 6 months and is eligible for conditional renewal
- Gender Dysphoria: Approval will be indefinite per length of requested treatment, and renewal will not be required
- All other indications: Coverage will be provided for 12 months and may be renewed

B. Quantity Limits:

Drug Name	Strength	Quantity	Day Supply
Camcevi	42 mg	1 injection	180 days
Leuprolide acetate SubQ Solution	5 mg/mL (1 mg/0.2 mL multi-dose vial)	2 vials	28 days
Lupron Depot 1-month	3.75 mg, 7.5 mg	1 injection	28 days
Lupron Depot 3-month	11.25 mg, 22.5 mg	1 injection/vial	84 days
Lupron Depot 4-month	30 mg	1 injection	112 days

Drug Name	Strength	Quantity	Day Supply
Lupron Depot 6-month	45 mg	1 injection	168 days
Lupron Depot-Ped 1-month	7.5 mg, 11.25 mg, 15 mg	1 injection	28 days
Lupron Depot-Ped 3-month	11.25 mg, 30 mg	1 injection	84 days
Eligard 1-month	7.5 mg	1 injection	28 days
Eligard 3-month	22.5 mg	1 injection	84 days
Eligard 4-month	30 mg	1 injection	112 days
Eligard 6-month	45 mg	1 injection	168 days
Fensolvi 6-month	45 mg	1 injection	168 days
Trelstar 1-month	3.75 mg	1 injection	28 days
Trelstar 3-month	11.25 mg	1 injection	84 days
Trelstar 6-month	22.5 mg	1 injection	168 days
Triptodur 6-month	22.5 mg	1 injection	168 days
Supprelin LA	12-month	1 implant	365 days
Zoladex 1-month	3.6 mg	1 implant	28 days
Zoladex 3-month	10.8 mg	1 implant	84 days

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Diagnosis: Prostate Cancer

Initial Authorization

- □ Member is 18 years of age or older
- **□** Requesting provider is an oncologist or urologist
- □ Member has a diagnosis of advanced prostate cancer
- □ The quantity (dose) requested is in accordance with FDA-approved labeling, and if applicable or necessary, age and weight conditions are met
- □ If requesting a non-preferred drug, the member has failed <u>ONE</u> of the preferred formulations noted above

Diagnosis: Breast Cancer

Initial Authorization

- □ Member is 18 years of age or older
- □ Requesting provider is an oncologist

- □ Select <u>ONE</u> of the following:
 - □ Member is a pre- or peri-menopausal woman
 - □ Member is male with suppression of testicular steroidogenesis
- □ Member has hormone-receptor positive disease AND meets <u>ONE</u> of the following:
 - □ Medication will be used in combination with adjuvant endocrine therapy
 - □ Medication will be used combination with endocrine therapy for recurrent or metastatic disease
 - □ Medication will be used as palliative treatment for advanced disease
- □ The quantity (dose) requested is in accordance with FDA-approved labeling, and if applicable or necessary, age and weight conditions are met
- □ If requesting a non-preferred drug, the member has failed <u>ONE</u> of the preferred formulations noted above

Diagnosis: Gender Identity/Gender Dysphoria

Initial Authorization

- □ Select <u>ONE</u> of the following:
 - □ Member is 18 years of age or older and has a diagnosis of gender dysphoria
 - □ Member is less than 18 years of age and has a diagnosis of gender dysphoria. Provider please note: parental or legal guardian consent for un-emancipated members is required
- Provider attests member has the capacity to make informed treatment decisions and has assented to treatment after discussion of the potential benefits and risks
- □ Member has been assessed and diagnosed with gender dysphoria according to DSM-V criteria, by <u>ONE</u> of the following provider types:
 - □ A licensed mental health provider
 - An endocrinologist
 - □ A gender dysphoria-informed hormone prescriber, defined as a provider competent in the assessment of gender dysphoria who practices in conjunction with a multidisciplinary gender dysphoria care team
- Medication is prescribed by, or in consultation with, a licensed mental health provider, endocrinologist or other medical provider experienced in gender dysphoria hormone therapy
- Provider attests coexisting behavioral health and medical comorbidities or social problems that may interfere with diagnostic procedures or treatment are being appropriately treated and are not causing symptoms of gender dysphoria
- □ Member has experienced puberty development to at least <u>ONE</u> of the following:
 - □ Tanner stage 2 (stage 2 through 4)
 - □ Lab values for Luteinizing Hormone (LH), Follicle Stimulating Hormone (FSH), and the endogenous sex hormones consistent with at least Tanner stage 2 (must submit documentation)
- □ If requesting a non-preferred drug, the member has failed <u>ONE</u> of the preferred formulations noted above

Diagnosis: Central Precocious Puberty

Initial Authorization

- □ Member is less than 13 years of age
- Onset of secondary sexual characteristics associated with pubertal pituitary gonadotropin activation, occurring earlier than age 8 for girls and age 9 for boys (submit documentation, progress notes, medical documentation recording physical changes, Tanner staging)
- Diagnosis of central precocious puberty is confirmed by <u>ALL</u> the following:
 - □ Laboratory documentation of pubertal gonadal sex steroid level
 - □ Pubertal luteinizing hormone response simulation by native GnRH [Laboratory documentation demonstrating basal LH (>0.3 IU/L), and peak stimulated LH (>4-6 IU/L)]
 - □ X-Ray results of the estimated bone age of the non-dominant wrist and hand greater than 2 standard deviations beyond chronological age (submit laboratory and x-ray documentation)
- □ A tumor has been ruled out by lab tests such as diagnostic imaging of the brain (to rule out intracranial tumor), pelvic/testicular/adrenal ultrasound (to rule out steroid secreting tumors), and human chorionic gonadotropin levels (to rule out a chorionic gonadotropin secreting tumor)
- □ Medication will <u>NOT</u> be used in combination with growth hormone therapy
- □ The quantity (dose) requested is in accordance with FDA-approved labeling, and if applicable or necessary, age and weight conditions are met
- □ If requesting a non-preferred drug, the member has failed <u>ONE</u> of the preferred formulations noted above

Diagnosis: Gynecological Indications

Initial Authorization

- □ Member is 18 years of age or older
- □ If requesting a non-preferred drug the member has failed <u>ONE</u> of the preferred formulations notes above
- **D** Treatment is being prescribed by or in consultation with a specialist in obstetrics/gynecology

Select <u>ONE</u> of the following indications for use:

- □ **FOR UTERINE LEIOMYOMATA, FIBROIDS** (requires chart notes documenting symptomology/pelvic exam, transvaginal ultrasonography, sono-hysterography):
 - □ Member is premenopausal
 - □ Member has uterine leiomyomas (fibroids)
 - □ Member is using for the management of heavy menstrual bleeding
 - □ Member does <u>NOT</u> have any contraindications to therapy including osteoporosis, history of thrombotic or thromboembolic disorders, uncontrolled hypertension, or hepatic impairment/disease

(Continued on next page)

(Continued from previous page)

- □ Member has history of inadequate response to the following therapies, tried for at least three (3) months each (must submit chart note documentation of all therapy failures):
 - □ Oral contraceptives OR a selective progesterone receptor modulator OR intrauterine device
 - □ NSAIDs (non-steroidal anti-inflammatory drugs)
 - □ tranexamic acid 650 mg

<u>OR</u>

- □ Member has had surgery for uterine fibroids (i.e., ablation, myomectomy) and has persistent symptoms (must submit documentation of date/type of surgery or procedure)
- □ The quantity (dose) and administration frequency requested is in accordance with FDA-approved labeling, and if applicable or necessary, age and weight conditions are met
- □ FOR ENDOMETRIOSIS (requires chart notes detailing and recording treatment plan and/or symptomology of chronic pelvic pain (defined as noncyclical pain lasting 6 or more months that localizes to the anatomic pelvis, anterior abdominal wall at or below the umbilicus, the lumbosacral back, or the buttocks, and is of sufficient severity to cause functional disability or lead to medical care), amenorrhea, preoperative ablation treatment):
 - □ Member is premenopausal
 - □ Member has a diagnosis of moderate to severe pain associated with endometriosis
 - Diagnosis of endometriosis has been confirmed by direct visualization during surgery and/or histology
 - □ Member does <u>NOT</u> have any contraindications to therapy including osteoporosis, severe hepatic impairment/disease, or concomitant use of hormonal contraceptives
 - □ Member has history of inadequate response to the following therapies, tried for at least three (3) months each (must submit chart note documentation of all therapy failures):
 - □ NSAIDs (non-steroidal anti-inflammatory drugs)
 - □ Combination (estrogen/progesterone) oral contraceptive
 - □ Progestins

<u>OR</u>

- □ Member has had surgical ablation to prevent recurrence
- □ The quantity (dose) and administration frequency requested is in accordance with FDA-approved labeling, and if applicable or necessary, age and weight conditions are met

REAUTHORIZATION CRITERIA

*******NOTE: Gender identity/gender dysphoria diagnoses does NOT require reauthorization******

Diagnosis: Oncology diagnosis (Please submit chart notes and other supporting documents)

- □ Member requires continuation of therapy and is <u>NOT</u> experiencing disease progression
- Ongoing treatment is consistent with FDA-labeling or compendia support
- □ Member is <u>NOT</u> experiencing an FDA-labeled limitation of use or toxicity
- **D** The quantity (dose) requested is in accordance with FDA approved labeling

(Continued on next page)

Diagnosis: Central Precocious Puberty (Please submit chart notes and other supporting documents)

- $\Box \quad \text{Member is } \underline{\text{NOT}} \text{ over the age of } 13$
- Member has experienced disease response as indicated by lack of progression or stabilization of secondary sexual characteristics, decrease in growth velocity and bone age advancement, and improvement in final height prediction
- □ Member has experienced an absence of unacceptable toxicity from the drug (e.g., convulsions, development or worsening of psychiatric symptoms)

Diagnosis: Gynecological Indications (Please submit chart notes and other supporting documents)

□ Total duration of therapy (initial plus re-treatment for symptom recurrence) should not exceed 12 months **AND** will be used in combination with add-back therapy, unless clinically contraindicated

<u>OR</u>

□ The member's medical history and medical condition's current status requires longer treatment duration than otherwise recommended in published compendia/FDA labeling.

Please provide an explanation along with any pertinent progress notes of medical condition including recorded recurrence of symptoms, procedure/exam results:

AND

D Prescriber will order and review a bone density assessment prior to re-treatment

Medication being provided by: Please check applicable box below.

Location/site of drug administration:

NPI or DEA # of administering location: _____

<u>OR</u>

D Specialty Pharmacy – PropriumRx

For urgent reviews: Practitioner should call Sentara Health Plans Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health Plan's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. *Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *