

# SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

**Drug Requested:** Fasenra<sup>®</sup> SQ (benralizumab) (Pharmacy)

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

DEA OR NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight: \_\_\_\_\_ Date: \_\_\_\_\_

**Recommended Dosage:** 30 mg SubQ once every 4 weeks for the first 3 doses, then once every 8 weeks thereafter

\*Sentara Health Plans considers the use of concomitant therapy with Cinqair<sup>®</sup>, Dupixent<sup>®</sup>, Fasenra<sup>®</sup>, Nucala<sup>®</sup>, Tezspire<sup>™</sup> and Xolair<sup>®</sup> to be experimental and investigational. Safety and efficacy of these combinations have **NOT** been established and will **NOT** be permitted. In the event a member has an active Cinqair<sup>®</sup>, Dupixent<sup>®</sup>, Nucala<sup>®</sup>, Tezspire<sup>™</sup> or Xolair<sup>®</sup> authorization on file, all subsequent requests for Fasenra<sup>®</sup> will **NOT** be approved.

Medication will be (select **ONE** of the following):

- ☐ Self-Administered (pharmacy benefit)
- ☐ Administered by Provider (medical benefit)

(Continued on next page)

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Initial Authorization: 12 months**

- ☐ Prescribed by or in consultation with an allergist, immunologist or pulmonologist
- ☐ Member is 12 years of age or older
- ☐ Has the member been approved for Fasenra<sup>®</sup> previously through the Sentara medical department?  
☐ Yes   ☐ No
- ☐ Member has been diagnosed with severe eosinophilic phenotype defined by a baseline (pre-Fasenra<sup>®</sup>) peripheral blood eosinophil level  $\geq 150$  cells/microliter at the initiation of treatment
- ☐ Member is currently being treated with **ONE** of the following unless there is a contraindication or intolerance to these medications and must be compliant on therapy **for at least 90 consecutive days** within a year of request:
  - ☐ High-dose inhaled corticosteroid (ICS) (e.g., greater than 500 mcg fluticasone propionate equivalent/day) **AND** an additional asthma controller medication (e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline)
  - ☐ One maximally dosed combination ICS/LABA product (e.g., Advair<sup>®</sup> (fluticasone propionate/salmeterol), Dulera<sup>®</sup> (mometasone/formoterol), Symbicort<sup>®</sup> (budesonide/formoterol))
- ☐ Member has experienced **ONE** of the following (check box that applies):
  - ☐ More than > 2 exacerbations requiring additional medical treatment (e.g., an increase in oral corticosteroid dose, emergency department, urgent care visits or hospitalizations) within the past 12 months
  - ☐ Any prior intubation for an asthma exacerbation
- ☐ Member has a baseline forced expiratory volume (FEV1) < 80% predicted normal (< 90% for members 12-17 years old) submitted within year of request
- ☐ Provider must submit member blood eosinophil count after a trial and failure of at least 90 consecutive days of therapy with high dose inhaled corticosteroids **AND** long-acting inhaled beta-2 agonist. A failure of these medications is defined as a blood count > 150 cells/microliter (**submit labs collected within the past 12 months**)

Eosinophil count: \_\_\_\_\_ Date: \_\_\_\_\_

(Continued on next page)

**Reauthorization: 12 months.** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ☐ Member has experienced a sustained positive clinical response to Fasenra® therapy as demonstrated by at least **ONE** of the following (**check all that apply; chart notes must be submitted**):
  - ☐ Increase in percent predicted Forced Expiratory Volume (FEV1) from baseline (pre-treatment)
  - ☐ Reduction in the dose of inhaled corticosteroids required to control asthma
  - ☐ Reduction in the use of oral corticosteroids to treat/prevent exacerbation
  - ☐ Reduction in asthma symptoms such as chest tightness, coughing, shortness of breath or nocturnal awakenings
- ☐ Member is currently being treated with **ONE** of the following unless there is a contraindication or intolerance to these medications:
  - ☐ High-dose inhaled corticosteroid (ICS) (e.g., greater than 500 mcg fluticasone propionate equivalent/day) **AND** an additional asthma controller medication (e.g., leukotriene receptor antagonist, long-acting beta-2 agonist (LABA), theophylline)
  - ☐ One maximally dosed combination ICS/LABA product (e.g., Advair® (fluticasone propionate/salmeterol), Dulera® (mometasone/formoterol), Symbicort® (budesonide/formoterol))

**Medication being provided by a Specialty Pharmacy - PropriumRx**

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****