SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name</u> (<u>preprinted stamps not valid</u>) on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete</u>, correct, or legible, the authorization process can be delayed.

The Sentara Health Plans Oncology Program is administered by OncoHealth

❖ For any oncology indications, the most efficient way to submit a prior authorization request is through the OncoHealth OneUM Provider Portal at https://oneum.oncohealth.us. Fax to 1-800-264-6128.
OncoHealth can also be contacted at Phone: 1-888-916-2616

ANTIEMETIC/ANTIVERTIGO DRUGS

Drug Requested: (Check below the drug that applies)

PREFERRED MEDICATIONS (***PREFERRED Dronabinol and Diclegis Require Prior Authorization)										
	Diclegis ® (doxylamine succinate/vitamin B6) ***	k	ondansetro 8mg)/tab/s		ODT (4mg,	- 1	neclizine (OTC, Rx)			
	metoclopramide (tab/sol	ln/vial)	□ Phenadoz [©] over 2 yea		pp (AG) (members f age)	<u> </u>	Prochlorperazine tab			
	promethazine (AG) (merover 2 years of age)	mbers dronabinol capsule***								
All Non-Preferred Medications Require Prior Authorization										
	Akynzeo®	□ Aloxi®			Antivert®		Aponvie TM			
	aprepitant capsule/pack	□ Ba	rhemsys®		Bonjesta [®]		Cesamet [®]			
	Cinvanti [™]		mpazine [®] pp/tab		Compro®		dimenhydrinate tab, vial			
	Emend® Bi Pak	□ Em	nend® susp		Emend [®] Tri-fold pack		Focinvez TM			
	Fosaprepitant vial	□ gra	nisetron		Kytril [®]		Marinol [®]			
	metoclopramide ODT		lansetron ng ODT		palonosetron (generic Aloxi®)		prochlorperazine supp, vial			
	promethazine 50mg supp, vial, ampule	□ Re	glan®		Sancuso® patch		scopolamine (generic Transderm-Scop®)			
	Sustol [®]	□ Syı	ndros™		Transderm-Scop®		trimethobenzamide			
	Varubi [®]	□ Vis	taril [®]		Zofran [®] ODT/soln/tab					

(Continued on next page)

Memb	er Name:								
Memb	er Sentara #:								
Prescri	iber Name:								
Prescri	iber Signature:								
Office	Contact Name:								
NPI #:									
	G INFORMATION: Authorization may be delayed if incomp								
Orug N	Name/Form/Strength:								
		Length of Therapy:							
Oosing	Schedule: Length of T	herapy:							
	Schedule: Length of T ICD Code, i								
Diagno Veigh	osis: ICD Code, i	if applicable: veight obtain	 ed:						
Veight DIA met f	t (if applicable): Date was GNOSIS AND CLINICAL CRITERIA: Check below all for approval. To support each line checked, all documentation, includer chart notes, must be provided or request may be denied.	if applicable: veight obtain that apply. A uding lab resu	ed:	a must	be s,				
Veight DIA met f and/c	To Code, in the contraction of t	that apply. Auding lab resumegestrol ace	ed:	a must gnostics suspen	be s, nsion OR No				
Veight DIA met f and/c	To Code, is the first of the control	that apply. Auding lab resumegestrol ace	ed:	a must gnostics suspen	be s, nsion OR No o <u>TWO</u>				
DIA met f and/c	ICD Code, is the different conventional antiemetics (e.g., promethazine, prochlor etc.)? ICD Code, is the different conventional antiemetics (pregnancy-related nausea/vomiting)?	that apply. Auding lab resumegestrol ace	ed:	a must gnostics suspen	be s, nsion OR No o TWO lopramic				
DIA met f and/o	To Code, is the contraction of t	that apply. Auding lab resumegestrol ace	ed:	a must gnostice suspen constions to metoc	be s, nsion OR No o TWO lopramic				
Diagno Weight DIA met f and/c 1. 2.	To Code, it (if applicable): GNOSIS AND CLINICAL CRITERIA: Check below all a for approval. To support each line checked, all documentation, includer chart notes, must be provided or request may be denied. If diagnosis is AIDS-related wasting, has member tried and failed a contraindication, intolerance, drug-drug interaction? Member has tried and failed therapeutic doses of, or has adverse edifferent conventional antiemetics (e.g., promethazine, prochlor etc.)? Member has hyperemesis (pregnancy-related nausea/vomiting)? Bonjesta®/Diclegis®: length of approval, Estimated Delivery Date	that apply. Auding lab resumegestrol ace effects or continuer perazine, me	ed:	a must gnostice suspen constions to metoc	be s, nsion OR No o TWO lopramic No No				

PA Antiemetic/Antivertigo (Medicaid)

(Continued from previous page)

			Yes		No
For ondansetron 16 mg ODT : Has the member tried and failed or been intoler	rant to ondansetron 8 mg ODT?		Yes		No
Provide clinical evidence that the <u>Preferred</u> drug(s) will not provide adequate benefit and list pharmaceutical drugs attempted and outcome.					

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *