

Comprehensive Care Gap Documentation Guide

**For Medicare and Medicaid
All 2025 Measures**



Table of Contents

Introduction	1
Telehealth/Telemedicine Services	2
Adherence to Antipsychotic Medications for Individuals With Schizophrenia (SAA)	3
Adults' Access to Preventive/Ambulatory Health Services (AAP)	3
Adult Immunization Status (AIS-E)*	4
Advance Care Planning (ACP) *Medicare Only	5
Antibiotic Utilization for Respiratory Conditions (AXR)	6
Appropriate Testing for Pharyngitis (CWP)	6
Appropriate Treatment for Upper Respiratory Infection (URI)	7
Asthma Medication Ratio (AMR)	8
Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis (AAB)	8
Blood Pressure Control for Patients With Diabetes (BPD)	9
Blood Pressure Control for Patients with Hypertension (BPC-E)	10
Breast Cancer Screening (BCS-E)	11
Cardiac Rehabilitation (CRE)	12
Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC)	13
Care for Older Adults (COA) *Medicare Only	14
Care for Older Adults (COA) *Medicare Only - continued	15
Cervical Cancer Screening (CCS-E)	16
Child and Adolescent Well-care Visits (WCV)	17
Childhood Immunization Status (CIS-E)	18
Childhood Immunization Status (CIS-E) - continued	19
Chlamydia Screening	19
Colorectal Cancer Screening (COL-E)	20
Controlling High Blood Pressure (CBP)	21
Deprescribing of Benzodiazepines in Older Adults (DBO) *Medicare Only	22
Depression Remission or Response for Adolescents and Adults (DRR-E)	23
Depression Screening and Follow-up for Adolescents and Adults (DSF-E)*	24
Diabetes Monitoring for People With Diabetes and Schizophrenia (SMD)	24
Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD)	25
Documented Assessment After Mammogram (DBM-E)*	26
Emergency Department Visits for Hypoglycemia in Older Adults With Diabetes (EDH) *Medicare Only	27
Eye Exam for Patients With Diabetes (EED)	28

Table of Contents (2)

Fall Risk Management (FRM) *Medicare Only	29
Follow-up After Abnormal Mammogram Assessment (FMA-E)*	29
Follow-up After Emergency Department Visit for Mental Illness (FUM)*	30
Follow-up After Emergency Department Visit for People With Multiple High-risk Chronic Conditions (FMC) *Medicare Only	31
Follow-up After Emergency Department Visit for Substance Use (FUA)	32
Follow-up After High-intensity Care for Substance Use Disorder (FUI)	33
Follow-up After Hospitalization for Mental Illness (FUH)	34
Follow-up Care for Children Prescribed ADHD Medication (ADD-E)	35
Glycemic Status Assessment for Patients With Diabetes (GSD)	36
Immunizations for Adolescents (IMA-E)	37
Initiation and Engagement of Substance Use Disorder Treatment (IET)	38
Kidney Health Evaluation for Patients With Diabetes (KED)*	39
Lead Screening in Children (LCS) *Medicaid Only	40
Management of Urinary Incontinence in Older Adults (MUI) *Medicare Only	41
Medical Assistance With Smoking Cessation and Tobacco Use Cessation (MSC)	42
Medication Adherence for Cholesterol (Statins)	43
Medication Adherence for Diabetes Medications	44
Medication Adherence for Hypertension (RAS Antagonists)	45
Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM-E)*	46
Non-recommended PSA-based Screening in Older Men (PSA) *Medicare Only	47
Oral Evaluation Dental Services (OED) *Medicaid Only	47
Osteoporosis Screening in Older Women (OSW) *Medicare Only	48
Osteoporosis Management in Women Who Had a Fracture (OMW) *Medicare Only	49
Pediatric Quality Indicator 14: Asthma Admission Rate (PDI)	50
Pharmacotherapy Management of COPD Exacerbation (PCE)	51
Pharmacotherapy for Opioid Use Disorder (POD)	51
Physical Activity in Older Adults (PAO) *Medicare Only	52
Postpartum Depression Screening and Follow-up (PDS-E)	52
Potentially Harmful Drug-Disease Interactions in Older Adults (DDE) *Medicare Only	53
Prenatal and Postpartum Care (PPC)	54
Prenatal Depression Screening and Follow-up (PND-E)*	55
Prenatal Immunization Status (PRS-E)	56
Prevention Quality Indicator 05 (as calculated by DMAS): COPD or Asthma in Older Adults Admission Rate (PQI)	56
Prevention Quality Indicator 08 (as calculated by DMAS): Heart Failure Admission Rate (PQI)	57

Table of Contents (3)

Risk of Continued Opioid Use (COU)*	58
Social Need Screening and Intervention (SNS-E)	59
Statin Therapy for Patients With Cardiovascular Disease (SPC)	60
Statin Therapy for Patients With Diabetes (SPD)	61
Topical Fluoride for Children (TFC)	62
Transitions of Care (TRC) *Medicare Only	63
Unhealthy Alcohol Use Screening and Follow-up (ASF-E)*	64
Use of First-line Psychosocial Care for Children and Adolescents on Antipsychotics (APP)*	65
Use of High-risk Medications in Older Adults (DAE) *Medicare Only	66
Use of Imaging Studies for Low Back Pain (LBP)	67
Use of Opioids at High Dosage (HDO)	68
Use of Opioids From Multiple Providers (UOP)*	68
Utilization of the PHQ-9 To Monitor Depression Symptoms for Adolescents and Adults (DMS-E)*	69
Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents	70
Well-child Visits in the First 30 Months of Life (W30)	72
Glossary	73

Introduction

Preventive and chronic care management are critical steps along the pathway to helping your patients, our members, achieve optimal health. Sentara Health Plans is proud to partner with you to accomplish this very achievable goal. Electronic medical records may provide a means to track gaps in care and reminders of needed services. The Care Gap Documentation Guide is designed to help providers easily document the closure of care gaps.

This resource is organized for ease of use as follows:

- Measure definition.
- Identification of applicable quality program(s).
- Helpful tips to achieve performance measure.
- Codes recommended to document gap closure.

For additional information or assistance, you may contact your network management trainer.

Key:

VBC Measure – Value-Based Care Program Contractual Measure

STARS 2024 Focus Measure



Telehealth/Telemedicine Services

Please visit this **link** to review our guide on telehealth/telemedicine services;

As physicians, you can help improve the quality of care by:

- Encouraging your patients to schedule preventive exams.
- Reminding your patients to follow up with ordered tests and procedures.
- Making sure necessary services are being performed in a timely manner.
- Submitting claims with proper HEDIS codes.
- Accurately documenting all services and results (if appropriate) in the patient's medical record.

We need to work together to improve and maintain a higher quality of care. When our members are healthy, everyone benefits!



Adherence to Antipsychotic Medications for Individuals With Schizophrenia (SAA)

Definition: Adults ages 18 and older.

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members with diagnoses of schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for at least 80% of their treatment period.



Exclusion Criteria: Members with a diagnosis of dementia; those who did not have at least two antipsychotic medication-dispensing events; members in hospice or using hospice services during the measurement year and/or members who died anytime during the measurement year.

Adults' Access to Preventive/Ambulatory Health Services (AAP)

Definition: Adults ages 20 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

An ambulatory or preventive care visit

- In the measurement year **for Medicaid and Medicare members.**
- In the measurement year or two years prior **for commercial members.**



Exclusion Criteria: Members in hospice or using hospice services during the measurement year. Members who died anytime during the measurement year.

Adult Immunization Status (AIS-E)*

Definition: Adults ages 19 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

The percentage of members who are up to date on recommended routine vaccines for influenza, tetanus, and diphtheria (Td) or tetanus, diphtheria, and acellular pertussis (Tdap), zoster, pneumococcal and hepatitis B.



- **Influenza vaccine** on or between July 1 of the year prior to the measurement period and June 30 of the measurement period or members with anaphylaxis due to influenza vaccine any time before or during the measurement year.
- **Td or Tdap vaccine** at least one vaccine between 9 years prior to the start of the measurement period and the end of the measurement period; or members with a history of anaphylaxis or encephalitis due to diphtheria, tetanus, or pertussis vaccine.
- **Herpes zoster vaccine** received two doses of the recombinant vaccine at least 28 days apart on October 1, 2017, through the end of the measurement period or members with anaphylaxis due to the herpes zoster vaccine any time before or during the measurement period.
- **Pneumococcal vaccine** received at least one on or after the member's 19th birthday, before or during the measurement period; or had anaphylaxis due to pneumococcal vaccine any time before or during the measurement period.
- **Childhood hepatitis B** received at least three doses of the vaccine with different dates of service on or before their 19th birthday; or a hepatitis B vaccine series on or after their 19th birthday to include either a recommended two-dose adult hepatitis B vaccine administered at least 28 days apart or any other recommended adult hepatitis B vaccine administered on different dates of service.
- **Hepatitis B** received a hepatitis B surface antigen, antibody, or total antibody to hepatitis B core antigen test with a positive result, hepatitis B illness or anaphylaxis due to the hepatitis vaccine any time before or during the measurement period.

Exclusion Criteria: Members in hospice or using hospice services during the measurement year.

**Developed with support from the Department of Health and Human Services (DHHS), Office of the Assistant Secretary for Health (OASH), National Vaccine Program Office (NVPO) and The Hepatitis Education Project.*



Advance Care Planning (ACP) *Medicare Only

Definition: Adults ages 66–80 with advanced illness, an indication of frailty, or receiving palliative care; and adults ages 81 years and older who had advance care planning during the measurement year.

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Advance Care Planning is the discussion or documentation about preferences for resuscitation, life-sustaining treatment, or end-of-life care.



Exclusion Criteria: Members in hospice or using hospice services during the measurement year.

Codes

CPT Codes To Identify Advance Care Planning:

99483, 99497



Antibiotic Utilization for Respiratory Conditions (AXR)

Definition: Ages 3 months and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Diagnosis of a respiratory condition that resulted in an antibiotic dispensing event.

Intake Period: July 1 of the year prior to the measurement year to June 30 of the measurement year.



Exclusion Criteria: Members in hospice or using hospice services during the measurement year.

Note: This measure is designed to capture the frequency of antibiotic utilization for respiratory conditions. Organizations should use this information for internal evaluation only. NCQA does not view higher or lower service counts as indicating better or worse performance.

Appropriate Testing for Pharyngitis (CWP)

Definition: Ages 3 years and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

- Diagnosis of pharyngitis.
- Prescribed an antibiotic.
- Received a group A Streptococcus (strep) in the seven-day period from three days prior to, through three days after the episode date.



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

ICD-10 Codes To Identify Pharyngitis:

J02.0, J02.8–J02.9, J03.00–J03.01, J03.80–J03.81, J03.90–J03.91

Appropriate Treatment for Upper Respiratory Infection (URI)

Definition: Ages 3 months and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Diagnosis of upper respiratory infection (URI) and not prescribed an antibiotic.



Exclusion Criteria: Members in hospice or using hospice services during the measurement year. Members who died anytime during the measurement year.

Codes

ICD-10 Codes To Identify URI:

J00, J06.0, J06.9, J40

Pharyngitis:

J02.0, J02.8–J02.9, J03.00–J03.01, J03.80, J03.81, J03.90, J03.91



Asthma Medication Ratio (AMR)

Definition: Ages 5–64

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members identified as having persistent asthma with a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.



Exclusion Criteria: Members who had any diagnosis that requires a different treatment approach than members with asthma anytime during the member's history through December 31 of the measurement year. Members who had no asthma controller or reliever medications prescribed during the measurement year. Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

ICD-10 Codes To Identify Asthma:

J45.21, J45.22, J45.30, J45.31, J45.32, J45.40, J45.41, J45.42, J45.50, J45.51, J45.52, J45.901, J45.902, J45.909, J45.991, J45.998

Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis (AAB)

Definition: Ages 3 months and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Diagnosis of acute bronchitis/bronchiolitis and not prescribed an antibiotic.



Exclusion Criteria: Members in hospice or using hospice services during the measurement year. Members with a diagnosis of any of the following comorbid conditions: emphysema, COPD, immune system disorder, HIV, or malignant neoplasms. Members who died anytime during the measurement year.

Codes

ICD-10 Codes to Identify Acute Bronchitis:

J20.3–J20.9, J21.0–J21.1, J21.8–J21.9



Blood Pressure Control for Patients With Diabetes (BPD)

VBC Measure

Definition: Adults ages 18–75

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Diagnosis of diabetes (type 1 or type 2) whose blood pressure (BP) was adequately controlled (<140/90 mm Hg) during the measurement year.

Exclusion Criteria: Members who did not have a diagnosis of diabetes in any setting during the measurement year or the year prior but did have:

- Polycystic ovarian syndrome.
- Gestational diabetes.
- Steroid-induced diabetes.

Members in hospice, using hospice services, or receiving palliative care anytime during the measurement year. Members who died anytime during the measurement year.



Codes

ICD-10 Codes To Identify Diabetes:

E10.10, E10.11, E10.21, E10.22, E10.29, E10.8, E10.9 E11.00–E11.9, E13.00–E13.9

CPT/CPT II Codes for BP Values:

Systolic less than 130: 3074

Systolic greater than/equal to 140: 3077F

Systolic 130-139: 3075

Diastolic less than 80: 3078F

Diastolic 80–89: 3079F

Diastolic greater than/equal to 90: 3080F

Blood Pressure Control for Patients with Hypertension (BPC-E)

Definition: Adults age 18-85

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members who had a diagnosis of hypertension (HTN) and whose most recent blood pressure (BP) was, 140/90 mm Hg during the measurement period.

Identify the most recent BP reading noted during the measurement year.

The BP reading must occur on or after the date when the second diagnosis of hypertension (identified using the event/diagnosis criteria) occurred.



Exclusion Criteria: Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Members with a nonacute inpatient admission during the measurement period. Members with evidence of end-stage renal disease (ESRD), dialysis, nephrectomy, or kidney transplant anytime during the member's history on or prior to December 31 of the measurement year. Members with a diagnosis of pregnancy anytime during the measurement year. Members who died anytime during the measurement year.

**This is a first-year measure*



Breast Cancer Screening (BCS-E)

VBC Measure

Definition: Women 50–74 years of age (denominator) who had a mammogram to screen for breast cancer (numerator) during measurement year

Applicable Quality Program(s): HEDIS, CMS Medicare Advantage Star Rating

Helpful Tips To Achieve Performance Measure:

A mammogram to screen for breast cancer on or between October 1, 2023–December 31, 2025.



Exclusion Criteria: Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Bilateral mastectomy or both a right and left unilateral mastectomies anytime during the member's history through the end of the measurement year. Members who had gender-affirming chest surgery with a diagnosis of gender dysphoria anytime during the member's history through the end of the measurement year. Members who died anytime during the measurement year.

Codes

ICD-10-CM Codes:

Absence of left breast: Z90.12

Absence of right breast: Z90.11

Bilateral mastectomy: OHTV0ZZ

History of bilateral mastectomy: Z90.13

Mammography: 77061, 77062, 77063, 77065, 77066, 77067

Unilateral mastectomy: OHTU0ZZ (left), OHTT0ZZ (right)

**STARS
2024
Focus
Measure**



Cardiac Rehabilitation (CRE)

Definition: Adults ages 18 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members who attended cardiac rehabilitation following a qualifying cardiac event, (myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, heart and heart/lung transplantation, or heart valve repair/replacement).



Four rates are reported:

1. **Initiation:** Percentage of members who attended two or more sessions of cardiac rehabilitation within 30 days after a qualifying event.
2. **Engagement 1:** Percentage of members who attended 12 or more sessions of cardiac rehabilitation within 90 days after a qualifying event.
3. **Engagement 2:** Percentage of members who attended 24 or more sessions of cardiac rehabilitation within 180 days after a qualifying event.
4. **Achievement:** Percentage of members who attended 36 or more sessions of cardiac rehabilitation within 180 days after a qualifying event.

Exclusion Criteria: Members in hospice or using hospice services, receiving palliative care, had PCI, or died anytime during the measurement year. Discharged from an inpatient setting with any of the following on the discharge claim 180 days after the episode date: MI, CABG, heart or heart/lung transplant, or heart valve repair or replacement.

Codes

CPT Codes:

93797-93798



Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC)

Definition: Adults ages 18-64

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Percentage of members with schizophrenia or schizoaffective disorder and cardiovascular disease, who had an LDL-C test during the measurement year.



Exclusion Criteria:

Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

CPT Codes to Identify LDL-C Screening:

80061, 83700, 83701, 83704, 83721

Care for Older Adults (COA) *Medicare Only

VBC Measure

Definition: Adults ages 66 and older

Applicable Quality Program(s): HEDIS, CMS Medicare Advantage Star Rating

Helpful Tips To Achieve Performance Measure:

Members who had each of the following during the measurement year:

Medication Review:

- At least one medication review conducted by a prescribing practitioner or clinical pharmacist during the measurement year and the presence of a medication list in the medical record, as documented through either administrative data or medical record review.
- A medication list, signed and dated during the measurement year by the appropriate practitioner type (prescribing practitioner or clinical pharmacist), meets criteria (the practitioner's signature is considered evidence that the medications were reviewed).
- Documentation must come from the same medical record and must include one of the following:
 - A medication list in the medical record and evidence of a medication review by a prescribing practitioner or clinical pharmacist and the date when it was performed.
 - Notation that the member is not taking any medication and the date when it was noted.

**A review of side effects for a single medication at the time of prescription alone is not sufficient. An outpatient visit is not required to meet criteria. Do not include medication lists or medication reviews performed in an acute inpatient setting.*



Care for Older Adults (COA) *Medicare Only - Continued

Functional Status Assessment:

- At least one functional status assessment during the measurement year, as documented through either administrative data or medical record review.
- Documentation in the medical record must include evidence of a complete functional status assessment and the date when it was performed.
- A functional status assessment limited to an acute or single condition, event, or body system (e.g., lower back, leg) does not meet criteria for a comprehensive functional status assessment. The components of the functional status assessment numerator may take place during separate visits within the measurement year. Do not include comprehensive functional status assessments performed in an acute inpatient setting.



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

CPT Codes for COA:

Medication review: 1160F, 90863, 99483, 99605, 99606

Medication list: 1159F

Transition care management: 99495, 99496

Functional Status Assessment: 1170F, 99483

STARS
2024
Focus
Measure



Cervical Cancer Screening (CCS-E)

Definition: Women ages 21–64

- PAP test (cervical cytology) within the last three years for women ages 21–64.
- Cervical high-risk human papillomavirus (hrHPV) testing within the last five years for women ages 30–64.
- Cervical cytology/high-risk human papillomavirus (hrHPV) cotesting within the last five years for women ages 30–64.

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Documentation in the record must include both of the following:

- Date the test was performed.
- The result or finding.



Note: *Evidence of hrHPV testing within the last 5 years also captures patients who had cotesting; therefore, additional methods to identify cotesting are not necessary.*

Exclusion Criteria: Documentation of hysterectomy with no residual cervix, cervical agenesis, or acquired absence of cervix. Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Members who died anytime during the measurement year. Members with sex assigned at birth of male at any time in the patient's history.

Codes

CPT Codes To Identify Cervical Cancer Screening:

Cervical Cytology: 88141–88143, 88147–88148, 88150, 88152, 88153, 88164–88167, 88174–88175

High Risk HPV: 87624, 87625

Child and Adolescent Well-care Visits (WCV)

VBC Measure

Definition: Members ages 3–21

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

At least one (1) comprehensive well-care visit with a PCP or OB/GYN during the measurement year.

The well-care visit must occur with a PCP or an OB/GYN practitioner, but the practitioner does not have to be the practitioner assigned to the member.



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

CPT Codes To Identify Well-child Visits:

99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99461

ICD-10 Codes:

Z00.0, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z00.5, Z76.1, Z76.2

Childhood Immunization Status (CIS-E)

VBC Measure

Definition: Members who turn 2 years old during the measurement year. Vaccines must be completed on or before the second birthday:

- 4 DTaP or anaphylaxis or encephalitis due to diphtheria, tetanus, or pertussis vaccine (do not count any before 42 days of age).
- 3 IPV or anaphylaxis due to the IPV vaccine (do not count any before 42 days of age).
- 1 MMR; history of measles, mumps, and rubella; or anaphylaxis due to the MMR vaccine (do not count any before 42 days of age).
- 3 HiB or anaphylaxis due to HiB vaccine (do not count any before 42 days of age).
- 3 hepatitis B, anaphylaxis due to hepatitis B vaccine, positive serology, or history of hepatitis B.
- 1 VZV, anaphylaxis due to the VZV vaccine, positive serology, or documented history of chicken pox disease.
- 4 pneumococcal conjugates or anaphylaxis due to the pneumococcal conjugate vaccine (do not count any before 42 days of age).
- 1 hepatitis A, anaphylaxis due to the hepatitis A vaccine, or documented hepatitis A illness.
- 2 or 3 rotavirus vaccines - depends on the vaccine administered - or documented anaphylaxis due to the rotavirus vaccine (do not count any before 42 days of age).
- 2 influenza with different dates of service or anaphylaxis due to the influenza vaccine. One of the two vaccinations can be a live attenuated influenza vaccine (LAIV) if administered on the child's second birthday (do not count any given prior to 6 months of age).

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Parental refusal is not an exclusion.

- Documentation of "immunizations are up-to-date" is not acceptable.
- Documentation of an immunization (such as the first Hep B) received "at delivery" or "in the hospital" may be counted.
- For documented history of illness, a seropositive test result, or anaphylaxis, there must be a note indicating the date of the event, which must have occurred by the member's second birthday.



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who had a contraindication to a childhood vaccine on or before their second birthday. Members who died anytime during the measurement year.

Childhood Immunization Status (CIS-E) - continued

VBC Measure

Codes

42 Days of Age Through Second Birthday:

- (4) DTaP - Recommended Codes - CPT: 90697, 90698, 90700, 90723
- (3) IPV - Recommended Codes - CPT: 90697, 90698, 90713, 90723
- (3) HiB - Recommended Codes - CPT: 90644, 90647, 90648, 90697, 90698, 90748
- (4) Pneumococcal Conjugate - Recommended Code - CPT: 90670, 90671
- (2 or 3) Rotavirus - Recommended Codes- (2 Dose) CPT: 90681 or (3 Dose) CPT: 90680

On or Between First and Second Birthdays:

- (1) VZV - Recommended Codes - CPT: 90710, 90716
- (1) MMR - Recommended Codes - CPT: 90707; MMRV - 90710
- (1) Hepatitis A - Recommended Code - CPT: 90633

On or Before Second Birthday:

- (3) Hepatitis B - Recommended Codes - CPT: 90697, 90723, 90740, 90744, 90747, 90748

6 Months of Age Through Second Birthday:

- (2) Influenza - Recommended Codes - CPT: 90655, 90657, 90661, 90673, 90674, 90685, 90686, 90687, 90688, 90689, 90765

On or Between Ninth and Thirteenth Birthdays:

- LAIV - Recommended Codes - 90660, 90672

Chlamydia Screening

Definition: Age 16-24

Applicable Quality Program(s): HEDIS

Helpful Tips to Achieve Performance Measure:

Members who were recommended for routine chlamydia screening were identified as sexually active and had at least one test for chlamydia during the measurement year.



Exclusion Criteria: Sex Assigned at Birth: (LOINC code 76689-9) Male (LOINC code LA2-8) any time in the member's history. Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

CPT Codes for Chlamydia Screening:

87110; 87270; 87320; 87490- 87492; 87810



Colorectal Cancer Screening (COL-E)

VBC Measure

Definition: Adults ages 45–75

Applicable Quality Program(s): HEDIS, CMS Medicare Advantage Star Ratings

Helpful Tips To Achieve Performance Measure:

One or more of the following screenings:

- Colonoscopy in past 10 years (measurement year and nine years prior).
- Flexible sigmoidoscopy in past five years (measurement year and four years prior).
- CT colonography (e.g., virtual colonoscopy) in the past five years (measurement year and four years prior).
- FIT-DNA (e.g., Cologuard) test in the past three years (measurement year and two years prior).
- Fecal occult blood test (iFOBT or gFOBT) annually (measurement year).



Exclusion Criteria: Diagnosis of colorectal cancer or total colectomy anytime during the member's history through the measurement year. Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Members who died anytime during the measurement year.

Codes

FOBT:

CPT/CPT II: 82270, 82274

HCPCS: G0328

FIT-DNA Test:

CPT/CPT II: 81528. This code is specific to the Cologuard® FIT-DNA test.

Flexible Sigmoidoscopy:

CPT/CPT II: 45330–45335, 45337, 45338, 45340–45342, 45346, 45347, 45349, 45350

Computed Tomography (CT) Colonography:

CPT/CPT II: 74261–74263

Colonoscopy:

CPT/CPT II: 44388–44394, 44401–44408, 45378–45382, 45384–45386, 45388–45393, 45398

STARS
2024
Focus
Measure



Controlling High Blood Pressure (CBP)

VBC Measure

Definition: Adults ages 18–85

Adequate control is defined as: <140/90

Applicable Quality Program(s): HEDIS, CMS Medicare Advantage Star Ratings

Helpful Tips To Achieve Performance Measure:

Identify the most recent BP reading noted during the measurement year.

The BP reading must occur on or after the date when the second diagnosis of hypertension (identified using the event/diagnosis criteria) occurred.



Exclusion Criteria: Members in hospice or receiving palliative care anytime during the measurement year. Members with evidence of end-stage renal disease (ESRD), dialysis, nephrectomy, or kidney transplant anytime during the member's history on or prior to December 31 of the measurement year. Members with a diagnosis of pregnancy anytime during the measurement year. Members who died anytime during the measurement year.

Codes

ICD-10 Codes To Identify HTN: I10

CPT II Codes for BP Values:

Systolic less than 130: 3074

Systolic 130-139: 3075

Systolic greater than/equal to 140: 3077F

Diastolic less than 80: 3078F

Diastolic 80–89: 3079F

Diastolic greater than/equal to 90: 3080

**STARS
2024
Focus
Measure**



Deprescribing of Benzodiazepines in Older Adults (DBO) *Medicare Only

Definition: Adults ages 67 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members who were dispensed benzodiazepines and experienced a 20% decrease or greater in benzodiazepine dose (diazepam milligram equivalent [DME] dose) during the measurement year.



Exclusion Criteria: Members with a diagnosis of seizure disorder, REM sleep behavior disorder, benzodiazepine withdrawal, or ethanol withdrawal on or before January 1 of the year prior to the measurement year and the ITE start date. Members in hospice using hospice services, or palliative care. Members who died anytime during the measurement year.

Note: A lower rate represents better performance for all rates.

Depression Remission or Response for Adolescents and Adults (DRR-E)

Definition: Ages 12 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Diagnosis of depression with an elevated PHQ-9 (total score ≥ 9) who had evidence of response or remission within four to eight months of the elevated score.



Three rates are reported:

1. **Follow-up PHQ-9** within four to eight months after the initial elevated PHQ-9 score.
2. **Depression Remission:** Achieved remission within four to eight months after the initial elevated PHQ-9 score of <5 .
3. **Depression Response:** Showed a response within four to eight months after the initial elevated PHQ-9 score, with PHQ-9 score reduction of at least 50%.

Exclusion Criteria: Members with any of the following anytime during the member's history through the end of the measurement year: bipolar disorder, personality disorder, psychotic disorder, or pervasive developmental disorder. Members in hospice or using hospice services anytime during the measurement year.

**Adapted with financial support from the Agency for Healthcare Research and Quality (AHRQ) and the Centers for Medicare & Medicaid Services (CMS) under the CHIPRA Pediatric Quality Measures Program Centers of excellence grant number U18HS020503, and with permission from the measure developer, Minnesota Community Measurement.*

Codes

ICD-10 Codes To Identify Major Depression and Dysthymia:

F32.0–F32.5, F32.9, F33.0–F33.3, F43.40–F43.42, F33.9, F34.1

CPT Codes To Identify Interactive Outpatient Encounters:

90791, 90792, 90832, 90834, 90837, 98960–98962, 98967–98972, 98980, 98981, 99078, 99202–99205, 99211–99215, 99220, 99242–99245, 99341–99345, 99347–99350, 99381–99387, 99391–99397, 99401–99404, 99411–99412, 99421–99423, 99441–99443, 99457, 99458, 99483, 99492–99494, 99510



Depression Screening and Follow-up for Adolescents and Adults (DSF-E)*

Definition: Ages 12 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Screened for clinical depression using a standardized tool and, if screened positive, received follow-up care within 30 days following a positive depression screen finding.



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members with a history of bipolar disorder anytime during the member's history through the end of the year prior to the measurement year. Members with depression that starts in the prior year through the measurement year.

**Adapted with financial support from the Centers for Medicare & Medicaid Services (CMS).*

Diabetes Monitoring for People With Diabetes and Schizophrenia (SMD)

Definition: Adults ages 18–64

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members with diagnosis of schizophrenia or schizoaffective disorder and diabetes who had both an LCL-C test and HbA1c test during the measurement year.



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

CPT Codes To Identify:

HbA1c Tests: 83036, 83037

LDL-C Screening: 80061, 83700, 83701, 83704, 83721



Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD)

Definition: Adults ages 18–64

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

- Diagnosis of schizophrenia, schizoaffective disorder, or bipolar disorder.
- Dispensed an antipsychotic medication.
- Diabetes screening (glucose or HbA1c) test during the measurement year.



Exclusion Criteria: Members with diabetes or who were dispensed insulin or oral hypoglycemics/ antihyperglycemics during the measurement year or the year prior to the measurement year. Members who had no antipsychotic medications dispensed during the measurement year. Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

ICD-10 Codes To Identify Diagnosis:

Bipolar: F30.10–F30.13, F30.2–F30.4, F30.8–F30.9, F31.0, F31.10–F31.13, F31.2, F31.30–F31.32, F31.4–F31.5, F31.60–F31.64, F31.70–F31.78,

Schizophrenia: F20.0–F20.3, F20.5, F20.81, F20.89, F20.9, F25.0–F25.1, F25.8–F25.9

CPT Codes To Identify Diabetes Screening:

Glucose Tests: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951

HbA1c Tests: 83036, 83037



Documented Assessment After Mammogram (DBM-E)*

Definition: Adults ages 40-74

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Percentage of episodes of mammograms documented in the form of a BI-RADS assessment within 14 days of a mammogram.



Exclusion Criteria: Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Members who died anytime during the measurement year.

**This is a first-year measure*

**This measure was supported by Cooperative Award NU380T000303 from the Centers for Disease Control and Prevention and the National Network of Public Health Institutes (NNPHI). Its contents are the sole responsibility of the authors (NCQA) and do not necessarily represent the official position of the Centers for Disease Control and Prevention, the US Department of Health and Human Services, the US government, or the NNPHI.*



Emergency Department Visits for Hypoglycemia in Older Adults With Diabetes (EDH) *Medicare Only

Definition: Members 67 years of age or older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members with diabetes (types 1 and 2) who had an ED visit for hypoglycemia during the measurement year.



Two rates are reported:

1. Members with diabetes (types 1 and 2) who had an ED visit for hypoglycemia during the measurement year.
2. Members with diabetes (types 1 and 2) who had at least one dispensing event of insulin within each six-month treatment period from July 1 of the year prior to the measurement year through December 31 of the measurement year.

Exclusion Criteria: Members who do not have a diagnosis of diabetes in any setting during the measurement year or the year prior to the measurement year and who had a diagnosis of polycystic ovarian syndrome, gestational diabetes, or steroid-induced diabetes in any setting during the measurement year or the year prior to the measurement year. Members in hospice or using hospice services during the measurement year.



Eye Exam for Patients With Diabetes (EED)

VBC Measure

Definition: Adults ages 18–75

Applicable Quality Program(s): HEDIS, CMS Medicare Advantage Star Ratings

Helpful Tips To Achieve Performance Measure:

Diagnosis of diabetes (Types 1 and 2) who had a retinal exam. Screening or monitoring for diabetic retinal disease as identified by administrative data.

This includes diabetics who had one of the following:

- Retinal or dilated eye exam by an eye care professional (optometrist or ophthalmologist) in the measurement year.
- Negative retinal or dilated eye exam (negative for retinopathy) by an eye care professional (optometrist or ophthalmologist) in the measurement year.



Exclusion Criteria: Bilateral eye enucleation anytime during the member's history through December 31 of the measurement year. Members in hospice, using hospice services, or receiving palliative care anytime during the measurement year. Members who died anytime during the measurement year.

Codes

ICD-10 Codes To Identify Diabetes:

E10.10, E10.11, E10.21, E10.22, E10.29, E11.00–E11.9; E13.00–E13.9

CPT/CPT II Codes for Diabetic Retinal Screening:

92002, 92004, 92012, 92014, 92018, 92019, 92134, 92201, 92202, 92230, 92235, 92250, 99203–92205, 99213–99215, 99242–99245, 2022F–2026F, 2033F

STARS
2024
Focus
Measure



Fall Risk Management (FRM) *Medicare Only

Definition: Adults ages 65 and older.

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Two components of this measure that assess different facets of fall risk management:

1. Discussing fall risk

- Seen by a practitioner in the past 12 months.
- Discussed falls or problems with balance or walking with their current practitioner.

2. Managing fall risk

- Had a fall or problems with balance or walking in the past 12 months.
- Seen by a practitioner in the past 12 months.
- Received a recommendation for how to prevent falls or treat problems with balance or walking from their current practitioner.



Follow-up After Abnormal Mammogram Assessment (FMA-E)*

Definition: Adults ages 65 and older.

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Percentage of episodes with inconclusive or high-risk BI-RADS assessments that received appropriate follow-up within 90 days of the assessment.

Exclusion Criteria: Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Members who died anytime during the measurement year.



**This is a first-year measure*

**This measure was supported by Cooperative Award NU380T000303 from the Centers for Disease Control and Prevention and the National Network of Public Health Institutes (NNPHI). Its contents are the sole responsibility of the authors (NCQA) and do not necessarily represent the official position of the Centers for Disease Control and Prevention, the US Department of Health and Human Services, the US government, or the NNPHI.*



Follow-up After Emergency Department Visit for Mental Illness (FUM)*

VBC Measure

Definition: Ages 6 years and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Percentage of emergency department visits with a principal diagnosis of mental illness or any diagnosis of intentional self-harm and had a mental health follow-up service.



Two rates are reported:

1. The percentage of ED visits for which the member received follow-up within seven days of the ED visit (eight total days).
2. The percentage of ED visits for which the member received follow-up within 30 days of the ED visit (31 total days).

Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

Behavioral Health Outpatient Visit:

CPT: 98960–98962, 99078, 99201–99205, 99211–99215, 99217–99220 99241–99245, 99341–99345, 99347–99350, 99381–99387, 99391–99397, 99401–99404, 99411, 99412, 99483, 99492–99494, 99510

**Adapted from an NCQA measure with financial support from the Office of the Assistant Secretary for Planning and Evaluation (ASPE) under Prime Contract No. HHSP23320100019WI/ HHSP23337001T, in which NCQA was a subcontractor to Mathematica. Additional financial support was provided by the Substance Abuse and Mental Health Services Administration (SAMHSA).*

Follow-up After Emergency Department Visit for People With Multiple High-risk Chronic Conditions (FMC)

***Medicare Only**

VBC Measure

Definition: Ages 18 and older

Applicable Quality Program(s): HEDIS, CMS Medicare Advantage Star Rating

Helpful Tips To Achieve Performance Measure:

Members with emergency department (ED) visits who have multiple high-risk chronic conditions such as COPD/asthma, dementia, CKD, major depression, heart failure, MI, atrial fibrillation, or stroke who had a follow-up service within seven days of the ED visit (eight total days).



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

Outpatient and Telehealth Visit:

CPT: 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99281, 99282, 99283, 99284, 99285, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99429, 99455, 99456, 99483

**STARS
2024
Focus
Measure**



Follow-up After Emergency Department Visit for Substance Use (FUA)

VBC Measure

Definition: Ages 13 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

ED visit with principal diagnosis of substance use disorder (SUD) or any diagnosis of drug overdose for which there was a follow-up.



Two rates are reported:

1. Follow-up visit within seven days of the ED visit (eight total days).
2. Follow-up visit within 30 days of the ED visit (31 total days).

Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

Outpatient and Telehealth Visit:

CPT: 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99281, 99282, 99283, 99284, 99285, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99429, 99455, 99456, 99483



Follow-up After High-intensity Care for Substance Use Disorder (FUI)

Definition: Ages 13 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

- Acute inpatient hospitalization, residential treatment, or withdrawal management visits.
- Diagnosis of substance use disorder.
- Follow-up visit or service for substance use disorder.



Two rates are reported:

1. The percentage of visits or discharges for which members received follow-up for substance use disorder within seven days after the visit or discharge.
2. The percentage of visits or discharges for which members received follow-up for substance use disorder within 30 days after the visit or discharge.

Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.



Follow-up After Hospitalization for Mental Illness (FUH)

Definition: Ages 6 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Percentage of discharges for members hospitalized for a principal diagnosis of mental illness, or any diagnosis of intentional self-harm, and had a mental health follow up service.



Two rates are reported:

1. Follow-up visit within seven days of discharge.
2. Follow-up visit within 30 days of discharge.

Do not include services that occur on the date of discharge.

Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year.
Members who died anytime during the measurement year.



Follow-up Care for Children Prescribed ADHD Medication (ADD-E)

Definition: Children ages 6–12

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members with a newly prescribed ADHD medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was prescribed.



Two rates are reported:

1. **Initiation Phase:** One follow-up visit with a practitioner with prescribing authority within 30 days.
2. **Continuation and Maintenance Phase:** Remained on the medication for at least 210 days and had two additional visits with a practitioner within 270 days (nine months) after the initiation phase ended.

Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members with a diagnosis of narcolepsy anytime during their history through the measurement year. Members who died anytime during the measurement year.



Glycemic Status Assessment for Patients With Diabetes (GSD)

VBC Measure

Definition: Adults ages 18–75

Applicable Quality Program(s): HEDIS, CMS Medicare Advantage Star Ratings

Helpful Tips To Achieve Performance Measure:

Diagnosis of diabetes (Types 1 and 2) whose most recent glycemic status (hemoglobin A1c (HbA1c) or glucose management indicator (GMI) was at the following levels from measurement year:

- Glycemic Status (<8.0%).
- Glycemic Status (<9.0%).



Exclusion Criteria: Members in hospice or using hospice services or palliative care anytime during the measurement year. Members who died anytime during the measurement year.

Codes

ICD-10 Codes To Identify Diabetes:

E10.10, E10.11, E10.21, E10.22, E10.29, E10.8, E10.9 E11.00–E11.9, E13.00–E13.9

CPT/CPT II for HbA1c:

83036, 83037, 3044F, 3046F, 3051F, 3052F

STARS
2024
Focus
Measure



Immunizations for Adolescents (IMA-E)

Definition: Children who turn 13 years old during the measurement year

- 1 dose meningococcal vaccine between the 11th and 13th birthdays, or anaphylaxis due to the vaccine anytime on or before the member's 13th birthday, and
- 1 tetanus, diphtheria toxoids, and acellular pertussis (Tdap) vaccine between the 10th and 13th birthdays; or anaphylaxis or encephalitis due to the vaccine anytime on or before the 13th birthday, and
- 2-dose or 3-dose HPV vaccination series between the 9th and 13th birthdays, or anaphylaxis due to the vaccine anytime on or before the 13th birthday.

To align with ACIP recommendations:

- Meningococcal: the quadrivalent (serogroups A, C, W and Y) and pentavalent meningococcal vaccines (serogroups A, C, W, Y and B) are included in the measure.
- HPV: the minimum interval for the two-dose HPV vaccination schedule is 150 days, with a 4-day grace period (146 days).

***Note:** The measure calculates a rate for each vaccine and two combination rates.

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

- All vaccines must be completed on or before the 13th birthday.
- For documented history of anaphylaxis, there must be a note indicating the date of the event, which must have occurred by the member's 13th birthday.



Exclusion Criteria: Members in hospice or using hospice services during the measurement year.
Members who died anytime during the measurement year.

- Parental refusal is not an exclusion.

Codes

CPT Codes:

Meningococcal CPT: 90619, 90733, 90734

Tdap CPT: 90715

HPV-CPT: 90649–90651

Initiation and Engagement of Substance Use Disorder Treatment (IET)

VBC Measure

Definition: Ages 13 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

New substance use disorder (SUD) episodes that results in treatment initiation and engagement.



Two rates are reported:

1. **Initiation of SUD Treatment:** Within 14 days of the diagnosis through an inpatient SUD admission, outpatient visit, intensive outpatient encounter or partial hospitalization, telehealth, or medication treatment.
2. **Engagement of SUD Treatment:** Evidence of treatment engagement within 34 days of the initiation.

Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

Codes To Identify AOD Visits:

CPT: 98960–98962, 98980, 98981, 98966–98968 99078, 99202–99205, 99211–99215, 99241–99245, 99341–99345, 99347–99350, 99381–99387, 99391–99397, 99401–99404, 99408, 99409, 99411, 99412, 99483, 99492–99494, 99510



Kidney Health Evaluation for Patients With Diabetes (KED)*

Definition: Adults ages 18–85

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Diagnosis of diabetes (type 1 and type 2) who received a kidney health evaluation during the measurement year, defined by:

- An estimated glomerular filtration rate (eGFR) **and**
- A urine albumin-creatinine ratio (uACR).



Exclusion Criteria: Members with evidence of ESRD or dialysis anytime during the member's history on or prior to December 31 of the measurement year. Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Members who died anytime during the measurement year.

Codes:

ICD-10 Codes To Identify Diabetes:

E10.10, E10.11, E10.21, E10.22, E10.29, E11.00–E11.9, E13.00–E13.9

CPT Codes:

eGFR: 80047–80048, 80050, 80053, 80069, 82565

uACR: 82043, 82570

**This measure was developed by NCQA with input from the National Kidney Foundation.*

STARS
2024
Focus
Measure



Lead Screening in Children (LCS) *Medicaid Only

Definition: Children who turn 2 years old during the measurement year

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

At least one or more capillary or venous lead blood test by their second birthday.

Documentation in the record must include both of the following:

- Date the test was performed.
- Result of finding.



Exclusion Criteria: Members in hospice or using hospice services during the measurement year.
Members who died anytime during the measurement year.

Codes

CPT Codes: 83655



Management of Urinary Incontinence in Older Adults (MUI) *Medicare Only

Definition: Adults ages 65 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Three components that assess the management of urinary incontinence in older adults:



1. Discussing urinary incontinence

- Reported having urine leakage in the past six months.
- Discussed their urinary leakage problem with a healthcare provider.

2. Discussing treatment of urinary incontinence

- Reported having urine leakage in the past six months.
- Discussed treatment options for their current urine leakage problem.

3. Impact of urinary incontinence

- Reported having urine leakage in the past six months.
- Reported that urine leakage made them change their daily activities or interfered with their sleep a lot.

Exclusion Criteria: Evident from CMS administrative records of a hospice start date.

Note: A lower rate indicates better performance for this indicator.

Medical Assistance With Smoking Cessation and Tobacco Use Cessation (MSC)

Definition: Current smoker/tobacco user ages 18 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

This measure assesses **three different components** of providing medical assistance with smoking and tobacco use cessation:



1. **Advising smokers and tobacco users to quit**
 - Received advice to quit during the measurement year.
2. **Discussing Cessation Medications**
 - Discussed or were recommended cessation medications during the measurement year.
3. **Discussing cessation strategies**
 - Discussed or were provided cessation methods or strategies during the measurement year.

Medication Adherence for Cholesterol (Statins)

VBC Measure

Definition: Any member 18 years or older who has at least two fills of a designated medication(s) as defined in each measure. The measurement is calculated using Proportion of Days Covered (PDC) using Pharmacy Data Encounter (PDE) data. PDC is the number of days the member has medication on hand/is covered by the medication based on the dates of prescription fills (numerator) compared to the number of days in the time period (denominator).

Applicable Quality Program(s): CMS Medicare Advantage Star Ratings

Helpful Tips To Achieve Performance Measure:

The measurement is calculated using Proportion of Days Covered (PDC) using Pharmacy Data Encounter (PDE) data. PDC is the number of days the member has medication on hand/ is covered by the medication based on the dates of prescription fills (numerator) compared to the number of days in the time period (denominator). Members will only be recognized when filling medication through Sentara Health Plans.

This measure is calculated daily and is not a “gap” that can be closed.

- **Write for 90-day supplies:** This will ensure that a member is “covered” for 90 days. This gives the member a better chance to make the 80% PDC threshold throughout the year.
- **Update prescriptions:** If the dose changes, make sure to update the prescription with the new directions.
- **Ask members about barriers to medication use:** Cost of medications is well-known as a barrier, but other factors are just as prevalent. Can members get to the pharmacy? Can members read the small labels? Do members know how to take the medications?

MAC Medication Inclusion: Statin medications

Exclusion Criteria: Hospice enrollment, ESRD diagnosis or coverage dates.



Medication Adherence for Diabetes Medications

VBC Measure

Definition: Any member 18 years or older who has at least two fills of a designated medication(s) as defined in each measure.

The measurement is calculated using Proportion of Days Covered (PDC) with Pharmacy Data Encounter (PDE) data. PDC is the number of days the member has medication on hand/is covered by the medication based on the dates of prescription fills (numerator) compared to the number of days in the time period (denominator).

Applicable Quality Program(s): CMS Medicare Advantage Star Ratings

Helpful Tips To Achieve Performance Measure:

This measure is calculated daily and is not a “gap” that can be closed.

- **Write for 90-day supplies:** This will ensure that a member is “covered” for 90 days. This gives the member a better chance to make the 80% PDC threshold throughout the year.
- **Update prescriptions:** If the dose changes, make sure to update the prescription with the new directions.
- **Ask members about barriers to medication use:** Cost of medications is well-known as a barrier, but other factors are just as prevalent. Can members get to the pharmacy? Can members read the small labels? Do members know how to take the medications?
- Only medications filled through Sentara Health Plans will be recognized.
- This measure is calculated daily and is not a “gap” that can be closed.

Medication Inclusion: Biguanides, sulfonylureas, thiazolidinediones, and dipeptidyl peptidase (DPP)-IV inhibitors, incretin mimetics, meglitinides, and sodium glucose cotransporter 2 (SGLT2) inhibitors.



Medication Adherence for Hypertension (RAS Antagonists)

VBC Measure

Definition: The measurement is calculated using Proportion of Days Covered (PDC) with Pharmacy Data Encounter (PDE) data. PDC is the number of days the member has medication on hand/is covered by the medication based on the dates of prescription fills (numerator) compared to the number of days in the time period (denominator).

Applicable Quality Program(s): CMS Medicare Advantage Star Ratings

Helpful Tips To Achieve Performance Measure:

Any member 18 years or older who has at least two fills of a designated medication(s) as defined in each measure. The measurement is calculated using PDC with PDE data. PDC is the number of days the member has medication on hand/ is covered by the medication based on the dates of prescription fills (numerator) compared to the number of days in the time period (denominator). Only medications filled through Sentara Health Plans will be recognized.

This measure is calculated daily and is not a “gap” that can be closed.

- **Write for 90-day supplies:** This will ensure that a member is “covered” for 90 days. This gives the member a better chance to make the 80% PDC threshold throughout the year.
- **Update prescriptions:** If the dose changes, make sure to update the prescription with the new directions.
- **Ask members about barriers to medication use:** Cost of medications is well-known as a barrier, but other factors are just as prevalent. Can members get to the pharmacy? Can members read the small labels? Do members know how to take the medications?

Medication Inclusion: Angiotensin converting enzyme inhibitor (ACEI), angiotensin receptor blocker (ARB), or direct renin inhibitor medications.

Exclusion Criteria: Hospice enrollment, ESRD diagnosis or coverage dates, one or more prescriptions for sacubitril/valsartan.



Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM-E)*

Definition: Children and adolescents ages 1–17

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Children and adolescents who had two or more antipsychotics prescriptions and had metabolic testing.

Three rates are reported:

1. Children and adolescents on antipsychotics who received blood glucose testing.
2. Children and adolescents on antipsychotics who received cholesterol testing.
3. Children and adolescents on antipsychotics who received blood glucose and cholesterol testing.

Exclusion Criteria: Members in hospice or using hospice services during the measurement year.
Members who died anytime during the measurement year.



Codes

CPT Codes To Identify Cholesterol Tests Other Than LDL: 82465, 83718, 83722, 84478

**Developed with financial support from the Agency for Healthcare Research and Quality (AHRQ) and the Centers for Medicare & Medicaid Services (CMS) under the CHIPRA Pediatric Quality Measures Program Centers of Excellence grant number U18 HS020503.*



Non-recommended PSA-based Screening in Older Men (PSA) *Medicare Only

Definition: Men ages 70 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Screened unnecessarily for prostate cancer using prostate-specific antigen (PSA)-based screening during the measurement year.



Exclusion Criteria: Prostate cancer diagnosis anytime during the member's history through December 31 of the measurement year. Dysplasia of the prostate anytime during the measurement year or the year prior to the measurement year. A PSA test during the year prior to the measurement year where laboratory data indicate an elevated (>4.0 ng/ml) or abnormal result. Dispensed prescription for a 5-alpha reductase inhibitor during the measurement year. Members in hospice or using hospice services during the measurement year. Members who died anytime during the measurement year.

Codes

CPT codes for PSA-based Screening:

84152–84154

Oral Evaluation Dental Services (OED) *Medicaid Only

Definition: Members 21 years of age and younger

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members who received a comprehensive or periodic oral evaluation with a dental provider during the measurement year.



**This measure has been included in and/or adapted for HEDIS with the permission of the Dental Quality Alliance (DQA) and American Dental Association (ADA). © 2023 DQA on behalf of ADA, all rights reserved.*

Exclusion Criteria: Members in hospice or using hospice services during the measurement year. Members who died anytime during the measurement year.

Osteoporosis Screening in Older Women (OSW)

*Medicare Only

Definition: Women ages 65–75

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Received one or more osteoporosis screening tests on or between the member's 65th birthday and December 31 of the measurement year.



Exclusion Criteria: Members who had a claim/encounter for osteoporosis therapy anytime in the member's history through December 31 of the year prior to the measurement year. Members who were prescribed a prescription to treat osteoporosis anytime on or between January 1, 2022–December 31, 2024. Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Members who died anytime during the measurement year.

Codes

CPT Codes:

76977, 77078, 77080–77081, 77085



Osteoporosis Management in Women Who Had a Fracture (OMW) *Medicare Only

VBC Measure

Definition: Women ages 65–85

Applicable Quality Program(s): HEDIS, CMS Medicare Advantage Star Ratings

Helpful Tips To Achieve Performance Measure:

Suffered a fracture and had one of the following in the six months after the fracture:

- A bone mineral density (BMD) test **or**
- A prescription for a drug to treat osteoporosis.



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who received palliative care anytime during the intake period through the end of the measurement year. Members who died anytime during the measurement year.

Codes

Bone Mineral Density Tests:

CPT/CPT II: 76977, 77078, 77080, 77081, 77085, 77086

Osteoporosis Medications:

HCPCS: J0897, J1740, J3110, J3489

STARS
2024
Focus
Measure



Pediatric Quality Indicator 14: Asthma Admission Rate (PDI)

VBC Measure

Definition: Members ages 2 through 17 with a principal ICD-10-CM diagnosis code for asthma (ACSASTD)

Applicable Quality Program(s): AHRQ

Helpful Tips To Achieve Performance Measure:

- Development of a written asthma action plan in partnership with patient/family.
- Monitor medication compliance.



Exclusion Criteria: Excludes cases with a diagnosis code for cystic fibrosis and anomalies of the respiratory system, obstetric admissions, and transfers from other institutions.

Codes

Asthma Diagnosis Codes:

J4521, J4552, J4522, J45901, J4531, J45902, J4532, J45990, J4541, J45991, J4542, J45998, J4551

Pharmacotherapy Management of COPD Exacerbation (PCE)

Definition: Adults ages 40 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Discharged from an acute inpatient admission or an ED visit with a primary diagnosis of COPD on or between January 1–November 30 of the measurement year.

Prescribed appropriate medications (or already had an active prescription for):

- A systemic corticosteroid within 14 days of the event.
- A bronchodilator within 30 days of the event.



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

ICD-10 Codes To identify COPD:

J.41.0, J41.1, J42, J43.0–J43.2, J43.8, J44.1, J44.81, J44.89, J44.9

Pharmacotherapy for Opioid Use Disorder (POD)

Definition: Ages 16 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Diagnosis of opioid use disorder (OUD) pharmacotherapy events that lasted at least 180 days among members 16 years of age and older with a diagnosis of OUD and a new OUD pharmacotherapy event.



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Physical Activity in Older Adults (PAO) *Medicare Only

Definition: Adults ages 65 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Two components of this measure that assess different facets of promoting physical activity in older adults:



1. **Discussing Physical Activity**

- Had a doctor's visit in the past 12 months.
- Spoke with a doctor or other healthcare provider about their level of exercise or physical activity.

2. **Advising Physical Activity**

- Had a doctor's visit in the past 12 months.
- Received advice to start, increase, or maintain their level of exercise or physical activity.

Exclusion Criteria: Evidence from CMS guidelines administrative records of a hospice start date.

Postpartum Depression Screening and Follow-up (PDS-E)

Applicable quality program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Screened for clinical depression during the postpartum period, and if screened positive, received follow-up care.



Two rates are reported:

1. **Depression Screening:** Screened for clinical depression using a standardized instrument during the postpartum period (7–84 days following date of delivery).
2. **Follow-up on Positive Screen:** Received follow-up care within 30 days of a positive depression screen finding.

Exclusion Criteria: Deliveries in which members were in hospice or using hospice services anytime during the measurement period. Members who died during the measurement year.

Potentially Harmful Drug-Disease Interactions in Older Adults (DDE) *Medicare Only

Definition: Adults ages 65 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Evidence of an underlying disease, condition, or health concern, and who were dispensed an ambulatory prescription for a potentially harmful medication, concurrent with, or after the diagnosis.



Three rates are reported:

1. **A history of falls** and a prescription for antiepileptics, antipsychotics, benzodiazepines, nonbenzodiazepine hypnotics, or antidepressants (SSRIs, tricyclic antidepressants, SNRIs).
2. **Dementia** and a prescription for antipsychotics, benzodiazepines, nonbenzodiazepine hypnotics, tricyclic antidepressants, or anticholinergic agents.
3. **Chronic kidney disease** and a prescription for COX-2 selective NSAIDs or nonaspirin NSAIDs.

Exclusion Criteria: Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Members with a diagnosis of psychosis, schizophrenia, schizoaffective disorder, or bipolar disorder on or between January 1, 2024 to December 1, 2025. Members who died anytime during the measurement year.

Note: A lower rate represents better performance for all rates.

Prenatal and Postpartum Care (PPC)

VBC Measure

Definition: Members who delivered a live infant between October 8 of the year prior to the measurement year and October 7 of the measurement year.

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Two components to assess prenatal and postpartum care:

1. **Timeliness of prenatal care** – Received a prenatal care visit in the first trimester, on or before the enrollment start date, or within 42 days of enrollment in the organization.
2. **Postpartum care** – Had a postpartum visit on or between 7 and 84 days after delivery.



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.

Codes

Timeliness of Prenatal Care Recommended Codes:

CPT: 99202-99205, 99211-99215, 99242-99245, 99421-99423, 99483, 98970-98972, 99457, 99458, 98980, 98981, 98957, 98967

Postpartum Care Recommended Codes:

CPT: 57170, 58300, 59430, 99501

Prenatal Depression Screening and Follow-up (PND-E)*

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Screened for clinical depression while pregnant and, if screening is positive, received follow-up care.



Two rates are reported:

1. **Depression Screening:** Screened for clinical depression during pregnancy using a standardized instrument.
2. **Follow-up on Positive Screen:** Received follow-up care within 30 days of a positive depression screening finding.

Exclusion Criteria: Deliveries in which members were in hospice or using hospice services anytime during the measurement period. Deliveries that occurred at less than 37 weeks gestation. Members who died during the measurement year.

**Developed with support from the California HealthCare Foundation (CHCF). CHCF works to ensure that people have access to the care they need, when they need it, at a price they can afford. Visit <https://www.chcf.org/> to learn more. Also supported by the Zoma Foundation.*

Prenatal Immunization Status (PRS-E)

Definition: Deliveries in the measurement year

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Received the following recommended vaccines during the measurement year:

- Influenza.
- Tetanus, diphtheria toxoids, and acellular pertussis (Tdap) during the pregnancy (including on the delivery date).



Exclusion Criteria: Deliveries in which members were in hospice or using hospice services anytime during the measurement period. Deliveries that occurred at less than 37 weeks gestation. Members who died during the measurement year.

Prevention Quality Indicator 05 (as calculated by DMAS): COPD or Asthma in Older Adults Admission Rate (PQI)

VBC Measure

Definition: Admissions with a principal diagnosis of chronic obstructive pulmonary disease (COPD) or asthma per 100,000 population, ages 40 years and older.

Applicable Quality Program(s): AHRQ

Helpful Tips To Achieve Performance Measure:

Discharges, for patients ages 40 years and older, with either:

- Principal ICD-10-CM diagnosis code for COPD (ACCOPDD*) (excluding acute bronchitis) **or**
- Principal ICD-10-CM diagnosis code for asthma (ACSASTD*).



Exclusion Criteria: Excludes obstetric admissions and transfers from other institutions.

Codes

Primary Diagnosis Codes for COPD and Asthma:

J41.0, J41.1, J41.8, J42, J43.0, J43.1, J43.2, J43.8, J43.9, J44.0, J44.1, J44.9, J47.0, J47.1, J47.9, J45.21, J45.22, J45.31, J45.32,



Prevention Quality Indicator 08 (as calculated by DMAS): Heart Failure Admission Rate (PQI)

VBC Measure

Definition: Admissions with a principal diagnosis of heart failure per 100,000 population, ages 18 years and older

Applicable Quality Program(s): AHRQ

Helpful Tips To Achieve Performance Measure:

Discharges, for patients ages 18 years and older, with a principal ICD-10-CM diagnosis code for heart failure.



Exclusion Criteria: Discharges, for patients ages 18 years and older, with a principal ICD-10-CM diagnosis code for heart failure.

Codes

Primary Diagnosis Codes for Heart Failure:

I09.81, I11.0, I13.0, I13.2, I50.1, I50.20, I50.21, I50.22, I50.23, I50.30, I50.31, I50.32, I50.33, I50.4, I50.41, I50.42, I50.43, I50.9, I50.810, I50.811, I50.812, I50.813, I50.814, I50.82, I50.83, I50.84, I50.89



Risk of Continued Opioid Use (COU)*

Definition: Adults ages 18 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

A new episode of opioid use that puts them at risk for continued opioid use.

Two rates are reported:

1. The percentage of members with at least 15 days of prescription opioids in a 30-day period.
2. The percentage of members with at least 31 days of prescription opioids in a 62-day period.



Exclusion Criteria: Members in hospice or using hospice services during the measurement year. Members who had at least one of the following during the 12 months (1 year) prior to the prescription start date through 61 days after the prescription start date: cancer, sickle cell disease, or palliative care. Members who died anytime during the measurement year.

Note: A lower rate represents better performance for all rates.

**Adapted with financial support from CMS and with permission from the measure developer, Pharmacy Quality Alliance (PQA).*



Social Need Screening and Intervention (SNS-E)

Definition: Members \leq 17–65 years

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members who were screened, using prespecified instruments, at least once during the measurement year for unmet food, housing, and transportation needs, and received a corresponding intervention within one month if they screened positive.



Food Screening: The percentage of members who were screened for food insecurity.

Food Intervention: The percentage of members who received a corresponding intervention within 30 days (1 month) of screening positive for food insecurity.

Housing Screening: The percentage of members who were screened for housing instability, homelessness, or housing inadequacy.

Housing Intervention: The percentage of members who received a corresponding intervention within 30 days (1 month) of screening positive for housing instability, homelessness, or housing inadequacy.

Transportation Screening: The percentage of members who were screened for transportation insecurity.

Transportation Intervention: The percentage of members who received a corresponding intervention within 30 days (1 month) of screening positive for transportation insecurity.

Exclusion Criteria: Members in hospice or using hospice services during the measurement year. Medicare members 66 years of age and older by the end of the measurement year who meet either of the following:

- Enrolled in Institutional SNP (I-SNP) during the measurement year.
- Living long-term in an institution during the measurement year, as identified by the LTI flag in the Monthly Membership Detail Data File.
- Members who died during the measurement year.



Statin Therapy for Patients With Cardiovascular Disease (SPC)

VBC Measure

Definition: Males ages 21–75; females ages 40–75

Applicable quality program(s): CMS Medicare Advantage Star Ratings

Helpful Tips To Achieve Performance Measure:

Identified as having clinical atherosclerotic cardiovascular disease (ASCVD) during the measurement year.



Two rates are reported:

1. **Received Statin Therapy:** Members who were dispensed at least one high or moderate-intensity statin medication during the measurement year.
2. **Statin Adherence 80%:** Members who remained on a high or moderate-intensity statin medication for at least 80% of the treatment period (from prescription date through end of year).

Exclusion Criteria: Diagnosis of pregnancy or had IVF during the measurement year or the year prior. Diagnosis of ESRD, dialysis, or cirrhosis during the measurement year or the year prior. Diagnosis of myalgia, myositis, myopathy, or rhabdomyolysis during the measurement year. Dispensed at least one prescription for clomiphene during the measurement year or the year prior. Member in hospice or using hospice services or receiving palliative care anytime during the measurement year. Member died anytime during the measurement year.

Codes

Moderate or High-intensity Statin Therapy:

Atorvastatin: 10–80mg
Amlodipine-atorvastatin: 10–80mg
Rosuvastatin: 5–40mg
Simvastatin: 20–80mg
Ezetimibe-simvastatin: 20–80mg
Pravastatin: 40–80mg
Lovastatin: 40mg
Fluvastatin 40–80mg
Pitavastatin 1–4mg

STARS
2024
Focus
Measure

ICD-10 Codes for Myalgia/Myositis/Myopathy/Rhabdomyolysis:

G72.0, G72.2, G72.9, M60.80–M60.812, M60.819, M60.821–M60.822, M60.829, M60.831–M60.832, M60.839, M60.841–M60.842, M60.849, M60.851–M60.852, M60.859, M60.861–M60.862, M60.869, M60.871–M60.872, M60.879, M60.88–M60.89, M60.9, M62.82, M79.1–M79.12, M79.18



Statin Therapy for Patients With Diabetes (SPD)

VBC Measure

Definition: Adults ages 40–75

Applicable Quality Program(s): CMS Medicare Advantage Star Ratings

Helpful Tips To Achieve Performance Measure:

Identified as having diabetes and does not have clinical atherosclerotic cardiovascular disease (ASCVD) who met the following criteria:

Two rates are reported:

1. **Received Statin Therapy:** Members who were dispensed at least one statin medication of any intensity during the measurement year.
2. **Statin Adherence 80%:** Members who remained on a statin medication of any intensity for at least 80% of the treatment period (from prescription date through end of year).

Exclusion Criteria: Members diagnosed with myalgia, myositis, myopathy, or rhabdomyolysis in the measurement year. Members with any of the following: IVD diagnosis, pregnancy/IVF, prescribed clomiphene, ESRD or dialysis, cirrhosis. Members who had at least one of the following during the year prior to the measurement year: discharged from an inpatient setting with MI or had a CABG, PCI, or another revascularization procedure. Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Members who died anytime during the measurement year.



Codes

Myalgia/Myositis/Myopathy/Rhabdomyolysis:

G72.0, G72.2, G72.9, M60.80-M60.812, M60.819, M60.821, M60.822, M60.829, M60.831, M60.832, M60.839, M60.841, M60.842, M60.849, M60.851, M60.852, M60.859, M60.861, M60.862, M60.869, M60.871, M60.872, M60.879, M60.88, M60.89, M60.9, M62.82, M79.11, M79.12, M79.18



Topical Fluoride for Children (TFC)

Definition: Members ages 1–4 years

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members who received at least two fluoride varnish applications during the measurement year.

**This measure has been included in and/or adapted for HEDIS with the permission of the Dental Quality Alliance (DQA) and American Dental Association (ADA). © 2024 DQA on behalf of ADA, all rights reserved.*



Exclusion Criteria: Members in hospice or using hospice services anytime during the measurement year. Members who died anytime during the measurement year.



Transitions of Care (TRC) *Medicare Only

VBC Measure

Definition: Adults ages 18 and older

Applicable Quality Program(s): HEDIS, CMS Medicare Advantage Star Rating

Helpful Tips To Achieve Performance Measure:



1. Notification of Inpatient Admission

- On the day of admission through two days after (three total days).

2. Receipt of Discharge Information

- On the day of discharge through two days after (three total days).

3. Patient Engagement After Inpatient Discharge

- (E.g., office visits, home visits, telehealth) provided within 30 days after discharge.

4. Medication Reconciliation Post Discharge

- On the date of discharge through 30 days after discharge (31 total days).

Exclusion Criteria: Members in hospice or using hospice services during the measurement year.
Members who died during the measurement year.

Codes

CPT Codes for Transitional Care Management Services and Medication Reconciliation:

99483, 99495, 99496, 111F

CPT Codes for Patient Engagement:

98970-98972, 98980, 98981, 98966-98968, 99202-99205, 99211-99215, 99242-99245,
99341, 99342, 99344, 99345, 99347-99349, 99350, 99381-99384, 99385-99387,
99393-99397, 99401-99404, 99412, 99411, 99421-99423, 99429, 99441-99443,
99456, 99455, 99457, 99458, 99483

**STARS
2024
Focus
Measure**



Unhealthy Alcohol Use Screening and Follow-up (ASF-E)*

Definition: Adults ages 18 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

- Had a systematic screening for unhealthy alcohol use.
- If screened positive, received brief counseling or other follow-up care within two months of screening positive for unhealthy alcohol use.



Exclusion Criteria: Members with alcohol use disorder that starts during the year prior to the measurement year. Members with a history of dementia anytime during the member's history through the end of the measurement year. Members in hospice or using hospice services during the measurement year. Members who died during the measurement year.

**Adapted with financial support from the Substance Abuse and Mental Health Services Administration (SAMHSA) and with permission from the measure developer, the American Medical Association (AMA).*



Use of First-line Psychosocial Care for Children and Adolescents on Antipsychotics (APP)*

Definition: Ages 1–17 years

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members who had a **new** prescription for an antipsychotic medication and had documentation of psychosocial care as first-line treatment.



Exclusion Criteria: Members for whom first-line antipsychotic medications may be clinically appropriate: members with a diagnosis of schizophrenia, schizoaffective disorder, bipolar disorder, other psychotic disorder, autism, or other developmental disorder during the measurement year. Members in hospice or using hospice services during the measurement year. Members who died anytime during the measurement year.

Codes:

CPT Codes To Identify Psychosocial Care:

90832–90834, 90836–90840, 90845–90847, 90849, 90853, 90875–90876, 90880

**Developed with financial support from the Agency for Healthcare Research and Quality (AHRQ) and CMS under the CHIPRA Pediatric Quality Measures Program Centers of Excellence grant number U18HS020503, from a measure developed by MedNet Medical Solutions.*



Use of High-risk Medications in Older Adults (DAE)

*Medicare Only

Definition: Adults ages 67 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members who had a least two dispensing events for the same high-risk medication.

Three rates are reported:

1. The percentage of Medicare members 67 years of age and older who had at least two dispensing events for high-risk medications to avoid from the same drug class.
2. The percentage of Medicare members 67 years of age and older who had at least two dispensing events for high-risk medications to avoid from the same drug class, except for appropriate diagnoses.
3. Total rate (the sum of the two numerators divided by the denominator, deduplicating for members in both numerators).



Exclusion Criteria: Members in hospice or using hospice services or receiving palliative care during the measurement year. Members who died anytime during the measurement year.

Note: A lower rate represents a better performance.

Use of Imaging Studies for Low Back Pain (LBP)

Definition: Adults ages 18–75

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Principal diagnosis of low back pain who did not have an imaging study (plain X-ray, MRI, CT) within 28 days of the diagnosis.



Exclusion Criteria: Members in hospice or using hospice services or receiving palliative care anytime during the measurement year. Members who died anytime during the measurement year. Diagnosis of cancer, HIV, spondylopathy, major organ transplant, or history of major organ transplant, osteoporosis therapy, or a dispensed prescription to treat osteoporosis or lumbar surgery anytime during the member's history through 28 days after the IESD. Trauma or a fragility fracture anytime during the 90 days prior to the IESD through 28 days after the IESD. IV drug abuse, neurological impairment, or spinal infection anytime during the 365 days prior to IESD through 28 days after IESD. Prolonged use of corticosteroids: 90 consecutive days of corticosteroid treatment anytime during the 365 days prior to the IESD and ends on the IESD.

Index Episode Start Date (IESD) is defined as the earliest date of service for an eligible encounter during the intake period (January 1 through December 3, 2025) with a principal diagnosis of low back pain.

Codes

ICD-10 Codes To Identify Uncomplicated Low Back Pain:

M47.26–M47.28, M47.816–M47.818, M47.896–M47.898, M48.061–M48.07, M48.08, M51.16– M51.17, M51.26, M51.27–M51.36, M51.37, M51.86, M51.87, M53.2X6–M53.2X8–M53.88, M54.16– M54.9, M99.03–M99.84, S33.100A–S33.9XXA, S39.002A–S39.92XS

Note: A higher score indicates appropriate treatment of low back pain (the portion for whom imaging did not occur).



Use of Opioids at High Dosage (HDO)

Definition: Adults ages 18 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Received a prescription for opioids at a high dosage (average morphine milligram equivalent dose [MME] ≥ 90) for ≥ 15 days during the measurement year.



Exclusion Criteria: Members with cancer or sickle cell disease anytime during the measurement year. Members in hospice or using hospice services or receiving palliative care during the measurement year. Members who died anytime during the measurement year.

Note: A lower rate indicates better performance.

Use of Opioids From Multiple Providers (UOP)*

Definition: Adults ages 18 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Received a prescription for ≥ 15 days from multiple providers during the measurement year.



Three rates are reported: Opioid prescriptions received from and/or filled at:

1. **Multiple Prescribers** (four or more different prescribers).
2. **Multiple Pharmacies** (four or more different pharmacies).
3. **Multiple Prescribers and Multiple Pharmacies** (both four or more different prescribers and four or more different pharmacies).

Exclusion Criteria: Members in hospice or using hospice services during the measurement year. Members who died anytime during the measurement year.

Note: A lower rate represents better performance for all rates.

**Adapted with financial support from CMS and with permission from the measure developer, Pharmacy Quality Alliance (PQA).*



Utilization of the PHQ-9 To Monitor Depression Symptoms for Adolescents and Adults (DMS-E)*

Definition: Ages 12 and older

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Members with diagnosis of major depression or dysthymia, and had an outpatient encounter with a PHQ-9 score present in their record in the same assessment period as the encounter.



Exclusion Criteria: Members with any of the following anytime during the member's history through the end of the measurement year: bipolar disorder, personality disorder, psychotic disorder, or pervasive developmental disorder. Members in hospice or using hospice services during the measurement year.

**Adapted with financial support from the Agency for Healthcare Research and Quality (AHRQ) and the Centers for Medicare & Medicaid Services (CMS) under the CHIPRA Pediatric Quality Measures Program Centers of excellence grant number U18HS020503, and with permission from the measure developer, Minnesota Community Measurement.*



Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents

Definition: Members 3–17 years of age who had an outpatient visit with a PCP or an OB/GYN during the measurement year and had evidence of the following documented:

- BMI percentile*.
- Counseling for nutrition.
- Counseling for physical activity.

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

BMI Percentile documentation must include date, height, and weight.

- **BMI percentile** - May be plotted on age-growth chart.
- **Weight and height** - Must be taken during the measurement year.

Counseling for nutrition documentation must include a note indicating the **date** and **at least one of the following**:

- Discussion of current nutrition behaviors (e.g., eating habits, dieting behaviors).
- Checklist indicating nutrition was addressed.
- Counseling or referral for nutrition education.
- Educational materials on nutrition during a face-to-face visit.
- Anticipatory guidance for nutrition.
- Weight or obesity counseling.
- Referral to WIC.

Counseling for physical activity documentation includes a note indicating the **date** and **at least one of the following**:

- Discussion of current physical activity behaviors (e.g., exercise routine, participation in sports activities, exam for sports participation).
- Checklist indicating physical activity was addressed.
- Counseling or referral for physical activity.
- Member received educational materials on physical activity during a face-to-face visit.
- Anticipatory guidance specific to the child's physical activity.
- Weight or obesity counseling.



Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents - continued

Documentation of meeting Developmental Milestones only does not meet HEDIS® criteria for Physical Activity Counseling.

Services specific to an acute or chronic condition do not count toward the counseling indicators for either nutrition or physical activity.



Exclusion Criteria: Members in hospice or using hospice services during the measurement year. Members who have a diagnosis of pregnancy anytime during the measurement year. Members who died anytime during the measurement year.

Codes

ICD-10 Codes:

BMI Percentile: Z68.51-Z68.54

Counseling for Nutrition: Z71.3

Counseling for Physical Activity: Z02.5, Z71.82

CPT Codes:

Counseling for Nutrition: 97802–97804

**Because BMI norms for youth vary with age and gender, this measure evaluates whether BMI percentile is assessed rather than an absolute BMI value.*



Well-child Visits in the First 30 Months of Life (W30)

Definition: Children who turned 15 months or 30 months during measure year.

Applicable Quality Program(s): HEDIS

Helpful Tips To Achieve Performance Measure:

Well-child visits with a PCP/pediatrician during the last 15 months.

Two rates are reported:

1. **Six (6)** or more well-child visits on different dates of service **on or before the 15-month birthday.**
2. **Two (2)** or more well-child visits on different dates of service **between the child's 15-month birthday plus one day and the 30-month birthday.**



Exclusion Criteria: Members in hospice or using hospice services during the measurement year.
Members who died anytime during the measurement year.

Codes

CPT Codes To Identify Well-child Visits:

99381–99385, 99391–99395, 99461

ICD-10 Codes:

Z00.00–Z00.01,
Z00.110–Z00.111, Z00.121,
Z00.129, Z00.2, Z00.3,
Z02.5, Z02.84, Z76.1–Z76.2



Glossary

Attributed Member: Member for whom the provider is held accountable in regards to care.

CPT Category II Code: Tracking codes, ending with an "F," which facilitate data collection related to quality and performance measurement.

CPT Code: Medical code set used to report medical, surgical, diagnostic procedures, and other services by physicians/providers/facilities to health insurance companies and accreditation organizations.

Denominator: The number of members who qualify for the measure criteria.

Drug Tiers: A way for insurance providers to determine medicine costs. The higher the tier, the higher the cost of the medicine for the member in general.

HCPCS Code: Healthcare Common Procedure Coding System (often pronounced hick picks). A set of codes, beginning with a letter, used to report supplies, materials, drugs, procedures, and other services.

HEDIS: Health Care Effectiveness Data and Information Set. Standardized performance measures developed by NCQA (National Committee for Quality Assurance).

ICD-10-CM (Diagnosis Code): A code system used by physicians and other healthcare providers to classify and code all diagnoses, signs, and symptoms.

ICD-10 (Procedure Code): A code system used to report procedures performed by physicians and other healthcare providers in a facility/hospital setting.

Numerator: The number of members who meet compliance criteria.

PMPM: Per member per month. Usual unit of measure that payers remit to providers.

Measurement Year: January 1 through December 31.

Stars: CMS rating system used to measure how well Medicare Advantage and Part D plans perform in several areas, including quality of care and customer satisfaction. Stars ratings range from one to five, with one being the lowest score and five being the highest.

Step Therapy: Trying less expensive options before "stepping up" to drugs that cost more.



sentarahealthplans.com/providers



Provider Services 1-800-881-2166

Medical: 1-800-229-8822

Behavioral Health: 1-800-648-8420

