

Genetic Testing—Pre-Treatment or Post-Intervention

Table of Content
<u>Purpose</u>
<u>Description & Definitions</u>
<u>Criteria</u>
<u>Coding</u>
<u>Document History</u>
<u>References</u>
<u>Special Notes</u>
<u>Keywords</u>

<u>Effective Date</u>	2/2009
<u>Next Review Date</u>	2/15/2024
<u>Coverage Policy</u>	Medical 34B
<u>Version</u>	1

Member-specific benefits take precedence over medical policy and benefits may vary across plans. Refer to the individual’s benefit plan for details*.

Purpose:

This policy addresses genetic testing for the prevention diagnosis and treatment of various conditions.

Description & Definitions:

Autosomal recessive: A genetic condition that appears only in individuals who have received two copies of an autosomal gene, one copy from each parent. The gene is on an autosome, a nonsex chromosome. The parents are carriers who have only one copy of the gene and do not exhibit the trait because the gene is recessive to its normal counterpart gene

X-linked recessive inheritance - hereditary pattern in which a recessive gene on the X chromosome results in the manifestation of characteristics in male offspring and a carrier state in female offspring.

Criteria:

Genetic Testing is considered **medically necessary** for the prevention diagnosis and treatment of patients who meet **All** of the following:

- There is an approved mutation specific treatment available
- There is sufficient Published Scientific Evidence or 3rd party Consensus in the Medical Community that the results of the specific genetic testing improves clinical outcomes
- After completion of a thorough history, physical examination, pedigree analysis, genetic counseling, relevant diagnostic and biochemical tests (if any), the patient meets criteria for **1 or more** of the following approved tests:
 - **HPV DNA** is covered without criteria
 - **Factor V Leiden** is covered without criteria. (81241)
 - **Prothrombin/Factor II Gene Mutation Analysis** is covered without criteria. (81240)
 - **PCR (Polymerase Chain Reaction) Testing for Hepatitis C or B** is covered without criteria or Pre-Authorization. – May also be addressed by Avalon
 - **Thiopurine S-Methyltransferase (TPMT)** is covered without criteria.
 - **Fabry Disease** is medical necessary for individual **genetic** female on a case by case basis and requires Medical Director approval.
 - **Hemophilia A (Factor VIII) or Hemophilia B (Factor IX)** is covered when **All** of the following are met:

- The individual displays clinical features, or is at direct risk of inheritance;
- The result of the test will directly impact the treatment being delivered to the individual
- After history, physical examination, and completion of conventional diagnostic studies (e.g. Factor VIII, Factor IX levels), a definitive diagnosis remains uncertain
- **Aplastic Anemia (FISH) fluorescence in situ hybridization** is covered without criteria.
- **HLA-B27** is medically necessary when used to rule out autoimmune disorders.
- **Hereditary Hemochromatosis** - HFE gene testing may be indicated when **ALL** of the following are present: (81256)
 - Diagnosis or screening of HFE-hereditary hemochromatosis, as indicated by **1 or more** of the following
 - Confirmation of diagnosis in adult with clinical suspicion of HFE-related hemochromatosis, as indicated by **1 or more** of the following:
 - Transferrin-iron saturation higher than 45%
 - Serum ferritin above upper limit of normal, and other more common causes of elevated ferritin (eg, acute inflammation, alcohol abuse, metabolic syndrome) have been evaluated with inconclusive results
 - Unexplained chronic liver disease combined with increased transferrin saturation
 - Porphyria cutanea tarda
 - Chondrocalcinosis
 - Hepatocellular carcinoma
 - Late-onset type 1 diabetes;
 - Screening of siblings or parents of individual homozygous for C282Y mutation
 - Screening of reproductive partner of individual with HFE- related hemochromatosis
 - Testing is accompanied by genetic counseling
- **HLA haplotype testing to detect genes specific to Celiac disease** are considered medically necessary for patients suspected of having the disease but who have indeterminate serology and/or inconclusive biopsy results (examples include Prometheus HLA-DQA1, HLA-DQB1, HLA-DQ2, HLA-DQ8 testing) are covered without medical review.
- **Severe Combined Immunodeficiency (SCID) Genetic Testing** for individuals with inconclusive screening results from an immune function test or T cell and B cell counts considered eligible for transplant to confirm diagnosis.
- **HLA-A29 Uveitis - HLA I TYPING ALLELE HR** screening will be paid upon request when ordered to rule out or confirm Birdshot chorioretinopathy (BSCR), a rare form of autoimmune posterior uveitis that affects visual function that if left untreated can lead to sight-threatening complications and loss of central vision

The following tests are considered **not medically necessary** for any use other than those indication in the clinical criteria, to include but not limited to:

- Acute Porphyria: HMBS, CPOX, PPOX genes
- Alport gene testing; (COL4A3 and COL4A4)
- Atypical Hemolytic Uremic Syndrome Complement Panel
- Autoimmune lymphoproliferative syndrome ALPS and FAS gene testing ;
- BetaGlobin analysis;
- BIRC4 test is testing for X-linked lymphoproliferative syndrome (XLP), caused by alterations, also known as "mutations," at a specific area within an individual's genetic information.
- CLCN1 DNA Sequencing Test for Myotonia congenital;
- Charcot Marie Tooth is not medically necessary;
- deCODE T2™, is not medically necessary
- deCODE Glaucoma test, is not medically necessary
- Idiopathic Hypercalcemia of Infancy (CYP24A1 gene)
- IFNL3 & INFNL4 Gene Analysis Associated with Hepatitis C Virus Clearance (**81400**)
- Kennedy's disease (KD) or X-linked spinal and bulbar muscular atrophy (SBMA) or spinobulbar muscular atrophy or X-Linked bulbo-spinal atrophy.
- Lactose intolerance (Prometheus Lab LactoType);
- MaculaRisk PGx

- MTHFR (5,10-methylenetetrahydrofolate reductase) (eg, hereditary hypercoagulability) gene analysis, common variants (eg, 677T, 1298C) **(81291)**
- Narcolepsy gene testing DRB1*15, DQA1*0102, and DQB1*0602
- Porphyria cutanea tarda: UROD gene
- PROMETHEUS Crohn's Prognostic **(81599)**
- PROMETHEUS IBD sgi Diagnostic
- Prometheus Thiopurine Metabolites
- Proove Narcotic Risk, Drug Metabolism and Pain Perception Profiles **(81225, 81226, 81227)**
- Repeat/Duplicative genetic testing
- Transforming Growth Factor Beta-Induced (TGFBI) **(81333)**
- Whole Exome Sequencing **(WES)** all indications,
- Exome Sequence Analysis **(81415, 81416, 81417)**
- Whole-genome sequencing in which a member's entire DNA is sequenced,
- Genome Sequence Analysis **(81425, 81426, 81427)**

Coding:

Medically necessary with criteria:

Coding	Description
81240	F2 (prothrombin, coagulation factor II) (eg, hereditary hypercoagulability) gene analysis, 20210G>A variant
81241	F5 (coagulation factor V) (eg, hereditary hypercoagulability) gene analysis, Leiden variant
81256	HFE (hemochromatosis) (eg, hereditary hemochromatosis) gene analysis, common variants (eg, C282Y, H63D)
81381	HLA Class I typing, high resolution (ie, alleles or allele groups); one allele or allele group (eg, B*57:01P), each
81383	HLA Class II typing, high resolution (ie, alleles or allele groups); one allele or allele group (eg, HLA-DQB1*06:02P), each
81400	Molecular pathology procedure, Level 1 (eg, identification of single germline variant [eg, SNP] by techniques such as restriction enzyme digestion or melt curve analysis)
81404	Molecular pathology procedure, Level 5 (eg, analysis of 2-5 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of 6-10 exons, or characterization of a dynamic mutation disorder/triplet repeat by Southern blot analysis)
81479	Unlisted molecular pathology procedure
85245	Clotting; factor VIII, VW factor, ristocetin cofactor
85246	Clotting; factor VIII, VW factor antigen
85247	Clotting; factor VIII, von Willebrand factor, multimeric analysis
85250	Clotting; factor IX (PTC or Christmas)
81238	F9 (coagulation factor IX) (eg, hemophilia B), full gene sequence
81599	Unlisted multianalyte assay with algorithmic analysis

Considered Not Medically Necessary:

Coding	Description
--------	-------------

81225	CYP2C19 (cytochrome P450, family 2, subfamily C, polypeptide 19) (eg, drug metabolism), gene analysis, common variants (eg, *2, *3, *4, *8, *17)
81226	CYP2D6 (cytochrome P450, family 2, subfamily D, polypeptide 6) (eg, drug metabolism), gene analysis, common variants (eg, *2, *3, *4, *5, *6, *9, *10, *17, *19, *29, *35, *41, *1XN, *2XN, *4XN)
81227	CYP2C9 (cytochrome P450, family 2, subfamily C, polypeptide 9) (eg, drug metabolism), gene analysis, common variants (eg, *2, *3, *5, *6)
81291	MTHFR (5,10-methylenetetrahydrofolate reductase) (eg, hereditary hypercoagulability) gene analysis, common variants (eg, 677T, 1298C)
81333	TGFBI (transforming growth factor beta-induced) (eg, corneal dystrophy) gene analysis, common variants (eg, R124H, R124C, R124L, R555W, R555Q)
81415	Exome (eg, unexplained constitutional or heritable disorder or syndrome); sequence analysis
81416	Exome (eg, unexplained constitutional or heritable disorder or syndrome); sequence analysis, each comparator exome (eg, parents, siblings) (List separately in addition to code for primary procedure)
81417	Exome (eg, unexplained constitutional or heritable disorder or syndrome); re-evaluation of previously obtained exome sequence (eg, updated knowledge or unrelated condition/syndrome)
81425	Genome (eg, unexplained constitutional or heritable disorder or syndrome); sequence analysis
81426	Genome (eg, unexplained constitutional or heritable disorder or syndrome); sequence analysis, each comparator genome (eg, parents, siblings) (List separately in addition to code for primary procedure)
81427	Genome (eg, unexplained constitutional or heritable disorder or syndrome); re-evaluation of previously obtained genome sequence (eg, updated knowledge or unrelated condition/syndrome)

Document History:

Revised Dates:

- 2022: February
- 2020: December
- 2020: October
- 2019: February
- 2019: January
- 2018: June
- 2017: June
- 2016: August
- 2016: July
- 2016: May
- 2016: April
- 2015: December
- 2015: November
- 2015: October
- 2015: August
- 2015: July
- 2015: April
- 2015: March
- 2015: February
- 2014: December
- 2014: November
- 2014: October
- 2014: August
- 2014: July
- 2014: June
- 2014: May
- 2014: March
- 2013: December
- 2013: November
- 2013: October
- 2013: July
- 2013: January
- 2012: September
- 2012: July
- 2012: March
- 2011: September
- 2011: August
- 2011: July
- 2011: June
- 2011: March

Reviewed Dates:

- 2010: August
- 2014: July
- 2015: December

- 2020: December

Effective Date:

- February 2009

References:

Specialty Association Guidelines; Government Regulations; Winifred S. Hayes, Inc; UpToDate; Literature Review; Specialty Advisors; National Coverage Determination (NCD); Local Coverage Determination (LCD).

(2018). Retrieved January 14, 2019, from Hayes: <http://www.hayesinc.com/hayes/>

(2018, May 14). Retrieved Jan 16, 2019, from MCG: <https://careweb.careguidelines.com/ed22/index.html>

(2019, 01 01). Retrieved January 14, 2019, from UpToDate: <https://www.uptodate.com/home>

(2019). Retrieved from National Comprehensive Cancer Network: https://www.nccn.org/professionals/physician_gls/default.aspx#detection

(2019). Retrieved from DynamedPlus: <http://www.dynamed.com/resultlist?q=TGFBI&filter=all>

(2019, Jan). Retrieved Jan 24, 2019, from Centers for Medicare and Medicaid Services: <https://www.cms.gov/medicare-coverage-database/details/lcd-details.aspx?LCDId=35025&ver=80&SearchType=Advanced&CoverageSelection=Both&NCSelection=NCA%7cCAL%7cNCD%7cMEDCAC%7cTA%7cMCD&ArticleType=SAD%7cEd&PolicyType=Both&s=53&CptHcpcsCode=81333&kq=true&bc=l>

(2019). Retrieved from American Optometric Association: <https://www.aoa.org/search?q=TGFBI>

(2019). Retrieved Jan 25, 2019, from American Ophthalmological Society: https://www.aosonline.org/search/SearchForm?Search=tgfbf&action_results=Go

Garcia-Castellanos, R., Nielsen, N. S., Runager, K., Thogersen, I. B., Lukassen, M. V., Poulsen, E. T., . . . Gomis-Ruth, F. X. (2017, Nov 7). Structural and Functional Implications of Human Transforming Growth Factor Beta-Induced Protein, TGFBIp, in Corneal Dystrophies. *Structure*, 25(11), pp. 1740-1750.e2. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0969212617302927?via%3Dihub>

Identification and characterization of transforming growth factor beta-induced in circulating tumor cell subline from pancreatic cancer cell line. (2018, Nov). *Cancer Science*, 109(11), pp. 3623-3633. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6215881/>

S.K., G., M.F., S., H.W., Y., & Y.S., L. (2018). Enhanced Expression of TGFBI promotes the Proliferation and Migration of Glioma Cells. *Cellular Physiology and Biochemistry*, 49, pp. 1138-1150. Retrieved from <https://www.karger.com/Article/FullText/493293>

Genetics Home Reference. CYP24A1 gene: cytochrome P450 family 24 subfamily A member 1. Published: Jan 2, 2019; Cited: Jan 4, 2018. <https://ghr.nlm.nih.gov/gene/CYP24A1>

Rodd, C. & Goodyer, P. Hypercalcemia of the newborn: etiology, evaluation, and management. *Pediatr Nephrol* (1999) 13:542–547. 23 Nov 1998; Cited: Jan 04, 2019. https://www.researchgate.net/publication/12848142_Hypercalcemia_of_the_newborn_Etiology_evaluation_and_management/download

Shane, E. Diagnostic approach to hypercalcemia. Updated: Aug 13, 2018; Cited: Jan 04, 2019. https://www.uptodate.com/contents/diagnostic-approach-to-hypercalcemia?search=idiopathic%20hypercalcemia&source=search_result&selectedTitle=3~150&usage_type=default&display_rank=3#H10

Lietman, S.A. et al. Hypercalcemia in Children and Adolescents. *Curr Opin Pediatr*. 2010 Aug; 22(4): 508–515. doi: 10.1097/MOP.0b013e32833b7c23. Cited: Jan 04, 2019. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2967024/>

Paolis, E.D. et al. A rapid screening of a recurrent CYP24A1 pathogenic variant opens the way to molecular testing for Idiopathic Infantile Hypercalcemia (IIH). *Clin Chim Acta*. 2018 Jul;482:8-13. doi: 10.1016/j.cca.2018.03.024. Cited: Jan 04, 2019. <https://www.ncbi.nlm.nih.gov/pubmed/29574006>

Kagi, L. et al. Regulation of vitamin D metabolizing enzymes in murine renal and extrarenal tissues by dietary phosphate, FGF23, and 1,25(OH)2D3. *Plos One*. Published: May 17, 2018. Cited: Jan 04, 2018. <https://doi.org/10.1371/journal.pone.0195427>

ABO blood group but not haemostasis genetic polymorphisms significantly influence thrombotic risk: a study of 180 homozygotes for the Factor V Leiden mutation. *Br J Haematol* 2006; 135:697.

Aloj G, et al., 2012, Severe Combined Immunodeficiencies: New and Old Scenarios, *Internat Revs of Immunol* 31:43-65.

Buckley, RH, 2004, The multiple causes of human SCID, *J Clin Imm* 114:1409-11.

Butte MJ et al., 2007, IL-7 receptor deficient SCID with a unique intronic mutation and post-transplant autoimmunity due to chronic GVHD, *Clinical Immunology* 125(2):159-64.

Fischer A et al., 2005, Severe combined immunodeficiency, a model disease for molecular immunology and therapy, *Immunological Reviews*. 203:98-109.

Fisher, M, Fernández, JA, Amerisco, SF, et al. Activated protein C resistance in ischemic stroke not due to factor V arginine506 to glutamine mutation. *Stroke* 1996; 27:1163.

Huck K, Hanenberg H, Gudowius S, et al. Delayed diagnosis and complications of Fanconi anaemia at advanced age--a paradigm. *Br J Haematol* 2006; 133:188.

Lensen, R, Bertina, RM, Vandenbroucke, JP, Rosendaal, FR. High factor VIII levels contribute to the thrombotic risk in families with factor V Leiden. *Br J Haematol* 2001; 114:380.

Levine RL, Belisle C, Wadleigh M, et al. X-inactivation-based clonality analysis and quantitative JAK2V617F assessment reveal a strong association between clonality and JAK2V617F in PV but not ET/MMM, and identifies a subset of JAK2V617F-negative ET and MMM patients with clonal hematopoiesis. *Blood* 2006; 107:4139.

Marsh JC, Ball SE, Cavenagh J, et al. Guidelines for the diagnosis and management of aplastic anaemia. *Br J Haematol* 2009; 147:43.

Morange, PE, Tregouet, DA, Frere, C, et al. Biological and genetic factors influencing plasma factor VIII levels in a healthy family population: results from the Stanislas cohort. *Br J Haematol* 2005; 128:91.

Puck, JM, 2007, Population-based newborn screening for severe combined immunodeficiency: steps toward implementation, *J Allergy Clin Immunol* 120:760-768.

Tefferi A. The history of myeloproliferative disorders: before and after Dameshek. *Leukemia* 2008; 22:3.

Vannucchi AM, Guglielmelli P, Tefferi A. Advances in understanding and management of myeloproliferative neoplasms. *CA Cancer J Clin* 2009; 59:171.

Special Notes: *

This medical policy expresses Sentara Health Plan's determination of medical necessity of services, and they are based upon a review of currently available clinical information. Medical policies are not a substitute for clinical judgment or for any prior authorization requirements of the health plan. These policies are not an explanation of benefits.

Medical policies can be highly technical and complex and are provided here for informational purposes. These medical policies are intended for use by health care professionals. The medical policies do not constitute medical advice or medical care. Treating health care professionals are solely responsible for diagnosis, treatment and medical advice. Sentara Health Plan members should discuss the information in the medical policies with their treating health care professionals. Medical technology is constantly evolving and these medical policies are subject to change without notice, although Sentara Health Plan will notify providers as required in advance of changes that could have a negative impact on benefits.

Keywords:

F2, Prothrombin, F5, Factor V, Hemochromatosis, HFE, TGFBI, HLA, Clotting Factor, von Willebrand factor, Transforming growth factor beta-induced, Severe Combined Immunodeficiency, SCID, HLA-A29, Uveitis, X-linked lymphoproliferative syndrome, SHD1A, XIAP, HLA-DQA1, HLA-DQB1, HLA-DQ2, HLA-DQ8, HLA-B27, Aplastic Anemia, Hemophilia A, Factor VIII, Hemophilia B, Factor IX, Fabry Disease, Thiopurine S-Methyltransferase, TPMT, Leiden, autoimmune disorder, Autosomal recessive, Birdshot chorioretinopathy, BSCR, C282Y, Celiac disease, Factor II Gene Mutation Analysis, Factor V Leiden, FISH, Genetic Testing, Hepatitis B, Hepatitis C, Hereditary Hemochromatosis, HLA haplotype testing, HLA I TYPING ALLELE HR, HLA-A29 Uveitis, HPV DNA, mutation, PCR (Polymerase Chain Reaction) Testing, Post – Intervention, Pre-Treatment, prevention, Prometheus, SCID, X-linked recessive inheritance