# SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

## **Drug Requested:** SYNAGIS® (palivizumab)

(RSV)

If approved, an authorization will be given for a specific number of injections, to be **ORDERED between October 1st and March 31st. RSV season for Virginia (as well as West Virginia, Maryland, DC, Delaware, and Pennsylvania) begins in late October and ends in April.** \*\*Typically, RSV season begins
November and ends in March. However, the duration of the Synagis season remains 5 consecutive months for all geographic areas in the United States.

Member Name:	
Member Sentara #:	Date of Birth:
Prescriber Name:	
	Date:
Office Contact Name:	
DI N I	For Number
Phone Number:	Fax Number:
NPI #:	
NPI #: DRUG INFORMATION: Author	
NPI #:	orization may be delayed if incomplete.
NPI #:	orization may be delayed if incomplete.
NPI #:	Derization may be delayed if incomplete.  Length of Therapy:  ICD Code, if applicable:
NPI #:	Derization may be delayed if incomplete.  Length of Therapy:  ICD Code, if applicable:

<u>Recommended Dosing</u>: Synagis<sup>®</sup> will be authorized according to the FDA recommended dose: Infants and Children <24 months - 15 mg/kg IM once monthly

(Continued on next page)

• Dosing Allowance: Synagis® is available in 50 mg and 100 mg vials. Due to the potential for significant waste, the following table should be utilized to determine the permitted dose (within 5% of calculated dose due to vial overfill) and vials to dispense. Any dosage increase must have corresponding weight charts and/or progress notes with current weight.

|--|

•	Synagis <sup>®</sup> has 1	been administered:	in an inpatient setting	: $\square$ Yes	s, date of last dose		No
---	----------------------------	--------------------	-------------------------	-----------------	----------------------	--	----

Weight-based Dose	<u>Dosage</u>	Dispense Units
0 - 3.5  kg	≤ 53 mg	1 vial of 50 mg/0.5mL
3.6 –7 kg	54 – 105 mg	1 vial of 100 mg/1ml
7.1 - 10.3  kg	106.5 – 154.5 mg	1 vial of 50 mg/0.5mL and 1 vial of 100 mg/1mL
10.4 - 13.6  kg	156 – 205 mg	2 vials of 100 mg/1mL
13.7 – 16.93 g	205.5 – 254 mg	1 vial of 50 mg/0.5mL and 2 vials of 100 mg/1mL
17 - 20.3  kg	255 – 305 mg	3 vials of 100 mg/1mL

- If Beyfortus<sup>™</sup> has been administered, please provide date & dose of administration:
- If Beyfortus<sup>™</sup> (nirsevimab) is administered, Synagis<sup>®</sup> (palivizumab) should <u>NOT</u> be administered later that season.

• If Enflonsia <sup>™</sup> h	nas been administered,	lease provide date & dose of administration	:
-------------------------------	------------------------	---	---

### Quantity Limit: 1 vial (50 mg/0.5 mL or 100 mg/1 mL) per 28 days

- Approval will be given for the current dosage and vial size(s). Throughout the RSV season, weight changes should be submitted on the Synagis request form when a different vial size(s) is/are required.
- Requests for doses exceeding the five (5) dose maximum or beyond the season end date will be **DENIED.**
- As defined by The National Respiratory and Enteric Virus Surveillance System (NREVSS): RSV season is over when virology is < 10% for 2 consecutive weeks.
- Monthly prophylaxis should be discontinued in any child who experiences a breakthrough RSV hospitalization.

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Diagnosis:	Preterm	Infants with	out Chronic	<b>Lung Disease</b>	e (CLD) of	Prematurity (	r
Congenital	Heart Di	sease (CHD)					

#### Please select ONE of the following:

- ☐ Infants without CLD or CHD born <28 weeks, 6 days and member's current age <12 months
- ☐ Infants without CLD or CHD born between  $\geq$  29 weeks to 31 weeks, 6 days and member's current age  $\leq$  6 months at start of (RSV) season

Provider please note: Infants without CLD or CHD born ≥ 32 weeks, 0 days' gestation: Synagis® is NOT RECOMMENDED.

□ Dia	agnosis: Preterm Infants with Chronic Lung Disease (CLD)
Please s	select ONE of the following:
	Infants with CLD <12 months (first year life) born <32 weeks, 0 days' gestation and require >21% supplemental O <sup>2</sup> for at least 28 days after birth
S	Infants with CLD <24 months and >12 months (second year life) born <32 weeks, 0 days' gestation with CLD of prematurity AND continued to require medical support during the 6-month period before the start of the second RSV season (must be verified by pharmacy paid claims and/or submitted chart notes):
	☐ Chronic systemic corticosteroid therapy: date of last use:
	☐ Diuretic therapy: date of last use:
	☐ Supplemental oxygen: date of last use:
	agnosis: Infants with Hemodynamically Significant Congenital Heart Disease HD)
Please s	select ONE of the following:
	Infants < 12 months old at the start of RSV season with hemodynamically significant heart disease defined by <b>ONE</b> of the following:
_	Acyanotic CHD, receiving treatment for congestive heart failure (CHF) and requires cardiac surgery  Moderate to severe pulmonary hypertension (PH, PAH)
	☐ Cyanotic CHD and Synagis is recommended by a pediatric cardiologist
	Infants in the first year or second year of life who are undergoing cardiac transplant or cardiac bypass <b>DURING</b> the RSV season
prophyl cardiac	er please note: For children who are receiving prophylaxis and who continue to require laxis after a surgical procedure, a post-operative dose of Synagis should be considered after bypass or at the conclusion of extra-corporeal membrane oxygenation for infants and children r than 24 months

venous return, Truncus arteriosus, Hypoplastic left heart syn	drome	
NON-APPROVABLE CARDIAC CONDITIONS		
Insignificant he modynamic he art disease (and the refore are NOT approvable indications):	Indications in which patients are NOT at an increased risk for RSV (and therefore are NOT approvable indications)	
Secundum atrial septal defect, small ventricular septal defect, pulmonic stenosis, uncomplicated aortic stenosis, mild coarctation of the aorta, patent ductus arteriosus	Lesions adequately corrected by surgery (unless the patient continues to require medications for CHF)     Mild cardiomyopathy who are NOT receiving medical therapy	

Examples of significant he modynamic cyanotic congenital he art disease:

Tetralogy of Fallot, Transportation of the great vessels, Ebstein's anomally, Tricuspid atresia, Total anomalous pulmonary

**EXAMPLES OF SIGNIFICANT AND APPROVABLE CARDIAC CONDITIONS** 

<ul> <li>Diagnosis: Children with Anatomic Pulmonary Abnormalities or Neuromuscular</li> <li>Disorder</li> </ul>
☐ Infants <12 months old (first year life) with a neuromuscular disorder(s) or congenital pulmonary anomaly that impairs the ability to clear secretions from upper airway
□ Provider has submitted name and ICD-10 code for anatomic pulmonary abnormality or neuromuscular disorder:
☐ Member must have <u>ONE</u> of the following:
<ul><li>pulmonary malformations</li></ul>
□ tracheoesophageal fistula
upper airway conditions
□ requires tracheostomy
□ Diagnosis: Immunocompromised Children
☐ Infants and children <24 months of age who are severely immunocompromised DURING the RSV season (i.e., receiving chemotherapy, undergoing solid organ or hematopoietic stem cell transplantation
EXAMPLES OF SEVERE IMMUNODEFICIENCIES/IMMUNOSUPPRESSION:
Advanced Acquired Immunodeficieny Syndrome (AIDS), Transplant, Chemotherapy, Severe Combined Immunodeficiency (SCID)
□ Diagnosis: Children with Cystic Fibrosis
Please select ONE of the following:
☐ Infants < 12 months old (first year of life) with Cystic Fibrosis with clinical evidence of CLD and/or nutritional compromise (e.g., requires total parenteral nutrition)
☐ Infants with Cystic Fibrosis <24 months and >12 months (second year life) with manifestations of severe lung disease (e.g., previous history of hospitalization for pulmonary exacerbation in the first year of life, abnormalities on chest x-ray or CT scan that persist when stable or patient weight for length is less than the 10 <sup>th</sup> percentile)
□ Authorization Criteria for additional dose(s) of Synagis. Check below all that apply. All criteria must be checked for approval. To support each line checked, all documentation (lab results, diagnostics, and/or chart notes) must be provided or request may be denied.
NOTE: For all requests received after March 31st – If all below conditions are met, the request will be

(Continued on next page)

be referred to a Medical Director for medical necessity review.

approved for an additional 1-month duration. **For all requests received prior to November** 1<sup>st</sup> – For members born between 32 to less than 35 weeks of gestation (without any significant medical conditions), if all below conditions are met, the request will only be approved for a maximum quantity of up to 3 doses. All other members will be approved for a quantity of 5 doses. If all the criteria below is **NOT** met, then the request will

	r requests to initiate treatment of Synagis prior to November 1 <sup>st</sup> :  Local virology data supplied from the National Respiratory & Enteric Virus Surveillance System (NREVSS) RSV Surveillance website OR recent surveillance data from a local/regional hospital (dated within < 14 days prior to the member's appointment) indicates an incidence of RSV greater than 10% (percent positive total antigen detection tests greater than 10%) for that locality Member meets the above stated criteria for their chronological and/or gestational age
Fo	requests to administer an additional dose of Synagis after March 31 <sup>st</sup> :  Member has <u>NOT</u> already received the maximum approvable five (5) doses of Synagis according to the member's chronological age, gestational age, and/or clinical situation  Local virology data supplied from the National Respiratory & Enteric Virus Surveillance System (NREVSS): RSV Surveillance website OR recent surveillance data from a local/regional hospital (dated within < 14 days prior to the member's appointment) indicates an incidence of RSV greater than 10% (percent positive total antigen detection tests greater than 10%) for that locality
lica	ation being provided by (check box below that applies):
Ph	nysician's office OR
	e of samples to initiate therapy does not meet step edit/preauthorization criteria.** <u>us therapies will be verified through pharmacy paid claims or submitted chart notes.</u> *
	Fo Blica