

SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization may be delayed.**

Prophylaxis Hereditary Angioedema (HAE) (Pharmacy)

Drug Requested (select applicable drug below):

| | |
|--|---|
| <input type="checkbox"/> Cinryze [®] (C1 Esterase Inhibitor Human) | <input type="checkbox"/> Haegarda [®] (C1 Esterase Inhibitor Human) |
| <input type="checkbox"/> Orladeyo [®] (berotralstat) | <input type="checkbox"/> Takhzyro [®] (lanadelumab) |

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code: _____

Weight: _____ Date: _____

Dosing Limit: (see below)

Haegarda: Administer 60 International Units (IU) per kg body weight twice weekly (every 3 or 4 days)

Quantity Limit:

Haegarda 2000 IU SDV kit: 16 kits per 28 days

Haegarda 3000 IU SDV kit: 8 kits per 28 days

Cinryze: Administer 1,000 units intravenous every 3 or 4 days

Quantity Limit:

Cinryze 500-unit vial: 16 vials per 30 days

(Continued on next page)

- ❑ **Orladeyo:** Adults and children ≥ 12 years of age - One capsule taken orally once daily with food
 - ❑ Quantity Limit:
 - ❑ 150 mg capsules: 1 capsule per day
 - ❑ 110 mg capsules: 1 capsule per day (For members with Moderate to severe impairment (Child-Pugh class B and C))
- ❑ **Takhzyro:** For children ≥ 12 years of age, adolescents, and adults – Administer 300 mg every 2 weeks. Dosing every 4 weeks for well controlled members (e.g., attack free) for > 6 months. The recommended dosage of Takhzyro in pediatric patients 2 to less than 6 years of age is 150 mg administered SC once every 4 weeks
 - ❑ Quantity Limit:
 - ❑ 300 mg vial/syringe per 14 days
 - ❑ 150 mg syringe per 28 days

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 12 months

Treatment of acute attacks of Hereditary Angioedema Criteria:

- ❑ Prescribed by or in consultation with a specialist in allergy, immunology, hematology, pulmonology or medical genetics

AND

- ❑ Provider attests the requested medication will **NOT** be used in combination with other prophylactic therapies targeting C1 inhibitors or kallikrein (e.g., Haegarda or Takhzyro)

AND

- ❑ Provider attests the patient is avoiding **BOTH** of the following possible triggers for HAE attacks:
 - ❑ Estrogen-containing oral contraceptive agents **AND** hormone replacement therapy
 - ❑ Antihypertensive agents containing ACE inhibitors

AND

- ❑ Member must meet medication specific age requirement:
 - ❑ Haegarda: Member must be at least 6 years of age
 - ❑ Cinryze: Member must be at least 6 years of age
 - ❑ Takhzyro: Member must be at least 2 years of age
 - ❑ Orladeyo: Member must be at least 12 years of age

AND

- ❑ Member has a history of **ONE** of the following:
 - ❑ **Three (3) or more** severe HAE attacks per month (i.e., airway swelling, debilitating cutaneous or gastrointestinal episodes)

(Continued on next page)

- ❑ Disablement for more than 5 days per month by HAE
- ❑ Recurrent laryngeal attacks caused by HAE

AND

- ❑ Treatment of member with “on-demand” therapy (i.e., Kalbitor[®], Firazyr[®], Ruconest[®], or Berinert[®]) did **NOT** provide satisfactory control or access to “on-demand therapy” is limited (**failure is defined as more than 5 attacks/month for 4 months consecutively within the same year**)

AND

- ❑ Member failed, is intolerant, or has a contraindication to attenuated (17 alpha-alkylated) androgens (e.g., Danazol[®]) for HAE prophylaxis

AND

- ❑ **Orladeyo (berotralstat) Criteria Only:**
 - ❑ Member has tried and failed the preferred prophylaxis HAE medication Takhzyro[®] (lanadelumab)

Member has ONE of the following clinical presentations that is consistent with a HAE subtype, confirmed by repeat blood testing (please submit chart notes for symptoms and lab values to confirm the HAE subtype):

II.A. ❑ HAE I: (all bullet points must apply)

- ❑ Low C1 inhibitor (C1-INH) antigenic level (C1-INH antigenic level below the lower limit of normal as defined by the laboratory performing the test)
- ❑ Low C4 level (C4 below the lower limit of normal as defined by the laboratory performing the test)
- ❑ Low C1-INH functional level (C1-INH functional level below the lower limit of normal as defined by the laboratory performing the test) **in addition to ONE** of the following:
 - ❑ Member has a family history of HAE
 - ❑ Acquired angioedema has been ruled out (i.e., patient onset of symptoms occur prior to 30 years old, normal C1q levels, patient does not have underlying disease such as lymphoma or benign monoclonal gammopathy [MGUS])

OR

II.B. ❑ HAE II (C1-Inhibitor dysfunction): (all bullet points must apply)

- ❑ Normal to elevated C1-INH antigenic level
- ❑ Low C4 level (C4 below the lower limit of normal as defined by the laboratory performing the test)
- ❑ Low C1-INH functional level (C1-INH functional level below the lower limit of normal as defined by the laboratory performing the test) **in addition to ONE** of the following:
 - ❑ Member has a family history of HAE
 - ❑ Acquired angioedema has been ruled out (i.e., onset of symptoms occurred prior to 30 years old, normal C1q levels, patient does not have underlying disease such as lymphoma or benign monoclonal gammopathy [MGUS])

(Continued on next page)

OR

II.C. HAE III with normal C1-INH: (all bullet points must apply)

- Normal C1-INH antigenic level
- Normal C4 level
- Normal C1-INH functional level
- Repeat blood testing during an attack has confirmed the member does **NOT** have abnormal lab values indicative of HAE I or HAE II
- Member had an inadequate response or intolerance to an adequate trial of prophylactic therapy with **ONE** of following:
 - antifibrinolytic agent: (tranexamic acid (TXA) **OR** aminocaproic acid)
 - 17 α - alkylated androgen: danazol
 - progestins (female members only)

AND

- ONE of the following:**
 - Member has a known HAE-causing mutation (e.g., mutation of coagulation factor XII gene [F12 mutation], mutation in the angiotensin-converting enzyme gene, mutation in the plasminogen gene or kininogen-1)
 - Member has a family history of HAE and documented evidence of lack of efficacy of chronic high-dose antihistamine therapy (e.g., cetirizine standard dosing at up to four times daily or an alternative equivalent, given for at least one month or an interval long enough to expect three or more angioedema attacks) **AND** corticosteroids

Reauthorization: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- Member must continue to meet initial criteria

AND

- Significant improvement in severity and duration of attacks has been achieved and sustained

AND

- Member has experienced an absence of unacceptable toxicity from the drug (e.g., hypersensitivity reactions)

AND

- For Takzyro Renewal Only: Has the member been attack free for greater than > 6 months?
 - Yes- approve dosing every 4 weeks
- No- approve dosing every 2 weeks

(Continued on next page)

Medication being provided by a Specialty Pharmacy - PropriumRx

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****

*Approved by Pharmacy and Therapeutics Committee: 9/19/2013

REVISED/UPDATED/REFORMATTED: 7/30/2017; 9/13/2017; 11/16/2017; 3/31/2018; 5/18/2018; 12/11/2018; (Reformatted) 3/16/2019; 7/6/2019; 9/20/2019;
4/1/2021; 6/30/2021; 4/7/2023; 11/20/2023