

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

Drug Requested: Non-Preferred Ustekinumab products (**PHARMACY BENEFIT ONLY**)
(ustekinumab SQ therapy is Self-Administered by member)

PREFERRED		
<input type="checkbox"/> Pyzchiva® (ustekinumab-ttwe) syringe/vial (Requires trial and failure of a preferred TNF-alpha inhibitor)		
NON-PREFERRED		
<input type="checkbox"/> Imuldosa™ (ustekinumab-srlf)	<input type="checkbox"/> Otulfi® (ustekinumab-aauz)	<input type="checkbox"/> Selarsdi® (ustekinumab-aekn)
<input type="checkbox"/> Steqeyma® (ustekinumab-stba)	<input type="checkbox"/> ustekinumab (generic Stelara®)	<input type="checkbox"/> ustekinumab-aekn (generic Selarsdi®)
<input type="checkbox"/> ustekinumab-ttwe (generic Pyzchiva®)	<input type="checkbox"/> Wezlana™ (ustekinumab-kfce)	<input type="checkbox"/> Yesintek™ (ustekinumab-kfce)
<input type="checkbox"/> Stelara® (ustekinumab) *see additional criteria for approval		

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

(Continued on next page)

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ **Length of Therapy:** _____

Diagnosis: _____ **ICD Code, if applicable:** _____

Weight (if applicable): _____ **Date weight obtained:** _____

Recommended Dose:

Indication	Dosage:
Adults with Moderate to Severe Chronic Plaque Psoriasis	<p>Weight</p> <ul style="list-style-type: none"> Less than or = 100 kg Initial (two 45 mg prefilled syringe/ 28 days) then continue with one 45 mg prefilled syringe/84 days Great than 100 kg Two 90 mg administered prefilled syringe/ 28 days then, one 90 mg administered prefilled syringe/ 84 days
Pediatric patients (6 years or older) with Moderate to Severe Chronic Plaque Psoriasis	<ul style="list-style-type: none"> <60 kg (0.75 mg/kg): Initial (two 45 mg prefilled syringe/28days) then one 45mg/84 days ≥ 60 to ≤ 100 kg: Initial (two 45 mg prefilled syringe/ 28 days) then one 45/ 84 days >100 kg: two 90 mg administered prefilled syringe/ 28 days, then one 90 mg/84 days
Active Psoriatic Arthritis	<ul style="list-style-type: none"> <60 kg (0.75 mg/kg): Initial (two 45 mg prefilled syringe/28days) then one 45mg/84 days ≥ 60 to ≤ 100 kg: Initial (two 45 mg prefilled syringe/ 28 days) then one 45/ 84 days >100 kg: two 90 mg administered prefilled syringe/ 28 days, then one 90 mg/84 days
Moderately to Severe Active Crohn's Disease	<ul style="list-style-type: none"> A single intravenous infusion using weight-based dosing Up to 55 kg 260 mg (2 vials) Greater than 55 kg to 85 kg 390 mg (3 vials) Greater than 85 kg 520 mg (4 vials) After initial IV dose: 90mg syringe every 56 days
Adult patient with Moderately to Severely Active Ulcerative Colitis	<ul style="list-style-type: none"> A single intravenous infusion using weight-based dosing Up to 55 kg 260 mg (2 vials) Greater than 55 kg to 85 kg 390 mg (3 vials) Greater than 85 kg 520 mg (4 vials) After initial IV dose: 90mg syringe every 56 days

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

(Continued on next page)

Diagnosis: Check below diagnosis that applies to qualify for approval or authorization may be delayed.

DIAGNOSIS: Adults with Active Psoriatic Arthritis

- Member has a diagnosis of moderate to severe psoriatic arthritis
- Member has tried and failed **ONE (1)** of the preferred drugs below:

<input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima® (adalimumab-bwwd)	<input type="checkbox"/> Enbrel®
---	----------------------------------
- Member has tried and failed Pyzchiva® syringe/vial (requires trial and failure of a preferred TNF-alpha inhibitor)

Diagnosis: Pediatric members (6 years and older) with psoriatic arthritis

- Member is 6 years or older
- Member has a diagnosis of psoriatic arthritis
- Member has tried and failed **ONE (1)** of the preferred drugs below:

<input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima® (adalimumab-bwwd)	<input type="checkbox"/> Enbrel®
---	----------------------------------
- Member has tried and failed Pyzchiva® syringe/vial (requires trial and failure of a preferred TNF-alpha inhibitor)

Diagnosis: Moderate to Severe Chronic Plaque Psoriasis

- Member is 6 years or older
- Member has a diagnosis of moderate to severe plaque psoriasis who are candidates for phototherapy or systemic therapy
- Member has tried and failed **ONE (1)** of the preferred drugs below:

<input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima® (adalimumab-bwwd)	<input type="checkbox"/> Enbrel®
---	----------------------------------
- Member has tried and failed Pyzchiva® syringe/vial (requires trial and failure of a preferred TNF-alpha inhibitor)

Diagnosis: Ulcerative Colitis

- Member has a diagnosis of moderate to severe active disease
- Trial and failure of **BOTH** of the preferred drugs below:

<input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima® (adalimumab-bwwd)	<input type="checkbox"/> Pyzchiva® syringe/vial (Requires trial and failure of a preferred TNF-alpha inhibitor)
---	---

(Continued on next page)

Diagnosis: Crohn's Disease

- Member has a diagnosis of moderate to severe active disease
- Trial and failure of **BOTH** of the preferred drugs below:

<input type="checkbox"/> adalimumab-adbm (Boehringer Ingelheim) OR Hadlima® (adalimumab-bwwd)	<input type="checkbox"/> Pyzchiva® syringe/vial (Requires trial and failure of a preferred TNF-alpha inhibitor)
--	---

***For Brand Stelara approval**

- Member has trial and failure of **ONE** preferred biosimilar agents for Humira:
 - adalimumab-adbm (Boehringer Ingelheim)
 - Hadlima® (adalimumab-bwwd)
- Member has tried and failed Pyzchiva® syringe/vial
- FDA Medwatch** form must be completed for the preferred ustekinumab biosimilar product, Pyzchiva® syringe/vial and a copy of submitted MedWatch form must be attached for approval consideration.

Induction Dose (If required) – Single IV induction dose

Authorization Criteria: To be reviewed for one-time approval under the medical benefit

- Medication will be used as induction therapy
- Medication being provided by:
- Location/site of drug administration:** _____
- NPI or DEA # of administering location:** _____
- Select **ONE** of the following one-time doses to be administered based on member's weight
 - ≤55 kg: 260 mg as single dose; 260 mg = 260 billable units
 - >55 kg to 85 kg: 390 mg as single dose; 390 mg = 390 billable units
 - >85 kg: 520 mg as single dose; 520 mg = 520 mg billable units

Medication being provided by Specialty Pharmacy - PropriumRx

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****
****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****