

# SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

**Drug Requested:** Opzelura™ (ruxolitinib) (Non-Preferred)

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

DEA OR NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight: \_\_\_\_\_ Date: \_\_\_\_\_

**Quantity Limit:** 60 grams per week and 8-week limit

**Other Indications:** Opzelura cream will not be approved for the indication of nonsegmental vitiligo in adult and pediatric patients  $\geq 12$  years old

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Length of Authorization: 1 year**

- Member must be 12 years of age or older and have an FDA-approved diagnosis for mild to moderate Atopic Dermatitis

**AND**

(Continued on next page)

- ❑ Prior documented trial and failure of 8 weeks for each trial (or contraindication) of:
  - ❑ One (1) topical corticosteroid of medium to high potency (e.g., mometasone, triamcinolone)  
**AND**
  - ❑ One (1) topical calcineurin inhibitors (tacrolimus or pimecrolimus)  
**AND**
  - ❑ Trial and failure of Eucrisa™  
**AND**
  - ❑ Trial and failure of Dupixent® (dupilumab)  
**AND**
  - ❑ Trial and failure of Adbry® (tralokinumab)

**Medication being provided by Specialty Pharmacy - PropriumRx**

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****