## SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

**Drug Requested: Cinqair® IV** (reslizumab) (Pharmacy) (Non-Preferred)

MEMBER & PRESCRIBER INF	FORMATION: Authorization may be delayed if incomplete.
Member Name:	
	Date of Birth:
Prescriber Name:	
	Date:
Office Contact Name:	
Phone Number:	
NPI #:	
DRUG INFORMATION: Authoriz	
	Length of Therapy:
	ICD Code, if applicable:
Weight (if applicable):	Date weight obtained:
Recommended Dosage: Dosage 3mg *The Health Plan considers the us	kg once every 4 weeks by intravenous infusion over 20 - 50 minute e of concomitant therapy with Cinqair®, Dupixent®,
and efficacy of these combinations	l Xolair <sup>®</sup> to be experimental and investigational. Safety s have <u>NOT</u> been established and will <u>NOT</u> be permitted ve Dupixent <sup>®</sup> , Fasenra <sup>®</sup> , Nucala <sup>®</sup> , Tezspire <sup>™</sup> or Xolair <sup>®</sup>

(Continued on next page)

authorization on file, all subsequent requests for Cinqair® will NOT be approved.

(Continued from previous page)

provi	ded or request may be denied. (Trials will be verified using pharmacy claims and/or submitted t notes.)
<u>Initi</u>	al Authorization: 6 months
1.	Has the member been approved for Cinqair <sup>®</sup> previously through Sentara medical department?  ☐ Yes ☐ No
2.	Is the member 18 years of age or older?  ☐ Yes ☐ No
3.	Does the member have a diagnosis of severe* asthma?  □ Yes □ No
4.	Does the member have asthma with an eosinophilic phenotype defined as blood eosinophils $\geq$ 400 cells/ $\mu$ L?
5.	
	□ Yes □ No
6.	Will Cinqair® be used for add on maintenance treatment in members regularly receiving <b>both</b> of the following:
	Medium to high dose inhaled corticosteroids AND
	• An additional controller medication (i.e. long-acting beta agonist, leukotriene modifiers)?
	□ Yes □ No
7.	Has the member had two or more exacerbations in the previous year requiring oral or injectable corticosteroid treatment (in addition to the regular maintenance therapy defined above) <b>OR</b> one exacerbation resulting in a hospitalization?
	□ Yes □ No
0	

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To

- 8. Does the member have at least one of the following for assessment of clinical status:
  - Use of systemic corticosteroids
  - Use of inhaled corticosteroids
  - Number of hospitalizations, ER visits, or unscheduled visits to healthcare provider due to condition
  - Forced expiratory volume in 1 second (FEV<sub>1</sub>)?
  - □ Yes □ No
- 9. Has the member tried and failed an adequate trial of the 2 different preferred products (Fasenra® and Xolair®)?

□ Yes □ No

(Continued on next page)

Reauthorization: 12 months Check below all that apply. All criteria must be met for approval. To
support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be
provided or request may be denied. (Trials will be verified using pharmacy claims and/or submitted
chart notes.)

1.	Has the member been assessed for toxicity?	
	□ Yes □ No	
2.	2. Does the member have improvement in asthma symptoms or asthma exacerbations as evidenced by decrease in one or more of the following:	
	Use of systemic corticosteroids	
	• Hospitalizations	
	• ER visits	

- Unscheduled visits to healthcare provider
- Improvement from baseline in forced expiratory volume in 1 second (FEV1)?
- □ Yes □ No

## \*Components of severity for classifying asthma as *severe* may include any of the following (not all-inclusive):

- Asthma that remains uncontrolled despite optimized treatment with high-dose ICS-LABA
- Asthma that requires high-dose ICS-LABA to prevent it from being uncontrolled
- Symptoms throughout the day
- Nighttime awakenings, often 7 times per week
- SABA use for symptom control occurs several times per day
- Extremely limited normal activities
- Lung function (percent predicted FEV<sub>1</sub>) < 60%
- Exacerbations requiring oral systemic corticosteroids are generally more frequent and intense relative to moderate asthma.

## Medication being provided by Specialty Pharmacy - PropriumRx

\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. \*\*

\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. \*