

# SENTARA COMMUNITY PLAN (MEDICAID)

## MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-844-305-2331.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization can be delayed.**

**Drug Requested:** Fasenra<sup>®</sup> SQ (benralizumab) (J0517) (Medical)

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Name/Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight (if applicable): \_\_\_\_\_ Date weight obtained: \_\_\_\_\_

Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

- Fasenra<sup>®</sup> 10mg/0.5ml single prefilled syringe and pen= 10 billable units
- Fasenra<sup>®</sup> 30mg/ml single prefilled syringe and pen= 30 billable units

### **Recommended Dosing:**

**Asthma, severe eosinophilic:**

#### **Adult and Adolescent Patients 12 Years of Age and Older:**

- 30 mg every 4 weeks for the first 3 doses followed by once every 8 weeks thereafter

#### **Pediatric Patients 6 Years to 11 Years of Age:**

- Weighing Less Than 35 kg: the recommended dosage is 10 mg every 4 weeks for first 3 doses followed by once every 8 weeks thereafter
- Weighing 35 kg or More: the recommended dosage is 30 mg every 4 weeks for first 3 doses followed by once every 8 weeks thereafter

**Eosinophilic granulomatosis with polyangiitis (EGPA):** 30 mg every 4 weeks

(Continued on next page)

**\*The Health Plan considers the use of concomitant therapy with Cinqair<sup>®</sup>, Dupixent<sup>®</sup>, Nucala<sup>®</sup>, Tezspire<sup>™</sup> and Xolair<sup>®</sup> to be experimental and investigational. Safety and efficacy of these combinations have **NOT** been established and will **NOT** be permitted. In the event a member has an active Cinqair<sup>®</sup>, Dupixent<sup>®</sup>, Nucala<sup>®</sup>, Tezspire<sup>™</sup> or Xolair<sup>®</sup> authorization on file, all subsequent requests for Fasenra<sup>®</sup> will **NOT** be approved.**

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Diagnosis: Asthma, severe eosinophilic**

**Initial Authorization: 6 months**

1. Has the member been approved for Fasenra<sup>®</sup> previously through the Sentara pharmacy department?  
 Yes  No
2. Is the member 6 years of age or older?  
 Yes  No
3. Does the member have a diagnosis of severe\* asthma?  
 Yes  No
4. Does the member have asthma with an eosinophilic phenotype defined as blood eosinophils  $\geq 150$  cells/ $\mu$ L?  
 Yes  No
5. Will coadministration with another monoclonal antibody be avoided (i.e. omalizumab, mepolizumab, reslizumab, dupilumab, tezepelumab-ekko)?  
 Yes  No
6. Will Fasenra<sup>®</sup> be used for add-on maintenance treatment in members regularly receiving both (unless otherwise contraindicated) of the following:
  - Medium to high dose inhaled corticosteroids **AND**
  - An additional controller medication (i.e. long-acting beta agonist, leukotriene modifiers)? Yes  No
7. Has the member had two or more exacerbations in the previous year requiring oral or injectable corticosteroid treatment (in addition to the regular maintenance therapy defined above) **or** one exacerbation resulting in hospitalization?  
 Yes  No

(Continued on next page)

- Does the member have at least one of the following for assessment of clinical status:
- Use of systemic corticosteroids
  - Use of inhaled corticosteroids
  - Number of hospitalizations, ER visits, or unscheduled visits to healthcare provider due to condition forced expiratory volume in 1 second (FEV1)?
- Yes    No

**Diagnosis: Asthma, severe eosinophilic**

**Reauthorization: 12 months.** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

1. Has the member been assessed for toxicity?  
 Yes    No
2. Does the member have improvement in asthma symptoms or asthma exacerbations as evidenced by a decrease in one or more of the following:
  - Use of systemic corticosteroids
  - Hospitalizations
  - ER visits
  - Unscheduled visits to healthcare provider
  - Improvement from baseline in forced expiratory volume in 1 second (FEV1)? Yes    No

**Diagnosis: Eosinophilic Granulomatosis Polyangiitis (EGPA)**

**Initial Authorization: 6 months**

1. Has the member been approved for Fasenra<sup>®</sup> previously through the Sentara pharmacy department?  
 Yes    No
2. Is the member 18 years of age and older?  
 Yes    No
3. Does the member have a diagnosis of EGPA (aka Churg-Strauss Syndrome)?  
 Yes    No
4. Member has blood eosinophils greater than or equal to 1000 cells/ $\mu$ L or greater than 10% of leukocytes  
 Yes    No

(Continued on next page)

5. Member is currently on maximally tolerated oral corticosteroid therapy or has an intolerance, hypersensitivity or contraindication to oral corticosteroid therapy  
 Yes  No
6. Member's physician has assessed baseline disease severity utilizing an objective measure/tool (e.g., Birmingham Vasculitis Activity Score [BVAS], history of asthma symptoms and/or exacerbations, duration of remission, rate of relapses).  
 Yes  No

**Diagnosis: Eosinophilic Granulomatosis Polyangiitis (EGPA)**

**Reauthorization: 12 months.** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

1. Member has been assessed for toxicity  
 Yes  No
2. Member has disease response as indicated by improvement in signs and symptoms compared to baseline as evidenced in one or more of the following:
- Member is in remission [defined as a Birmingham Vasculitis Activity Score (BVAS) score=0 and a prednisone/prednisolone daily dose of  $\leq 4$  mg]
  - Decrease maintenance dose of systemic corticosteroids
  - Improvement in BVAS score compared to baseline
  - Improvement in asthma symptoms or asthma exacerbations
  - Improvement in duration of remission or decrease in the rate of relapses

**\*Components of severity for classifying asthma as *severe* may include any of the following (not all inclusive):**

- Asthma that remains uncontrolled despite optimized treatment with high-dose ICS-LABA
- Asthma that requires high-dose ICS-LABA to prevent it from being uncontrolled
- Symptoms throughout the day
- Nighttime awakenings, often 7 times per week
- SABA use for symptom control occurs several times per day
- Extremely limited normal activities
- Lung function (percent predicted FEV1)  $< 60\%$
- Exacerbations requiring oral systemic corticosteroids are generally more frequent and intense relative to moderate asthma

(Continued on next page)

**Eosinophilic Granulomatosis Polyangiitis (EGPA) is defined as all of the following:**

- History or presence of asthma
- Blood eosinophil level > 10% or an absolute count > 1000 cells/mm<sup>3</sup>
- Two or more of the following criteria:
- Histopathologic evidence of eosinophilic vasculitis, perivascular eosinophilic infiltration, or eosinophil rich granulomatous inflammation
  - Neuropathy
  - Pulmonary infiltrates
  - Sinonasal abnormalities
  - Cardiomyopathy
  - Glomerulonephritis
  - Alveolar hemorrhage
  - Palpable purpura
  - Antineutrophil Cytoplasmic Antibody (ANCA) positivity

**Medication being provided by: Please check applicable box below.**

- Location/site of drug administration: \_\_\_\_\_  
NPI or DEA # of administering location: \_\_\_\_\_
- OR**
- Specialty Pharmacy – PropriumRx

For urgent reviews: Practitioner should call Sentara Health Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****