

SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Drug Requested: Skyclarys[®] (omaveloxolone)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

Quantity Limit: 3 capsules per day

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 12 months

- Member is 16 years of age or older
- Member has a diagnosis of Friedreich's ataxia as established by molecular genetic testing and detection of biallelic pathogenic variants in the FXN gene (**submit documentation**)
- Prescribed by or in consultation with a Neurologist, Geneticist or Physician who specializes in the treatment of Friedreich's ataxia
- Member exhibits clinical signs and symptoms of disease (e.g., ataxia, speech disturbance, sensory dysfunction muscle weakness, decline in coordination, frequent falling) that are consistent with Friedreich's ataxia

(Continued on next page)

- Member has a baseline modified Friedreich Ataxia Rating Scale (mFARS) score between 20-80 (if score is below 20 please send genetic test to document member is **NOT** a carrier): _____ (submit score)
- Member's B- Type natriuretic Peptide (BNP) is ≤ 200 pg/mL prior to initiating therapy and will be monitored periodically during treatment
- Member must **NOT** have uncontrolled diabetes (i.e., HbA1c $\geq 11\%$)
- Member will avoid concomitant therapy with the following:
 - Strong or moderate CYP3A4 inhibitors (e.g., fluconazole, itraconazole) [**NOTE: If therapy is unavoidable, members will be monitored closely for adverse reaction and/or dose modifications will be implemented**]
 - Strong and moderate CYP3A4 inducers (e.g., rifampin, carbamazepine, St. John's Wort)
- Member does **NOT** have a history of clinically significant left-sided heart disease and/or clinically significant cardiac disease (**NOTE: Excludes mild to moderate cardiomyopathy associated with Friedreich's ataxia; submit chart notes and/or lab documentation**)
- Member retains meaningful voluntary motor function (e.g., manipulate objects using upper extremities, ambulates)

Reauthorization: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- Member's modified Friedreichs Ataxia Rating scale (mFARS) score has shown an improvement or has demonstrated slowed progression of disease from baseline (e.g., bulbar function, upper/lower limb coordination, upright stability): _____ (submit score and/or clinical documentation obtained within the last 30 days)
- Member continues to avoid concomitant therapy with strong or moderate CYP3A4 inhibitors s (e.g., fluconazole, itraconazole) **AND** strong and moderate CYP3A4 inducers (e.g., rifampin, carbamazepine, St. John's Wort)
- Member does **NOT** have a history of clinically significant left-sided heart disease and/or clinically significant cardiac disease (**NOTE: Excludes mild to moderate cardiomyopathy associated with Friedreich's ataxia; submit chart notes and/or lab documentation**)
- Member retains meaningful voluntary motor function (e.g., manipulate objects using upper extremities, ambulates)

Medication being provided by a Specialty Pharmacy – Proprium Rx

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****
****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****