

SENTARA COMMUNITY PLAN (MEDICAID)

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-844-305-2331. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If information provided is not complete, correct, or legible, authorization can be delayed.

Drug Requested: Empaveli® (pegcetacoplan) SQ (J3490) (Medical)
Paroxysmal Nocturnal Hemoglobinuria (PNH)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____
Member Sentara #: _____ Date of Birth: _____
Prescriber Name: _____
Prescriber Signature: _____ Date: _____
Office Contact Name: _____
Phone Number: _____ Fax Number: _____
DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____
Dosing Schedule: _____ Length of Therapy: _____
Diagnosis: _____ ICD Code, if applicable: _____
Weight: _____ Date: _____

☐ Standard Review. In checking this box, the timeframe does not jeopardize the life or health of the member or the member's ability to regain maximum function and would not subject the member to severe pain.

Max quantity limits:

- 8 (eight) SQ infusions every 28 days
- Empaveli® 1080 mg/20 mL solution in single-use vials for injection supplied in 8-count cartons

Recommended Dosage:

- Maintenance – 1080 mg twice weekly
- Dosage Adjustment: For lactate dehydrogenase (LDH) levels > 2 levels ULN, adjust pegcetacoplan dosing regimen to 1080 mg every 3 days. Monitor LDH twice weekly for at least 4 weeks after a dose increase.

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

- ☐ Medication must be prescribed by or in consultation with a hematologist or nephrologist
- ☐ Prescriber must be enrolled in the Empaveli[®] Risk Evaluation and Mitigation Strategy (REMS) program
- ☐ Member must be 18 years of age or older
- ☐ Member must meet **ONE** of the following:
 - ☐ Empaveli[®] will be used as switch therapy **AND** member meets **ALL** the following:
 - ☐ Member failed Soliris[®] or Ultomiris[®] and must meet renewal criteria
 - ☐ Member does **NOT** have a systemic infection
 - ☐ Member must be vaccinated against encapsulated bacteria (*Streptococcus pneumoniae*, *Neisseria meningitidis*, and *Haemophilus influenzae type B*) **at least two weeks prior** to initiation of Empaveli[®] therapy and revaccinated according to current medical guidelines for vaccine use
 - ☐ Empaveli[®] will **NOT** be used in combination with other complement inhibitor therapies (e.g., Ultomiris[®], Soliris[®] or Fabhalta[®])

OR

- ☐ Member is treatment-naïve **AND** member meets **ALL** the following:
 - ☐ Member must have a diagnosis of Paroxysmal Nocturnal Hemoglobinuria (PNH) confirmed by detection of PNH clones of at least 10% by flow cytometry testing (**must submit labs**)
 - ☐ Flow cytometry pathology report must demonstrate at least two (2) different glycosylphosphatidylinositol (GPI) protein deficiencies (e.g., CD55, CD59, etc.) within two (2) different cell lines from granulocytes, monocytes, erythrocytes (**must submit labs**)
 - ☐ Member has laboratory evidence of significant intravascular hemolysis (i.e. LDH $\geq 1.5 \times$ ULN) **AND** has experienced **ONE** of the following additional indications for therapy (**must submit chart notes and labs**):
 - ☐ Member is transfusion dependent (defined by having a transfusion within the last 12 months) and symptomatic anemia
 - ☐ Presence of a thrombotic event (e.g., DVT, PE)
 - ☐ Presence of organ damage secondary to chronic hemolysis (i.e. renal insufficiency, pulmonary insufficiency, or hypertension)
 - ☐ Member is pregnant and potential benefit outweighs potential fetal risk
 - ☐ Member has abdominal pain requiring admission to hospital
- ☐ Member does **NOT** have a systemic infection
- ☐ Member must be administered a meningococcal vaccine **at least two weeks prior** to initiation of Empaveli[®] therapy and revaccinated according to current medical guidelines for vaccine use
- ☐ Empaveli[®] will **NOT** be used in combination with other complement inhibitor therapies (e.g., Ultomiris[®], Soliris[®] or Fabhalta[®])

(Continued on next page)

Reauthorization: 6 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ☐ Provider attests to an absence of unacceptable toxicity from the drug (e.g. serious meningococcal infections [septicemia and/or meningitis], infusion reactions)
- ☐ Member has experienced positive disease response indicated by at least **ONE** of the following (**check all that apply; results must be submitted to document improvement**):
 - ☐ Decrease in serum LDH
 - ☐ Stabilization/increase in hemoglobin level
 - ☐ Decrease in packed RBC transfusion requirement
 - ☐ Reduction in thromboembolic events

Medication being provided by (check box below that applies):

- ☐ Physician's office **OR** ☐ Specialty Pharmacy – Proprium Rx

For urgent reviews: Practitioner should call Sentara Health Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****