SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request</u>. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

| Drug Requested: (select ONE drug below) | |
|---|--|
| □ Cuvrior [™] (trientine tetrahydrocholoride) 300 mg tablets | □ Syprine® (trientine) 250 mg capsules |
| □ trientine 250 mg capsules | □ trientine 500 mg capsules |
| MEMBER & PRESCRIBER INFORMAT | TION: Authorization may be delayed if incomplete. |
| Member Name: | |
| Member Sentara #: | |
| Prescriber Name: | |
| Prescriber Signature: | |
| Office Contact Name: | |
| Phone Number: Fax Number: | |
| DEA OR NPI #: | |
| DRUG INFORMATION: Authorization may | be delayed if incomplete. |
| Drug Form/Strength: | |
| | Length of Therapy: |
| Diagnosis: | ICD Code, if applicable: |
| Weight: | |
| Quantity Limits: | |
| ☐ Cuvrior: 3,000 mg (10 tablets) per day | |
| ☐ trientine (all formulations): | |
| • Age > 12 years: 2,000 mg per day | |
| • Age \leq 12 years: 1,500 mg per day | |

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

| | Member must meet ONE of the following age requirements: |
|-------|--|
| | \square For Cuvrior requests: Member is ≥ 18 years of age |
| | \square For trientine requests (all formulations): Member is ≥ 6 years of age |
| | Medication must be prescribed by or in consultation with a gastroenterologist or hepatologist |
| | Member has a diagnosis of Wilson's disease |
| | Member's diagnosis of Wilson's disease has been confirmed by at least <u>TWO</u> of the following (submit labs or chart notes for documentation; check all that apply): |
| | ☐ Presence of Kayser-Fleisher rings |
| | ☐ Serum ceruloplasmin (CPN) < 20 mg/dL |
| | □ 24-hour urine copper > 40 mcg |
| | ☐ Liver biopsy with copper dry weight > 250 mcg/g |
| | Member has tried and failed generic penicillamine *requires prior authorization* at up to maximally indicated doses or clinically significant adverse effects are experienced (must submit completed MedWatch form and chart notes to document adverse event and/or treatment failure with penicillamine) |
| | For Cuvrior [™] or Brand Syprine [®] requests: Member has tried and failed generic trientine (*requires prior authorization*) at up to maximally indicated doses or clinically significant adverse effects are experienced (must submit completed MedWatch form and chart notes to document adverse event and/or treatment failure with trientine) |
| | For Cuvrior [™] requests <u>ALL</u> the following criteria must be met: |
| | ☐ Member is de-coppered [i.e., serum non-ceruloplasmin copper (NCC) level ≥ 25 and ≤ 150 mcg/L] |
| | ☐ Member is tolerant to penicillamine |
| | ☐ Member will discontinue penicillamine prior to initiating therapy with Cuvrior [™] |
| | Member's serum or urinary copper levels will be monitored during therapy along with LFT's, CBC, INR, serum non-ceruloplasmin bound copper plus monitoring for skin changes and fever during the first month of therapy |
| suppo | uthorization: 12 months. Check below all that apply. All criteria must be met for approval. To ort each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be ded or request may be denied. |
| | Member has experienced a positive response to therapy as demonstrated by ONE of the following: |
| | ☐ Member's serum copper level is maintained at <10 mcg free copper/dL of serum (submit current lab level for documentation) |
| | ☐ Member's urinary copper excretion is maintained at 200-500 mcg (3-8 micromoles) per day on 24-hour urinary copper assessment (submit current lab level for documentation) |
| | Member's serum or urinary copper levels will continue to be monitored during therapy along with LFT's, CBC, INR and serum non-ceruloplasmin bound copper |

Medication being provided by Specialty Pharmacy – Proprium Rx

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *