

# SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

**Drug Requested:** **Xeljanz<sup>®</sup>** (tofacitinib)/**Xeljanz<sup>®</sup> XR<sup>®</sup>** (tofacitinib xr) **(Non-Preferred)**

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Name/Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight (if applicable): \_\_\_\_\_ Date weight obtained: \_\_\_\_\_

### **Recommended Dose:**

Indication:	Dosage:
<b>Moderate to Severe Active Rheumatoid Arthritis</b>	<ul style="list-style-type: none"><li>• Xeljanz 5 mg twice daily (60 tabs/30 days) or</li><li>• Xeljanz XR 11 mg once daily (30 tabs/30 days)</li><li>• Patients with moderate and severe renal impairment or moderate hepatic impairment is Xeljanz 5 mg once daily (30 tabs/30 days)</li></ul>
<b>Psoriatic Arthritis (in combination with nonbiologic DMARDs)</b>	<ul style="list-style-type: none"><li>• Xeljanz 5 mg twice daily (60 tabs/30 days) or Xeljanz XR 11 mg once daily (30 tabs/30 days)</li><li>• Patients with moderate and severe renal impairment or moderate hepatic impairment is Xeljanz 5 mg once daily</li></ul>
<b>Polyarticular Course Juvenile Idiopathic Arthritis (pcJIA) (≥ 2 years)</b>	<ul style="list-style-type: none"><li>• Xeljanz 5 mg twice daily (60 tabs/30 days) or weight-based equivalent twice daily</li><li>• Xeljanz Oral Solution 5 mg twice daily (300ml/30 days) or weight-based equivalent twice daily</li></ul>

(Continued on next page)

Indication:	Dosage:
<b>Ulcerative Colitis</b>	<ul style="list-style-type: none"> <li>• Induction: Xeljanz 10 mg twice daily (60 tabs/30 days) or Xeljanz XR 22 (30 tabs/30 days) mg once daily for 8 weeks; evaluate patients and transition to maintenance therapy depending on therapeutic response. If needed, continue Xeljanz 10 mg twice daily (60 tabs/30 days) or Xeljanz XR 22 mg (30 tabs/30 days) once daily for a maximum of 16 weeks. Discontinue Xeljanz 10 mg twice daily or Xeljanz XR 22 mg once daily (30 tabs/30 days) after 16 weeks if adequate therapeutic response is not achieved.</li> <li>• Maintenance: Xeljanz 5 mg twice daily (60 tabs/30 days) or Xeljanz XR 11 mg once daily (30 tabs/30 days). For patients with loss of response during maintenance treatment, Xeljanz 10 mg twice daily or Xeljanz XR 22 mg once daily (30 tabs/30 days) may be considered and limited to the shortest duration, with careful consideration of the benefits and risks for the individual patient. Use the lowest effective dose needed to maintain response</li> </ul>
<b>Active Ankylosing Spondylitis</b>	<ul style="list-style-type: none"> <li>• Xeljanz 5mg twice daily (60 tabs/30 days) or Xeljanz XR 11mg once daily (30 tabs/30 days)</li> <li>• Patients with moderate and severe renal impairment or moderate hepatic impairment Xeljanz 5mg once daily (30 tabs/30 days)</li> </ul>

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**DIAGNOSIS:** (Check one of the diagnoses below to ensure authorization will not be delayed.)

☐ **Rheumatoid Arthritis – Moderate to Severe**

- ☐ Member is 18 years of age or older
- ☐ Trial and failure of, contraindication, or adverse reaction to methotrexate
- ☐ Trial and failure of at least **ONE (1) other DMARD** therapy (**check each tried**):

<input type="checkbox"/> auranofin	<input type="checkbox"/> azathioprine
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> Other: _____

- ☐ Patient has tried and failed **TWO (2)** of the preferred drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

(Continued on next page)

### ☐ Psoriatic Arthritis

- ☐ Member is 18 years of age or older
- ☐ Trial and failure of, contraindication, or adverse reaction to methotrexate
- ☐ Trial and failure of at least **ONE (1) other DMARD** therapy (**check each tried**):

<input type="checkbox"/> auranofin	<input type="checkbox"/> azathioprine
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> Other: _____

- ☐ Patient has tried and failed **TWO (2)** of the preferred drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

### ☐ Polyarticular Course Juvenile Idiopathic Arthritis

- ☐ Member is 2 years of age or older
- ☐ Member has a diagnosis of Polyarticular Course Juvenile Idiopathic Arthritis (pcJIA)
- ☐ Trial and failure of, contraindication, or adverse reaction to methotrexate
- ☐ Trial and failure of at least **ONE (1) other DMARD** therapy (**check each tried**):

<input type="checkbox"/> auranofin	<input type="checkbox"/> azathioprine
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> Other: _____

- ☐ Patient has tried and failed **TWO (2)** of the preferred drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

### ☐ Moderate-to-Severe Active Ulcerative Colitis

- ☐ Member is 18 years of age or older
- ☐ Member has a diagnosis of active ulcerative colitis
- ☐ Member has trial and failure of **BOTH** of the preferred drugs below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Infliximab
----------------------------------	-------------------------------------

(Continued on next page)

☐ **Ankylosing Spondylitis**

- ☐ Member is 18 years of age or older
- ☐ Member has a diagnosis of active ankylosing spondylitis
- ☐ Trial and failure of **TWO (2)** of the preferred drugs below:

☐ Humira®

☐ Enbrel®

☐ Infliximab

**Medication being provided by Specialty Pharmacy - PropriumRx**

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****