## SENTARA HEALTH PLANS

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (<u>including phone and fax #s</u>) on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

**<u>Drug Requested</u>**: **Rezdiffra**<sup>™</sup> (resmetirom)

steatohepatitis (NASH/MASH)

MEMBER & PRESCRIBER INF	<b>TORMATION:</b> Authorization may be delayed if incomplete.
Member Name:	
Member Sentara #:	Date of Birth:
Prescriber Name:	
Prescriber Signature:	Date:
Office Contact Name:	
Phone Number:	Fax Number:
NPI #:	
DRUG INFORMATION: Authoriz	zation may be delayed if incomplete.
Drug Name/Form/Strength:	
Dosing Schedule:	Length of Therapy:
Diagnosis:	ICD Code, if applicable:
Weight:	Date:
	mg once daily. $\geq$ 100 kg: 100 mg once daily. Coadministration with e to 80 mg daily for patients weighing $\geq$ 100 kg, or reduce dose to 60 strengths – 60, 80 & 100 mg)
	low all that apply. All criteria must be met for approval. To tion, including lab results, diagnostics, and/or chart notes, must be
<b>Initial Authorization: 12 months</b>	
☐ Member is 18 years of age or older	
☐ Medication is prescribed by or in co	onsultation with a hepatologist or gastroenterologist

(Continued on next page)

☐ Member has a diagnosis of nonalcoholic steatohepatitis or metabolic dysfunction—associated

ш	Provider must submit ONE of the following
	□ Biopsy results (completed within the last 6 months) documenting <b>BOTH</b> of the following:
	☐ Liver fibrosis stage F2 or F3
	□ Non-alcoholic fatty liver disease activity score (NAS) of ≥ 4 with a score of > 1 in all the following: steatosis, ballooning, and lobular inflammation
	Liver fibrosis stage F2 or F3 as determined by an elastography test, such as vibration-controlled transient elastography (i.e., FibroScan), magnetic resonance elastography (MRE), shear wave elastography; etc. (must submit current test results)
	In cases of indeterminate fibrosis stage (i.e., inconsistency between fibrosis stage and clinical presentation), a liver biopsy will be required to be submitted
	Member has three or more of the following metabolic risk factors that are managed according to standard of care (verified by medical chart notes, lab test results and/or pharmacy claims):
	□ Central obesity
	□ Hypertriglyceridemia
	□ Reduced high-density lipoprotein cholesterol
	□ Hypertension
	☐ Elevated fasting plasma glucose indicative of diabetes or pre-diabetes
	Current liver function (CMP) and CBC test results must be submitted
	Other causes of liver disease or hepatic steatosis have been ruled out (i.e., alcoholic steatohepatitis, acute fatty liver, autoimmune hepatitis, Hepatitis A, B or C, hemochromatosis, drug-induced liver disease)
	Member has adopted liver-protective lifestyle interventions such as optimizing weight loss, dietary changes, and exercise and is compliant with treatment for comorbidities (i.e. hyperlipidemia, hypertension, diabetes; etc.)
	Member does <u>NOT</u> have significant alcohol consumption (alcohol consumption of more than 20 g per day for women and more than 30 g per day for men)
	Member does <u>NOT</u> have evidence of cirrhosis, hepatic decompensation, or hepatocellular carcinoma (must submit documentation)
	Member has had an unsuccessful 6-month trial of Wegovy and documentation of insufficient clinical response (i.e. lack of MASH/NASH resolution, no improvement in fibrosis score; etc.) must be submitted
Reau	uthorization: 6 months. Check below all that apply. All criteria must be met for approval. To

☐ Member continues to meet <u>ALL</u> initial authorization criteria

provided or request may be denied.

support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be

(Continued on next page)

PA Rezdiffra (CORE) (Continued from previous page)

	Member has experienced <u>ONE</u> of the following as determined by an elastography test, such as vibration-controlled transient elastography (e.g., FibroScan), magnetic resonance elastography (MRE), shear wave elastography or biopsy:  □ MASH/NASH resolution <u>AND</u> no worsening of fibrosis □ No worsening of MASH/NASH <u>AND</u> improvement in fibrosis by ≥ 1 stage
Med	dication being provided by Specialty Pharmacy – Proprium Rx
	*Use of samples to initiate therapy does not meet step edit/preauthorization criteria.** evious therapies will be verified through pharmacy paid claims or submitted chart notes.*