

# SENTARA HEALTH PLANS

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization may be delayed.**

### Gastrointestinal (GI) Motility Drugs

<b>Drug Requested:</b> (select one drug below)		
<b>Non-Preferred</b>		
<input type="checkbox"/> <b>Ibsrela</b> <sup>®</sup> (tenapanor)	<input type="checkbox"/> <b>Motegrity</b> <sup>®</sup> (prucalopride)	<input type="checkbox"/> <b>Relistor</b> <sup>®</sup> (methylnaltrexone bromide)
<input type="checkbox"/> <b>Trulance</b> <sup>®</sup> (plecanatide)	<input type="checkbox"/> <b>Zelnorm</b> <sup>™</sup> (tegaserod)	

**MEMBER & PRESCRIBER INFORMATION:** Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

DEA OR NPI #: \_\_\_\_\_

**DRUG INFORMATION:** Authorization may be delayed if incomplete.

Drug Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code: \_\_\_\_\_

Weight: \_\_\_\_\_ Date: \_\_\_\_\_

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

(Continued on next page)

**❑ Approval of Motegrity® for diagnosis of Chronic Idiopathic Constipation (CIC)**

- ❑ Member has had trial and failure, contraindication, or intolerance to **ONE** of the following generic prerequisite therapies:

<input type="checkbox"/> lactulose	<input type="checkbox"/> polyethylene glycol (generic MiraLAX®)
------------------------------------	---

**AND**

- ❑ Member has had trial and failure, contraindication, or intolerance to lubiprostone (Amitiza®)

**AND**

- ❑ Member has had trial and failure, contraindication, or intolerance to Linzess®

**❑ Approval of Zelnorm™**

- ❑ Diagnosis of **Irritable Bowel Syndrome with Constipation (IBS-C)**

**AND**

- ❑ Member is < 65 years of age with no history of ischemic cardiovascular disease and has no more than one CVD risk factor. CVD risk factors are defined as active smoking, current hypertension/history of antihypertensive treatment, current hyperlipidemia/history of lipid lowering medication, history of diabetes mellitus, age >55 years, or obesity (BMI >30 kg/m<sup>2</sup>)

**AND**

- ❑ Provider attests that member does **NOT** have any of the following contraindications to therapy:
- History of myocardial infarction (MI), stroke, transient ischemic attack (TIA), or angina
  - History of ischemic colitis or other forms of intestinal ischemia
  - Severe renal impairment (eGFR < 15 mL/min/1.73 m<sup>2</sup>) or end-stage renal disease
  - Moderate and severe hepatic impairment (Child-Pugh B or C)
  - History of bowel obstruction, symptomatic gallbladder disease, suspected sphincter of Oddi dysfunction, or abdominal adhesions

**AND**

- ❑ Member has had trial and failure, contraindication, or intolerance to **ONE** of the following generic prerequisite therapies:

<input type="checkbox"/> lactulose	<input type="checkbox"/> polyethylene glycol (generic MiraLAX®)
------------------------------------	---

**AND**

- ❑ Member has had trial and failure, contraindication, or intolerance to lubiprostone (Amitiza®)

**AND**

(Continued on next page)

- Member has had trial and failure, contraindication, or intolerance to Linzess<sup>®</sup>

**Approval of Trulance<sup>®</sup> for diagnosis of Irritable Bowel Syndrome with Constipation (IBS-C) or Chronic Idiopathic Constipation (CIC)**

- Member has had trial and failure, contraindication, or intolerance to **ONE** of the following generic prerequisite therapies:

<input type="checkbox"/> lactulose	<input type="checkbox"/> polyethylene glycol (generic MiraLAX <sup>®</sup> )
------------------------------------	--

**AND**

- Member has had trial and failure, contraindication, or intolerance to lubiprostone (Amitiza<sup>®</sup>)

**AND**

- Member has had trial and failure, contraindication, or intolerance to Linzess<sup>®</sup>

**Approval of Ibsrela<sup>®</sup> for diagnosis of Irritable Bowel Syndrome with Constipation (IBS-C)**

- Member has had trial and failure, contraindication, or intolerance to **ONE** of the following generic prerequisite therapies:

<input type="checkbox"/> lactulose	<input type="checkbox"/> polyethylene glycol (generic MiraLAX <sup>®</sup> )
------------------------------------	--

**AND**

- Member has had trial and failure, contraindication, or intolerance to lubiprostone (Amitiza<sup>®</sup>)

**AND**

- Member has had trial and failure, contraindication, or intolerance to Linzess<sup>®</sup>

**AND**

- Member has had trial and failure, contraindication, or intolerance to Trulance<sup>®</sup> (**requires prior authorization**)

**Approval of Relistor<sup>®</sup>**

**Recommended Dosing:**

Weight of Adult Patient	Subcutaneous Dose	Injection Volume
Less than 38kg	0.15 mg/kg	See below
38kg to less than 62 kg	8mg	0.4 mL
62kg to 114kg	12mg	0.6 mL
More than 114kg	0.15 mg/kg	See below

\*Calculate injection volume by multiplying member weight in kilograms by 0.0075, then round up the volume to the nearest 0.1 mL\*

(Continued on next page)

- ❑ Select **ONE** of the following:
  - ❑ Member has a diagnosis of opioid-induced constipation (OIC) with advanced illness or pain caused by active cancer who require opioid dosage escalation for palliative care
  - ❑ Member has a diagnosis of opioid-induced constipation (OIC) with chronic non-cancer pain, including members with chronic pain related to prior cancer or its treatment who do not require frequent (e.g., weekly) opioid dosage escalation

**AND**

- ❑ Member has been on an opioid within the last 60 days of prior authorization request but not less than 4 weeks. Provider please note: Members receiving opioids for less than 4 weeks may be less responsive to Relistor<sup>®</sup>

**AND**

- ❑ Member has had trial and failure, contraindication, or intolerance to **ONE** of the following generic prerequisite therapies:

<input type="checkbox"/> lactulose	<input type="checkbox"/> polyethylene glycol (generic MiraLAX <sup>®</sup> )
------------------------------------	--

**AND**

- ❑ Member has had trial and failure, contraindication, or intolerance to lubiprostone (Amitiza<sup>®</sup>)

**AND**

- ❑ Member has had trial and failure, contraindication, or intolerance to both Movantik<sup>®</sup>  
**AND** Symproic<sup>®</sup>

*Not all drugs may be covered under every Plan*

*If a drug is non-formulary on a Plan, documentation of medical necessity will be required.*

***\*\*Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.\*\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****