

# SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

**Drug Requested:** Actemra® (tocilizumab) and Tyenne® (tocilizumab-aazg) SQ  
(Pharmacy) (Non-Preferred)

|   |   |
|---|---|
| <input type="checkbox"/> Actemra® (tocilizumab) | <input type="checkbox"/> Tyenne® (tocilizumab-aazg) |
|---|---|

### MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

NPI #: \_\_\_\_\_

### DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight (if applicable): \_\_\_\_\_ Date weight obtained: \_\_\_\_\_

| DIAGNOSIS   | Recommended Dose  |
|---|---|
| <input type="checkbox"/> Rheumatoid Arthritis (RA)                          | SUBCUTANEOUS <ul style="list-style-type: none"><li>Weight &lt;100kg: Two syringes per 28 days. Max dose is 4 syringes per 28 days</li><li>Weight &gt;100kg: Four syringes per 28 days</li></ul> |
| <input type="checkbox"/> Polyarticular Juvenile Idiopathic Arthritis (PJIA) | SUBCUTANEOUS <ul style="list-style-type: none"><li>Weight &lt;30kg: 162mg/dose once every 3 weeks</li><li>Weight ≥30kg: 162mg/dose once every 2 weeks</li></ul>                                 |
| <input type="checkbox"/> Systemic Juvenile Idiopathic Arthritis (SJIA)      | SUBCUTANEOUS <ul style="list-style-type: none"><li>Weight &lt;30kg: 162mg/dose once every 2 weeks</li><li>Weight ≥30kg: 162mg/dose every week</li></ul>   |

(Continued on next page)

| DIAGNOSIS   | Recommended Dose   |
|---|--|
| <input type="checkbox"/> <b>Giant Cell Arteritis (GCA)</b>  | <b>SUBCUTANEOUS</b> <ul style="list-style-type: none"> <li>162mg once a week, may consider 162mg once every 2 weeks</li> </ul> |
| <input type="checkbox"/> <b>Systemic Sclerosis- Associated Interstitial Lung Disease (SSc-ILD)- Actemra® only</b> | <b>SUBCUTANEOUS</b> <ul style="list-style-type: none"> <li>162mg once a week</li> </ul>  |

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

☐ **Diagnosis: Rheumatoid Arthritis (RA)**

- ☐ Prescriber is a Rheumatologist; **AND**
- ☐ Member has moderate to severe rheumatoid arthritis; **AND**
- ☐ Tried and failed methotrexate; **OR**
- ☐ Requested medication will be used in conjunction with methotrexate; **OR**
- ☐ Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or other contraindication); **AND**
- ☐ Tried and failed at least **ONE (1) DMARD** other than methotrexate and (check each tried)

|  |   |   |
|--|---|---|
| <input type="checkbox"/> sulfasalazine   | <input type="checkbox"/> azathioprine       | <input type="checkbox"/> leflunomide      |
| <input type="checkbox"/> auranofin       | <input type="checkbox"/> hydroxychloroquine | <input type="checkbox"/> gold salts       |
| <input type="checkbox"/> d-penicillamine | <input type="checkbox"/> cyclosporine       | <input type="checkbox"/> cyclophosphamide |
| <input type="checkbox"/> tacrolimus      | <input type="checkbox"/> Other: _____       |   |

**AND**

- ☐ Trial and failure of **TWO (2)** of the **PREFERRED** biologics below:

|                                  |                                  |                                     |
|----------------------------------|----------------------------------|-------------------------------------|
| <input type="checkbox"/> Humira® | <input type="checkbox"/> Enbrel® | <input type="checkbox"/> Infliximab |
|----------------------------------|----------------------------------|-------------------------------------|

☐ **Diagnosis: Polyarticular Juvenile Idiopathic Arthritis (PJIA)**

- ☐ Prescriber is a Rheumatologist; **AND**
- ☐ Member must be 2 years of age and older with active polyarticular juvenile idiopathic arthritis; **AND**

(Continued on next page)

- ☐ Tried and failed methotrexate; **OR**
- ☐ Requested medication will be used in conjunction with methotrexate; **OR**
- ☐ Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or other contraindication); **AND**
- ☐ Trial and failure of **TWO (2)** of the **PREFERRED** biologics below:

|                                  |                                  |
|----------------------------------|----------------------------------|
| <input type="checkbox"/> Humira® | <input type="checkbox"/> Enbrel® |
|----------------------------------|----------------------------------|

**AND**

- ☐ Trial and failure of at least **ONE (1) DMARD** therapy **and** (**check each tried**)

|  |   |   |
|--|---|---|
| <input type="checkbox"/> sulfasalazine   | <input type="checkbox"/> azathioprine       | <input type="checkbox"/> leflunomide      |
| <input type="checkbox"/> auranofin       | <input type="checkbox"/> hydroxychloroquine | <input type="checkbox"/> gold salts       |
| <input type="checkbox"/> d-penicillamine | <input type="checkbox"/> cyclosporine       | <input type="checkbox"/> cyclophosphamide |
| <input type="checkbox"/> tacrolimus      | <input type="checkbox"/> Other: _____       |   |

☐ **Diagnosis: Systemic Juvenile Idiopathic Arthritis (SJIA)**

- ☐ Prescriber is a Rheumatologist; **AND**
- ☐ Member must be 2 years of age and older with active systemic juvenile idiopathic arthritis; **AND**
- ☐ Tried and failed methotrexate; **OR**
- ☐ Requested medication will be used in conjunction with methotrexate; **OR**
- ☐ Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or other contraindication); **AND**
- ☐ Trial and failure of at least **ONE (1) DMARD** therapy **and** (**check each tried**)

|  |   |   |
|--|---|---|
| <input type="checkbox"/> sulfasalazine   | <input type="checkbox"/> azathioprine       | <input type="checkbox"/> leflunomide      |
| <input type="checkbox"/> auranofin       | <input type="checkbox"/> hydroxychloroquine | <input type="checkbox"/> gold salts       |
| <input type="checkbox"/> d-penicillamine | <input type="checkbox"/> cyclosporine       | <input type="checkbox"/> cyclophosphamide |
| <input type="checkbox"/> tacrolimus      | <input type="checkbox"/> Other: _____       |   |

(Continued on next page)

☐ **Diagnosis: Giant Cell Arteritis (GCA)**

- ☐ **For Actemra® & Tyenne® requests only:** Member must be 18 years of age or older with giant cell arteritis (GCA) diagnosis

☐ **Diagnosis: Systemic Sclerosis- Associated Interstitial Lung Disease (SSc-ILD)**

- ☐ **For Actemra® requests only:** Member must be 18 years of age or older with systemic sclerosis-associated interstitial lung disease (SSc-ILD)

**Medication being provided by Specialty Pharmacy - PropriumRx**

***\*Use of samples to initiate therapy does not meet step-edit/preauthorization criteria.\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****