

# SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

**Drug Requested:** Actemra® (tocilizumab) and Tyenne® (tocilizumab-aazg) SQ (Pharmacy)  
**(Non- Preferred)**

<input type="checkbox"/> Actemra® (tocilizumab)	<input type="checkbox"/> Tyenne® (tocilizumab-aazg)
---	---

<b>MEMBER &amp; PRESCRIBER INFORMATION:</b> Authorization may be delayed if incomplete.
---

Member Name: \_\_\_\_\_

Member Sentara #: \_\_\_\_\_ Date of Birth: \_\_\_\_\_

Prescriber Name: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Office Contact Name: \_\_\_\_\_

Phone Number: \_\_\_\_\_ Fax Number: \_\_\_\_\_

NPI #: \_\_\_\_\_

<b>DRUG INFORMATION:</b> Authorization may be delayed if incomplete.
--

Drug Name/Form/Strength: \_\_\_\_\_

Dosing Schedule: \_\_\_\_\_ Length of Therapy: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD Code, if applicable: \_\_\_\_\_

Weight (if applicable): \_\_\_\_\_ Date weight obtained: \_\_\_\_\_

DIAGNOSIS	Recommended Dose
<input type="checkbox"/> Rheumatoid Arthritis (RA)	<b>SUBCUTANEOUS</b> <ul style="list-style-type: none"> <li>Weight &lt;100kg: Two syringes per 28 days. Max dose is 4 syringes per 28 days</li> <li>Weight &gt;100kg: Four syringes per 28 days</li> </ul>
<input type="checkbox"/> Polyarticular Juvenile Idiopathic Arthritis (PJIA)	<b>SUBCUTANEOUS</b> <ul style="list-style-type: none"> <li>Weight &lt;30kg: 162mg/dose once every 3 weeks</li> <li>Weight ≥30kg: 162mg/dose once every 2 weeks</li> </ul>
<input type="checkbox"/> Systemic Juvenile Idiopathic Arthritis (SJIA)	<b>SUBCUTANEOUS</b> <ul style="list-style-type: none"> <li>Weight &lt;30kg: 162mg/dose once every 3 weeks</li> <li>Weight ≥30kg: 162mg/dose every 2 weeks</li> </ul>

(Continued on next page)

DIAGNOSIS	Recommended Dose
<input type="checkbox"/> <b>Giant Cell Arteritis (GCA)</b>	<b>SUBCUTANEOUS</b> <ul style="list-style-type: none"> <li>162mg once a week, may consider 162mg once every 2 weeks</li> </ul>
<input type="checkbox"/> <b>Systemic Sclerosis- Associated Interstitial Lung Disease (SSc-ILD)- Actemra<sup>®</sup> only</b>	<b>SUBCUTANEOUS</b> <ul style="list-style-type: none"> <li>162mg once a week</li> </ul>

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

**Diagnosis: Rheumatoid Arthritis (RA)**

- Prescriber is a Rheumatologist; **AND**
- Member has moderate to severe rheumatoid arthritis; **AND**
- Tried and failed methotrexate; **OR**
- Requested medication will be used in conjunction with methotrexate; **OR**
- Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or other contraindication); **AND**
- Tried and failed at least **ONE (1) DMARD** other than methotrexate and (check each tried)

<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> auranofin	<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> gold salts
<input type="checkbox"/> d-penicillamine	<input type="checkbox"/> cyclosporine	<input type="checkbox"/> cyclophosphamide
<input type="checkbox"/> tacrolimus	<input type="checkbox"/> Other: _____	

**AND**

- Trial and failure of **TWO (2)** of the **PREFERRED** biologics below:

<input type="checkbox"/> Humira <sup>®</sup>	<input type="checkbox"/> Enbrel <sup>®</sup>	<input type="checkbox"/> Infliximab
--	--	-------------------------------------

**Diagnosis: Polyarticular Juvenile Idiopathic Arthritis (PJIA)**

- Prescriber is a Rheumatologist; **AND**
- Member must be 2 years of age and older with active polyarticular juvenile idiopathic arthritis; **AND**

(Continued on next page)

- Tried and failed methotrexate; **OR**
- Requested medication will be used in conjunction with methotrexate; **OR**
- Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or other contraindication); **AND**
- Trial and failure of **TWO (2)** of the **PREFERRED** biologics below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®
----------------------------------	----------------------------------

**AND**

- Trial and failure of at least **ONE (1) DMARD** therapy **and (check each tried)**

<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> auranofin	<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> gold salts
<input type="checkbox"/> d-penicillamine	<input type="checkbox"/> cyclosporine	<input type="checkbox"/> cyclophosphamide
<input type="checkbox"/> tacrolimus	<input type="checkbox"/> Other: _____	

**Diagnosis: Systemic Juvenile Idiopathic Arthritis (SJIA)**

- Prescriber is a Rheumatologist; **AND**
- Member must be 2 years of age and older with active systemic juvenile idiopathic arthritis; **AND**
- Tried and failed methotrexate; **OR**
- Requested medication will be used in conjunction with methotrexate; **OR**
- Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or other contraindication); **AND**
- Trial and failure of at least **ONE (1) DMARD** therapy **and (check each tried)**

<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> auranofin	<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> gold salts
<input type="checkbox"/> d-penicillamine	<input type="checkbox"/> cyclosporine	<input type="checkbox"/> cyclophosphamide
<input type="checkbox"/> tacrolimus	<input type="checkbox"/> Other: _____	

(Continued on next page)

**❑ Diagnosis: Giant Cell Arteritis (GCA)**

- ❑ **For Actemra® & Tyenne® requests only:** Member must be 18 years of age or older with giant cell arteritis (GCA) diagnosis

**❑ Diagnosis: Systemic Sclerosis- Associated Interstitial Lung Disease (SSc-ILD)**

- ❑ **For Actemra® requests only:** Member must be 18 years of age or older with systemic sclerosis-associated interstitial lung disease (SSc-ILD)

**Medication being provided by Specialty Pharmacy - PropriumRx**

***\*Use of samples to initiate therapy does not meet step-edit/preauthorization criteria.\****

***\*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\****