SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information <u>(including phone and fax #s)</u> on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

<u>Drug Requested</u>: Actemra (tocilizumab) and Tyenne® (tocilizumab-aazg) SQ (Pharmacy) (Non-Preferred)

□ Actemra® (tocilizumab)	□ Tyenne ® (tocilizumab-aazg)			
MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.				
Member Name:				
Member Sentara #:				
Prescriber Name:				
Prescriber Signature:	Date:			
Office Contact Name:				
one Number: Fax Number:				
NPI #:				
DRUG INFORMATION: Authorization may be delayed if incomplete.				
Drug Name/Form/Strength:				
	Length of Therapy:			
Diagnosis:	ICD Code, if applicable:			
Weight (if applicable):	Date weight obtained:			
DIAGNOSIS	Recommended Dose			
□ Rheumatoid Arthritis (RA)	 SUBCUTANEOUS Weight <100kg: Two syringes per 28 days. Max dose is 4 syringes per 28 days Weight >100kg: Four syringes per 28 days 			
□ Polyarticular Juvenile Idiopathic Arthritis (PJIA)	 SUBCUTANEOUS Weight <30kg: 162mg/dose once every 3 weeks Weight ≥30kg: 162mg/dose once every 2 weeks 			
□ Systemic Juvenile Idiopathic Arthritis (SJIA)	 SUBCUTANEOUS Weight <30kg: 162mg/dose once every 3 weeks Weight ≥30kg: 162mg/dose every 2 weeks 			

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

□ D	□ Diagnosis: Rheumatoid Arthritis (RA)					
	Prescriber is a Rheumatologist; AND					
	Member has moderate to severe rheumatoid arthritis; AND					
	Tried and failed methotrexate; OR					
	Requested medication will be used in conjunction with methotrexate; OR					
	Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or					
_	other contraindication); AND	Darring d. d.	.1			
	Tried and failed at least ONE (1) DMARD other than	n methotrexate ai	nd (check each tried)		
	□ sulfasalazine	□ azathioprine		□ leflunomide		
	□ auranofin	□ hydroxychloroqu	iine	□ gold salts		
	□ d-penicillamine	□ cyclosporine		□ cyclophosphamide		
	□ tacrolimus	Other:				
	AND					
	☐ Trial and failure of TWO (2) of the PREFERRED biologics below:					
	☐ Humira [®]	□ Enbrel [®]		□ Infliximab		
□ Diagnosis: Polyarticular Juvenile Idiopathic Arthritis (PJIA)						
	Prescriber is a Rheumatologist;	AND				
	Member must be 2 years of age and older with active polyarticular juvenile idiopathic arthritis; AND					
	Tried and failed methotrexate; OR					
	Requested medication will be used in conjunction with methotrexate; OR					
	Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or					
	other contraindication); AND Trial and failure of TWO (2) of the PREFERRED biologics below:					
	☐ Humira®		□ Enbrel®			
	AND					

AND

(Continued on next page)

	Trial and failure of at least ONE (1) DMARD therapy and (check each tried)						
	□ sulfasalazine	□ azathioprine	□ leflunomide				
	□ auranofin	□ hydroxychloroquine	□ gold salts				
	□ d-penicillamine	□ cyclosporine	□ cyclophosphamide				
	□ tacrolimus	Other:					
ı Di	Diagnosis: Systemic Juvenile Idiopathic Arthritis (SJIA)						
	Prescriber is a Rheumatologist; AND						
	Member must be 2 years of age and older with active systemic juvenile idiopathic arthritis; AND						
	Tried and failed methotrexate; OR						
	Requested medication will be used in conjunction with methotrexate; OR						
	Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or other contraindication); AND						
	□ sulfasalazine	□ azathioprine	□ leflunomide				
	□ auranofin	□ hydroxychloroquine	□ gold salts				
	□ d-penicillamine	□ cyclosporine	□ cyclophosphamide				
	□ tacrolimus	□ Other:					

Use of samples to initiate therapy does not meet step-edit/preauthorization criteria.

Medication being provided by Specialty Pharmacy - PropriumRx

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *