# SENTARA HEALTH PLANS

## **PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\***

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process may be delayed.

### Drug Requested: Recorlev<sup>®</sup> (levoketoconazole)

#### MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name:	
	Date of Birth:
Prescriber Name:	
Prescriber Signature:	
Office Contact Name:	
Phone Number:	
DEA OR NPI #:	
DRUG INFORMATION: Authorization may be delayed if incomplete.   Drug Form/Strength:	
	Length of Therapy:
Diagnosis:	ICD Code, if applicable:
Quantity Limits: 240 tablets per 30 days	
	apply. All criteria must be met for approval. To support sults, diagnostics, and/or chart notes, must be provided
Initial Authorization: 6 months	
□ Member is 18 years of age or older	

- $\hfill\square$  Prescribed by or in consultation with an endocrinologist or neurosurgeon
- □ Member has a diagnosis of persistent or recurrent Cushing's disease
- □ Member must meet at least <u>ONE</u> of the following (chart notes must be submitted to document diagnosis and surgical history or contraindication to surgery):
  - □ Member has undergone pituitary surgery and must be at least 6 weeks post-surgery
  - □ Member has undergone irradiation and must be at least 4 years post-pituitary irradiation
  - □ Member is contraindicated to surgery AND irradiation

(Continued on next page)

- □ Member must have failed <u>90 days</u> of therapy with **ketoconazole** tablets (verified by chart notes or **pharmacy paid claims**). Please provide documentation to support failure of oral ketoconazole along with rationale for use of requested medication
- □ Member must have current mean urine free cortisol levels (mUFC) >1.5 times the upper limit of normal (ULN)
- □ Member will <u>NOT</u> use concurrent Cushing's disease treatment with Recorlev (e.g., Isturisa, ketoconazole, metyrapone, mifepristone, mitotane)
- □ Member has been assessed for QTc prolongation/Torsade de Pointes
- Member has been assessed for hypokalemia, hypomagnesemia and treatment, as needed, prior to initiating therapy
- □ Member is <u>NOT</u> taking any CYP3A4 and/or P-gp substrate medications (e.g., lovastatin, simvastatin, tacrolimus)
- □ Provider attests member has been counseled to avoid excessive alcohol consumption
- □ Member does <u>NOT</u> have a history of any of the following:
  - Cirrhosis
  - Liver disease
  - Cholelithiasis
  - Baseline QTc prolongation
  - Ventricular tachycardia/fibrillation

**<u>Reauthorization</u>: 12 months.** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- □ Member's current 24-hour urinary free cortisol level is below the upper limit of normal (labs must be submitted)
- □ Improvements in quality of life have been maintained while on Recorlev therapy
- Member will continue to be monitored for QTc prolongation, electrolyte imbalances and hepatic impairment

#### **Medication being provided by Specialty Pharmacy - PropriumRx**

\*\* Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. \*\*

\*<u>Previous therapies will be verified through pharmacy paid claims or submitted chart notes.</u>\*