

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

Immunomodulators Atopic Dermatitis

Drug Requested: (check box below that applies)

PREFERRED		
<input type="checkbox"/> Adbry™ (tralokinumab)	<input type="checkbox"/> Dupixent® (dupilumab) (QL, AG) (Refer to Dupixent PA form)	<input type="checkbox"/> Eucrisa™ (crisaborole)
<input type="checkbox"/> pimecrolimus	<input type="checkbox"/> tacrolimus (generic Protopic®)	
Non-Preferred		
<input type="checkbox"/> Cibinqo™ (abrocitinib) (Refer to Cibinqo PA form)	<input type="checkbox"/> Ebglyss™ (lebrikizumab-lbkz) (Refer to Berlys PA form)	<input type="checkbox"/> Nemluvio® (nemolizumab-ilto) (Refer to Nemluvio PA form)
<input type="checkbox"/> Opzelura™ (ruxolitinib) (QL, AG) (Refer to Opzelura PA form)	<input type="checkbox"/> Protopic® (tacrolimus)	<input type="checkbox"/> Vtama® (tapinarof)
<input type="checkbox"/> Zoryve® cream 0.15%	<input type="checkbox"/> Zoryve® foam 0.3%	

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

(Continued on next page)

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ **Length of Therapy:** _____

Diagnosis: _____ **ICD Code, if applicable:** _____

Weight (if applicable): _____ **Date weight obtained:** _____

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Length of Authorization: 1 year

- ☐ Member must have an FDA-approved diagnosis of **Atopic Dermatitis**
 - ☐ **Adbry™**: moderate to severe for ages ≥ 12 years
 - ☐ **Eucria™**: mild to moderate for ages equal to or > 3 months
 - ☐ **pimecrolimus**: mild to moderate for ages ≥ 2 years
 - ☐ **Protopic®** 0.03%: moderate to severe for ages ≥ 2 years
 - ☐ **Protopic®** 0.1%: moderate to severe for ages ≥ 16 years
 - ☐ **Zoryve® cream** 0.15%: mild to moderate for ages ≥ 6 years
 - ☐ **Vtama® cream** 0.1%: for ages ≥ 2 years
 - ☐ For **pimecrolimus** or **tacrolimus**:
 - ☐ Prior documented trial and failure of 8 weeks (or contraindication) to one (1) medium to high potency (e.g., mometasone, triamcinolone) topical corticosteroid
 - ☐ For **Eucria™** or **Adbry®**:
 - ☐ Prior documented trial and failure of 30 days (or contraindication) to one (1) medium to high potency (e.g., mometasone, triamcinolone) topical corticosteroid
- OR**
- ☐ Prior documented trial and failure of 30 days (or contraindication) to one (1) topical calcineurin inhibitor (tacrolimus or pimecrolimus)
 - ☐ For **Protopic®**:
 - ☐ Failure to topical medium to high potency corticosteroids (e.g., mometasone, triamcinolone)
 - ☐ Failure to **pimecrolimus** and **tacrolimus** (generic)

(Continued on next page)

- ☐ For Zorvyte[®] cream 0.15% or Vtama[®]:
 - ☐ Prior documented trial and failure of 8 weeks of each:
 - ☐ One (1) topical corticosteroid of medium to high potency (e.g., mometasone, triamcinolone)
 - ☐ One (1) topical calcineurin inhibitor (tacrolimus or pimecrolimus)
 - ☐ Trial and failure of Dupixent[®]
 - ☐ **OR** for Vtama[®] for plaque psoriasis
 - ☐ Member is 18 years of age or older and has a diagnosis of plaque psoriasis
 - ☐ Member has a history of failure, contraindication, or intolerance to calcipotriene cr/oint/soln
 - ☐ For Zoryve[®] foam 0.3%:
 - ☐ Member is 9 years of age or older and has a diagnosis of seborrheic dermatitis
 - ☐ Prior documented trial and failure of 30 days (or contraindication) to one (1) topical corticosteroid (i.e., clobetasol, fluocinonide or mometasone cream/ointment/solution) in the past 180 days
 - ☐ Prior documented trial and failure of 30 days (or contraindication) to one (1) topical antifungal (ciclopirox shampoo/gel, ketoconazole cream/shampoo, selenium sulfide 2.25% shampoo) in the past 180 days
- OR**
- ☐ Member is 12 years of age or older and has a diagnosis of plaque psoriasis
 - ☐ Member has a history of failure, contraindication, or intolerance to calcipotriene cr/oint/soln

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****