SENTARA HEALTH PLANS

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-844-668-1550</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization can be delayed</u>.

For Medicare Members: Medicare Coverage for outpatient (Part B) drugs is outlined in the Medicare Benefit Policy Manual (Pub. 100-2), Chapter 15, §50 Drugs and Biologicals. In addition, National Coverage Determination (NCD) and Local Coverage Determinations (LCDs) may exist and compliance with these policies is required where applicable. They can be found at: https://www.cms.gov/medicare-coverage-database/overview-and-quick-search.aspx. Additional indications may be covered at the discretion of the health plan.

The Sentara Health Plans Oncology Program is administered by OncoHealth

- ❖ For any oncology indications, the most efficient way to submit a prior authorization request is through the OncoHealth OneUM Provider Portal at https://oneum.oncohealth.us. Fax to 1-800-264-6128. OncoHealth can also be contacted by Phone: 1-888-916-2616.
- ❖ Commercial customers <u>NOT</u> enrolled in the OncoHealth program, please fax requests to Sentara Health plans at fax number 1-844-668-1550.

<u>Drug Requested</u>: Aucatzyl[®] (obecabtagene autoleucel) (Q2058) (Medical)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.				
Member Name:				
Member Sentara #:	Date of Birth:			
Prescriber Name:				
Prescriber Signature:	Date:			
Office Contact Name:				
Phone Number:				
NPI #:				
DRUG INFORMATION: Authoriza				
Drug Name/Form/Strength:				
Dosing Schedule:	Length of Therapy:			
Diagnosis:	ICD Code, if applicable:			
Weight (if applicable):	Date weight obtained:			
	the timeframe does not jeopardize the life or health of the member num function and would not subject the member to severe pain.			

(Continued on next page)

Dosing Limits

A. Quantity Limit (max daily dose) [NDC Unit]:

• The total recommended dose of Aucatzyl is 410×10^6 CD19 chimeric antigen receptor (CAR)-positive viable T cells

B. Max Units (per dose and over time) [HCPCS Unit]:

• One treatment (dose) per lifetime.

☐ Member is 18 years of age or older

- Aucatzyl contains a total recommended dose of 410 x 10⁶ CD19 CAR-positive viable T cells supplied in 3 to 5 infusion bag.
- The treatment regimen consists of a split dose infusion to be administered on Day 1 and Day 10 (\pm 2 days). Dose to be administered is determined by the patient bone marrow blast assessment.

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Coverage will be provided for one treatment course and may <u>NOT</u> be renewed.

•	
Provider is an oncologist and the administrating healthcare facility providers have received training on the management of cytokine release syndrome (CRS) and neurological toxicities	
Member has NOT received prior CAR-T therapy	
Member must meet ONE of the following:	
☐ Member has <u>NOT</u> received prior anti-CD19 therapy (e.g., blinatumomab)	
☐ Member previously received anti-CD19 therapy and re-biopsy indicates CD-19 positive disease	
Medication will be used as single agent therapy (not applicable to lymphodepleting or additional chemotherapy while awaiting manufacture)	
Member has been screened for hepatitis B virus (HBV), hepatitis C virus (HCV), and human immunodeficiency virus (HIV) in accordance with clinical guidelines prior to collection of cells (leukapheresis)	
Member does NOT have a clinically significant active systemic infection or inflammatory disorder	
Prophylaxis for infection has been followed according to local guidelines	
Member has <u>NOT</u> received live vaccines within 6 weeks prior to the start of lymphodepleting chemotherapy, during obecabtagene autoleucel treatment, and will <u>NOT</u> receive live vaccines until immune recovery following treatment	
Member has a diagnosis of relapsed or refractory B-Cell Precursor Acute Lymphoblastic Leukemia (ALL)	

(Continued on next page)

		one Marrow Blast Percentage has been assessed and the laboratory documentation has been provided to eet ONE of the following dose recommendations:	
		Bone Marrow Blast > 20%: Day 1 infusion with a dose of 10×10^6 ; Day 10 (±2 days) 100×10^6 Dose and 300×10^6 Dose	
		Bone Marrow Blast \leq 20%: Day 1 infusion with a dose of 100×10^6 ; Day 10 (\pm 2 days) 10×10^6 Dose and 300×10^6 Dose	
	Me	ember's condition meets ONE of the following:	
		Philadelphia Chromosome (Ph)-Positive disease with prior therapy that includes a tyrosine kinase inhibitor (e.g., bosutinib, dasatinib, imatinib)	
		Philadelphia Chromosome (Ph)-Negative disease	
Medication being provided by: Please check applicable box below.			
_	Loca	ation/site of drug administration:	

□ Specialty Pharmacy

<u>OR</u>

NPI or DEA # of administering location:

For urgent reviews: Practitioner should call Sentara Health Plans Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health Plan's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

Use of samples to initiate therapy does not meet step edit/preauthorization criteria.

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.