

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

The Sentara Health Plans Oncology Program is administered by OncoHealth

- ❖ **For any oncology indications,** the most efficient way to submit a prior authorization request is through the OncoHealth OneUM Provider Portal at <https://oneum.oncohealth.us>. Fax to 1-800-264-6128.

OncoHealth can also be contacted at Phone: 1-888-916-2616

Drug Requested: Jylamvo[®] (methotrexate) oral solution (Non-Preferred)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Name/Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

Recommended Dosage:

Indication:	Dosage:
Treatment of adult and pediatric patients with acute lymphoblastic leukemia (ALL) as a component of a combination chemotherapy maintenance regimen	<ul style="list-style-type: none">ALL: 20 mg/m² once weekly
Treatment of pediatric patients with polyarticular juvenile idiopathic arthritis (pJIA)	<ul style="list-style-type: none">Starting dose of 10 mg/m² one time weekly and adjust to optimal response

(Continued on next page)

Indication:	Dosage:
Treatment of adults with mycosis fungoides	<ul style="list-style-type: none">• 25-75mg once weekly (monotherapy), or 10mg/m² twice weekly as combination therapy
Treatment adults with relapsed or refractory non-Hodgkin lymphoma as part of a metronomic combination regimen	<ul style="list-style-type: none">• 2.5mg two to four times weekly
Treatment of adults with rheumatoid arthritis	<ul style="list-style-type: none">• Start at 7.5mg once weekly and adjust to optimal response
Treatment of adults with severe psoriasis	<ul style="list-style-type: none">• 10-25mg once weekly

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ☐ Member meets **ONE** of the following:
 - ☐ Dosing will not allow the use of preferred methotrexate tablets
 - ☐ Member is unable to swallow methotrexate tablets

Medication being provided by Specialty Pharmacy - PropriumRx

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****