SENTARA COMMUNITY PLAN (MEDICAID)

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-844-305-2331</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization can be delayed</u>.

Drug Requested: Check applicable box below. If not checked, authorization could be delayed.

<u>Drug Requested:</u> Immune Globulin Intravenous (IVIG) (Medical) (Miscellaneous Sjögren's Syndrome Neuropathies)

Prescriber Name:

Office Contact Name:

DEA OR NPI #: _____

| <i>D</i> 1 | Drug Requested. Check applicable box below. If not enecked, authorization could be delayed. | | | | |
|--|---|--|--|--|--|
| | Bivigam [®] [Immune Globulin Intravenous (Human), 10% Liquid] (J1556) | | Gammaked [™] [Immune Globulin Injection (Human), 10% Caprylate/Chromatography Purified] (J1561) | | |
| | Carimune® NF [Nanofiltered, Immune Globulin Intravenous (Human)] (J1566) | | Gamunex®-C [Immune Globulin Injection (Human), 10% Caprylate/Chromatography Purified] (J1561) | | |
| | Flebogamma® DIF [Human Normal Immunoglobulin (IVIg)] (J1572) | | Octagam [®] [Immune Globulin Intravenous (Human) liquid preparation] (J1568) | | |
| | Gammagard® Liquid [Immune Globulin Infusion (Human), 10% Solution, for intravenous and subcutaneous administration] (J1569) | | Panzyga® [Immune Globulin Intravenous (Human) – ifas 10% Liquid Preparation] (J1576) | | |
| | Gammagard® S/D [Immune Globulin Intravenous (Human) Solvent/Detergent Treated (Freeze-Dried Concentrate)] (J1556) | | Privigen® [Immune Globulin Intravenous (Human), 10% Liquid] (J1459) | | |
| | | | | | |
| MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete. | | | | | |
| Member Name: | | | | | |
| Mei | Member Sentara #: Date of Birth: | | | | |

(Continued on next page)

Prescriber Signature: Date:

Phone Number: _____ Fax Number: _____

| DRUG | INFORMATION: Authorizati | on may be delayed if incomplete. |
|-----------|---|---|
| Drug For | m/Strength: | |
| Dosing So | chedule: | Length of Therapy: |
| Diagnosis | s: | ICD Code, if applicable: |
| Weight: | | Date: |
| Height: _ | inches | |
| | | ne timeframe does not jeopardize the life or health of the member of function and would not subject the member to severe pain. |
| | RTANT* - If recommended a will be granted. | adjusted body weight is not accepted, a PARTIAL |
| _ | nould be calculated using adjusted be ideal body weight (IBW). | ody weight if the member's actual body weight is 20% higher than |
| • IB | Adjusted body weight = $IBW + 0.5$ (a) W (kg) for males = $50 + [2.3$ (height W (kg) for females = $45.5 + [2.3 \times 10^{-3}]$ (kg) | t in inches -60)] |
| and subse | quently stop IVIG therapy if improvitions for Primary Immunodeficiency | n the dose for renewal requests when improvement has occurred rement is sustained with a dose reduction (this does NOT apply to (PID) as long as immunoglobulin levels are maintained in the |
| approval | | IS: Check below all that apply. All criteria must be met for locumentation, including lab results, diagnostics, and/or chart enied. |
| □ Diag | gnosis: Small Fiber Sensory N | Neuropathy associated with Sjögren's Syndrome |
| Initial A | <u>Authorization</u> : 6 months (Do | se: 2 g/kg/month) |
| mo | onths due to toxicity OR failure to strongression. Check ALL that apply Tricyclic antidepressant | WO of the following first-line therapies for a minimum of 3 cabilize disease. Submit supporting document on toxicities and (include drug name, dose, duration): |
| | SNRI | |
| | Anticonvulsant | |
| | Topical lidocaine | |

(Continued on next page)

| | Diagnosis based on American-European Consensus Group Criteria (presence of 4 out of 6 items if histopathology or autoantibodies positive <u>OR</u> THREE (3) of the following are positive: ocular signs, histopathology, salivary gland involvement, autoantibodies): □ Ocular symptoms (<u>at least 1</u>): | |
|------------|--|---|
| | | □ Daily, persistent, troublesome dry eyes for more than 3 months |
| | | ☐ Recurrent sensation of sand or gravel in eyes |
| | | ☐ Use of tear substitutes more than 3 times a day |
| | | |
| | | ☐ Daily feeling of dry mouth for more than 3 months |
| | | ☐ Recurrent or persistent swollen salivary glands |
| | | ☐ Frequently drink liquids to aid in swallowing dry food |
| | | Ocular signs (at least 1): |
| | | □ Schirmer's I test without anesthesia (≤ 5 mm in 5 minutes) |
| | | □ Rose Bengal score or other ocular dry eye score (≥ 4 with van Bijsterveld's scoring) |
| | | Histopathology: In minor salivary glands (obtained through normal-appearing mucosa) focal lymphocytic sialadenitis, evaluated by an expert histopathologist, with a focus score ≥ 1, defined as several lymphocytic foci (which are adjacent to normal-appearing mucous acini and contain more than 50 lymphocytes) per 4 mm² of glandular tissue |
| | | Salivary gland involvement (at least 1): |
| | | ☐ Unstimulated whole salivary flow (≤ 1.5 mL in 15 minutes) |
| | | ☐ Parotid sialography showing presence of diffuse sialectasis without evidence of obstruction in major ducts |
| | | □ Salivary scintigraphy showing delayed uptake, reduced concentration and/or delayed excretion of tracer |
| | | Autoantibodies present (Ro or SSA, La or SSB, or both) |
| | Co | onfirmed small fiber neuropathy by skin biopsy from distal leg |
| | Ba | seline pain and disability score documentation has been submitted: |
| | | Baseline Visual Analogue Scale (VAS): |
| | | Baseline Modified Rankin Scale (MRS): |
| D | iag | nosis: Small Fiber Sensory Neuropathy associated with Sjögren's Syndrome |
| <u>Rea</u> | uth | orization: 12 months (Dose: 0.4 g/kg/month) |
| | Member demonstrated a clinical response to therapy based on improvement from baseline score of the objective clinical measuring tool VAS or MRS | |
| | IV | IG dose has been tapered down to lowest effective dose since initial approval |
| | | |

(Continued on next page)

3

| Diagnosis: Other Peripheral Neuropathies associated with Sjögren's Syndrome | | | | | |
|---|---|---|--|--|--|
| Init | ial A | Authorization: 6 months (Dose: 2 g/kg/month) | | | |
| | Diagnosis based on American-European Consensus Group Criteria (presence of 4 out of 6 items if histopathology or autoantibodies positive) OR THREE (3) of the following are positive: ocular signs, histopathology, salivary gland involvement, autoantibodies): | | | | |
| | | Ocular symptoms (<u>at least 1</u>): | | | |
| | | ☐ Daily, persistent, troublesome dry eyes for more than 3 months | | | |
| | | ☐ Recurrent sensation of sand or gravel in eyes | | | |
| | | ☐ Use of tear substitutes more than 3 times a day | | | |
| | | Oral symptoms (<u>at least 1</u>): | | | |
| | | □ Daily feeling of dry mouth for more than 3 months | | | |
| | | ☐ Recurrent or persistent swollen salivary glands | | | |
| | | ☐ Frequently drink liquids to aid in swallowing dry food | | | |
| | | Ocular signs (at least 1): | | | |
| | | ☐ Schirmer's I test without anesthesia (≤ 5 mm in 5 minutes) | | | |
| | | \square Rose Bengal score or other ocular dry eye score (≥ 4 with van Bijsterveld's scoring) | | | |
| | | Histopathology: In minor salivary glands (obtained through normal-appearing mucosa) focal lymphocytic sialadenitis, evaluated by an expert histopathologist, with a focus score ≥ 1, defined as several lymphocytic foci (which are adjacent to normal-appearing mucous acini and contain more than 50 lymphocytes) per 4 mm² of glandular tissue | | | |
| | | Salivary gland involvement (at least 1): | | | |
| | | ☐ Unstimulated whole salivary flow (≤ 1.5 mL in 15 minutes) | | | |
| | | □ Parotid sialography showing presence of diffuse sialectasis without evidence of obstruction in major ducts | | | |
| | | □ Salivary scintigraphy showing delayed uptake, reduced concentration and/or delayed excretion of tracer | | | |
| | | Autoantibodies present (Ro or SSA, La or SSB, or both) | | | |
| | Pe | ripheral neuropathy confirmed by electromyography | | | |
| | | Sensorimotor | | | |
| | | Sensory ataxic neuropathy (ganglionopathy) | | | |
| | | Nonataxic sensory polyneuropathy | | | |
| | Ba | Baseline pain and disability score documentation has been submitted: | | | |
| _ | | Baseline Visual Analogue Scale (VAS): | | | |
| | | Baseline Modified Rankin Scale (MRS): | | | |
| | _ | · / | | | |

(Continued on next page)

1

| □ Diagnosis: Other Peripheral Neuropathies associated with Sjögren's Syndrome Reauthorization: 12 months (Dose: 0.4 g/kg/month) | | | | |
|--|--|--|--|--|
| | | | | |
| ☐ IVIG dose has been tapered down to lowest effective dose since initial approval | | | | |
| | | | | |
| Medication being provided by (check box below that applies): | | | | |
| □ Location/site of drug administration: | | | | |
| NPI or DEA # of administering location: | | | | |
| OR | | | | |
| □ Specialty Pharmacy – Proprium Rx | | | | |
| For urgent reviews: Practitioner should call Sentara Health Plans Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health Plan's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function. | | | | |
| **Use of samples to initiate therapy does not meet step edit/preauthorization criteria.** *Previous therapies will be verified through pharmacy paid claims or submitted chart notes. | | | | |
| | | | | |