# SENTARA COMMUNITY PLAN (MEDICAID)

## PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST\*

**Directions:** The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If the information provided is not</u> complete, correct, or legible, the authorization process can be delayed.

## Drug Requested: Orencia<sup>®</sup> SQ (abatacept) (Pharmacy)

## MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name:	
Member Sentara #:	Date of Birth:
Prescriber Name:	
Prescriber Signature:	Date:
Office Contact Name:	
Phone Number:	
DEA OR NPI #:	
DRUG INFORMATION: Authorization Drug Form/Strength:	n may be delayed if incomplete.
Dosing Schedule:	Length of Therapy:
Diagnosis:	ICD Code, if applicable:
Weight:	Date:
DIAGNOSIS	Recommended Dose

DIAGNOSIS	<b>Recommended Dose</b>
Moderate to severe Active Rheumatoid	SUBCUTANEOUS
Arthritis (RA)	• 125 mg once a week (4 syringe/28 days)
Psoriatic Arthritis (PsA)	Pediatric dosing:
	• Weighing 10 kg to less than 25 kg: 50 mg once a week (1 syringe/28 days)
	• Weighing 25 kg to less than 50 kg: 87.5 mg once a week (1 syringe/28 days)
	• Weighing 50 kg or greater: 125 mg once a week (4 syringes/28 days)

DIAGNOSIS	Recommended Dose
<ul> <li>Juvenile Idiopathic Arthritis (JIA) in members 2 years and older</li> </ul>	<ul> <li>SUBCUTANEOUS</li> <li>Weighing greater 50 kg: 125 mg once a week (4 syringes/28 days)</li> <li>Weighing 25 kg to less than 50 kg: 87.5 mg (Four syringes of 87.5mg/28days) once a week</li> <li>Weighing 10kg to 25 kg: 50 mg once a week (Four syringes of 50 mg/28 days)</li> </ul>

**CLINICAL CRITERIA:** Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

### **Diagnosis - check applicable diagnosis below or authorization will be denied.**

#### **D** Moderate to severe Active Rheumatoid Arthritis (RA)

- □ Prescriber is a Rheumatologist
- □ Patient has been diagnosed with moderate to severe rheumatoid arthritis
- □ Trial and failure of, contraindication, or adverse reaction to methotrexate
- □ Trial and failure of at least <u>ONE (1)</u> other <u>DMARD therapy</u> including, but not limited to: (check <u>each</u> tried)

auranofin	□ sulfasalazine
azathioprine	□ leflunomide
hydroxychloroquine	□ other:

□ Patient has tried and failed **TWO (2)** of the following biologics:

□ Humira <sup>®</sup> □	Enbrel <sup>®</sup>	Infliximab
-------------------------	---------------------	------------

(Continued on next page)

### **Juvenile Idiopathic Arthritis (JIA)**

- □ Prescriber is a Rheumatologist
- Department has been diagnosed with moderate to severe active Juvenile Idiopathic Arthritis (JIA)
- □ Patient is 2 years of age and older
- **□** Trial and failure of, contraindication, or adverse reaction to methotrexate
- □ Trial and failure of at least <u>ONE (1)</u> other <u>DMARD therapy</u> including, but not limited to: (check <u>each</u> tried):

auranofin	sulfasalazine
azathioprine	leflunomide
hydroxychloroquine	other:

#### □ Patient has tried and failed <u>TWO (2)</u> of the biologics below:

|--|

### □ Active Psoriatic Arthritis (PsA)

- Dependence of the Prescriber is a Rheumatologist
- □ Patient is 2 years of age and older
- Department has been diagnosed with moderate to severe active Psoriatic Arthritis (PsA)
- **□** Trial and failure of, contraindication, or adverse reaction to methotrexate

(Continued on next page)

□ Trial and failure of at least <u>ONE (1)</u> other <u>DMARD therapy</u> including, but not limited to: (check <u>each</u> tried):

auranofin	□ sulfasalazine
azathioprine	□ leflunomide
hydroxychloroquine	□ other:

□ Patient has tried and failed <u>TWO (2)</u> of the following biologics:

□ Humira <sup>®</sup>	□ Enbrel <sup>®</sup>	Infliximab
-----------------------	-----------------------	------------

Medication being provided by (check applicable box(es) below):

Physician's office

 OR
 Specialty Pharmacy – PropriumRx

\*Use of samples to initiate therapy does not meet step-edit/preauthorization criteria.\* \*Previous therapies will be verified through pharmacy paid claims or submitted chart notes.\*