

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

Drug Requested: Orencia® SQ (abatacept) **(Pharmacy)**

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

DIAGNOSIS	Recommended Dose
<input type="checkbox"/> Moderate to severe Active Rheumatoid Arthritis (RA)	SUBCUTANEOUS • 125 mg once a week (4 syringe/28 days)
<input type="checkbox"/> Psoriatic Arthritis (PsA)	
<input type="checkbox"/> Juvenile Idiopathic Arthritis (JIA) in members 2 years and older	SUBCUTANEOUS • Weighing greater 50 kg: 125 mg once a week (4 syringe/ 28 days) • Weighing 25 kg to less than 50 kg: 87.5 mg (Four syringes of 87.5mg/28days) once a week • Weighing 10kg to 25 kg: 50 mg once a week (Four syringes of 50mg/28days)

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Diagnosis - check applicable diagnosis below or authorization will be denied.

☐ **Moderate to severe Active Rheumatoid Arthritis (RA)**

- ☐ Prescriber is a Rheumatologist

AND

- ☐ Patient has been diagnosed with moderate to severe rheumatoid arthritis

AND

- ☐ Trial and failure of, contraindication, or adverse reaction to methotrexate

AND

- ☐ Trial and failure of at least **ONE (1)** other **DMARD therapy** including, but not limited to: (check each tried)

<input type="checkbox"/> auranofin	<input type="checkbox"/> sulfasalazine
<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> other: _____

AND

- ☐ Patient has tried and failed **TWO (2)** of the following biologics:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

☐ **Juvenile Idiopathic Arthritis (JIA)**

- ☐ Prescriber is a Rheumatologist

AND

- ☐ Patient has been diagnosed with moderate to severe active Juvenile Idiopathic Arthritis (JIA)

AND

- ☐ Patient is 2 years of age and older

AND

- ☐ Trial and failure of, contraindication, or adverse reaction to methotrexate

AND

(Continued on next page)

- ☐ Trial and failure of at least **ONE (1)** other **DMARD therapy** including, but not limited to: (**check each tried**):

<input type="checkbox"/> auranofin	<input type="checkbox"/> sulfasalazine
<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> other: _____

AND

- ☐ Patient has tried and failed **TWO (2)** of the biologics below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®
----------------------------------	----------------------------------

☐ **Active Psoriatic Arthritis (PsA)**

- ☐ Prescriber is a Rheumatologist

AND

- ☐ Patient has been diagnosed with moderate to severe active Psoriatic Arthritis (PsA)

AND

- ☐ Trial and failure of, contraindication, or adverse reaction to methotrexate

AND

- ☐ Trial and failure of at least **ONE (1)** other **DMARD therapy** including, but not limited to: (**check each tried**):

<input type="checkbox"/> auranofin	<input type="checkbox"/> sulfasalazine
<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> other: _____

AND

- ☐ Patient has tried and failed **TWO (2)** of the following biologics:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

Medication being provided by (check applicable box(es) below):

- ☐ Physician's office OR ☐ Specialty Pharmacy – PropriumRx

****Use of samples to initiate therapy does not meet step-edit/preauthorization criteria.****
****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****