

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692.** No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If information provided is not complete, correct, or legible, authorization will be delayed.**

Drug Requested: Actemra[®] SQ (tocilizumab) (**self-administered**) (**Pharmacy**) (**Non-Preferred**)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

DIAGNOSIS	Recommended Dose
<input type="checkbox"/> Rheumatoid Arthritis (RA)	SUBCUTANEOUS <ul style="list-style-type: none"> • Weight <100kg: Two syringes per 28 days. Max dose is 4 syringes per 28 days • Weight >100kg: Four syringes per 28 days
<input type="checkbox"/> Polyarticular Juvenile Idiopathic Arthritis (PJIA)	SUBCUTANEOUS <ul style="list-style-type: none"> • Weight <30kg: 162mg/dose once every 3 weeks • Weight ≥30kg: 162mg/dose once every 2 weeks
<input type="checkbox"/> Systemic Juvenile Idiopathic Arthritis (SJIA)	SUBCUTANEOUS <ul style="list-style-type: none"> • Weight <30kg: 162mg/dose once every 3 weeks • Weight ≥30kg: 162mg/dose every 2 weeks
<input type="checkbox"/> Giant Cell Arteritis (GCA)	SUBCUTANEOUS <ul style="list-style-type: none"> • Max adult dose is 4 syringes per 28 days
<input type="checkbox"/> Systemic Sclerosis Associated Interstitial Lung Disease (SSc-ILD)	SUBCUTANEOUS <ul style="list-style-type: none"> • Max adult dose is 4 syringes per 28 days

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Diagnosis: Rheumatoid Arthritis (RA)

- Prescriber is a Rheumatologist; **AND**
- Member has moderate to severe rheumatoid arthritis; **AND**
- Tried and failed methotrexate; **OR**
- Requested medication will be used in conjunction with methotrexate; **OR**
- Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or other contraindication); **AND**
- Tried and failed at least **one DMARD** other than methotrexate and (**check each tried**)

<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> auranofin	<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> gold salts
<input type="checkbox"/> d-penicillamine	<input type="checkbox"/> cyclosporine	<input type="checkbox"/> cyclophosphamide
<input type="checkbox"/> tacrolimus	<input type="checkbox"/> Other: _____	

AND

- Trial and failure of **TWO (2)** of the **PREFERRED** biologics below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®	<input type="checkbox"/> Infliximab
----------------------------------	----------------------------------	-------------------------------------

Diagnosis: Polyarticular Juvenile Idiopathic Arthritis (PJIA)

- Prescriber is a Rheumatologist; **AND**
- Member must be 2 years of age and older with active polyarticular juvenile idiopathic arthritis; **AND**
- Tried and failed methotrexate; **OR**
- Requested medication will be used in conjunction with methotrexate; **OR**
- Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or other contraindication); **AND**
- Trial and failure of **TWO (2)** of the **PREFERRED** biologics below:

<input type="checkbox"/> Humira®	<input type="checkbox"/> Enbrel®
----------------------------------	----------------------------------

AND

(Continued on next page)

- Trial and failure of at least **ONE (1) DMARD** therapy **and** (check each tried)

<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> auranofin	<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> gold salts
<input type="checkbox"/> d-penicillamine	<input type="checkbox"/> cyclosporine	<input type="checkbox"/> cyclophosphamide
<input type="checkbox"/> tacrolimus	<input type="checkbox"/> Other: _____	

Diagnosis: Systemic Juvenile Idiopathic Arthritis (SJIA)

- Prescriber is a Rheumatologist; **AND**
- Member must be 2 years of age and older with active systemic juvenile idiopathic arthritis; **AND**
- Tried and failed methotrexate; **OR**
- Requested medication will be used in conjunction with methotrexate; **OR**
- Member has a contraindication to methotrexate (e.g., alcohol abuse, cirrhosis, chronic liver disease, or other contraindication); **AND**
- Trial and failure of at least **ONE (1) DMARD** therapy **and** (check each tried)

<input type="checkbox"/> sulfasalazine	<input type="checkbox"/> azathioprine	<input type="checkbox"/> leflunomide
<input type="checkbox"/> auranofin	<input type="checkbox"/> hydroxychloroquine	<input type="checkbox"/> gold salts
<input type="checkbox"/> d-penicillamine	<input type="checkbox"/> cyclosporine	<input type="checkbox"/> cyclophosphamide
<input type="checkbox"/> tacrolimus	<input type="checkbox"/> Other: _____	

Diagnosis: Giant Cell Arteritis (GCA)

- Member must be 18 years of age and older with giant cell arteritis (GCA) diagnosis

Diagnosis: Systemic Sclerosis-Associated Interstitial Lung Disease (SSc-ILD)

- Member must be 18 years of age and has a confirmed diagnosis of systemic sclerosis-associated interstitial lung disease (SSc-ILD)

Medication being provided by (check applicable box(es) below):

- Physician's office **OR** Specialty Pharmacy – PropriumRx

****Use of samples to initiate therapy does not meet step-edit/preauthorization criteria.****
****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****