

SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; **fax to 1-800-750-9692**. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. **If the information provided is not complete, correct, or legible, the authorization process can be delayed.**

Botulinum Toxin Injections[®], Type A

Drug Requested: (check applicable drug below)

☐ **Botox[®]** (onabotulinumtoxinA)

☐ **Xeomin[®]** (incobotulinumtoxinA)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight (if applicable): _____ Date weight obtained: _____

- **Maximum quantity limits:** 400 units in a 3-month period
- **Cosmetic indications are EXCLUDED**

NOTE: In treating adult patients for one or more indications, the maximum cumulative dose should not exceed 400 units, in a 3-month interval. In pediatric patients, the total dose should not exceed the lower of 10 units/kg body weight or 340 units, in a 3-month interval.

(Continued on next page)

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- ☐ Member must have ONE of the following diagnoses:
 - ☐ **Achalasia, Primary idiopathic esophageal AND meets ONE of the following:**
 - ☐ Member failed or had a clinically significant adverse reaction to conventional therapy (nitrates or calcium channel blockers)
 - ☐ Member is ineligible for surgical treatment due to advance age or multiple co-morbidities (poor surgical risk)
 - ☐ Member is at high risk of complications of pneumatic dilation or surgical myotome
 - ☐ Failure of prior myotomy or dilation
 - ☐ Member has an epiphrenic diverticulum or hiatal hernia, both of which increase the risk of dilation induced perforation
 - ☐ **Achalasia, Internal anal sphincter (IAS) AND meets BOTH of the following:**
 - ☐ Member has NOT responded to treatment with laxatives
 - ☐ Member has NOT responded to **OR** is NOT a candidate for anal sphincter myectomy
 - ☐ **Anal Fissure – Chronic AND meets the following criteria:**
 - ☐ Member failed (at least 60 days) topical nitroglycerin or topical calcium channel blocker
 - ☐ **Blepharospasm**
 - ☐ **Cerebral Palsy – Dynamic Contracture**
 - ☐ **Cerebral Palsy – Spasticity** (including diplegia, hemiplegia, paraplegia, or quadriplegia)
 - ☐ **Cervical Dystonia** (spasmodic torticollis) and **Mixed Cervical Dystonia**
 - ☐ **CVA-related spasticity** within 1 year of onset
 - ☐ **Drooling in Parkinson's disease**
 - ☐ **Essential hand tremor in patients who fail oral agents**
 - ☐ **Hand Dystonia**
 - ☐ **Hemifacial spasm**
 - ☐ **Motor tics**

(Continued on next page)

- ☐ **Neurogenic detrusor overactivity (NDO) and/or detrusor sphincter dyssynergia **AND** meets ALL the following:**
 - ☐ Member has diagnosis of incontinence due to NDO or detrusor sphincter dyssynergia associated with a neurologic condition (e.g., multiple sclerosis, spinal cord injury, brain injury) that has been confirmed by urodynamic testing (**submit documentation of diagnosis**)
 - ☐ Member meets ONE of the following age and prerequisite therapy requirements:
 - ☐ **For members aged 5-17 years:** Member has had a 30-day trial and failure of oxybutynin (oral or intravesical use) and one other oral systemic medication from the following classes: anticholinergics or beta-3 antagonists (**Must submit chart notes documenting therapy failures**)
 - ☐ **For members aged 17 years and older:** Member has had a 30-day trial and failure of two oral systemic medications from the following classes: anticholinergics or beta-3 antagonists (**Must submit chart notes documenting therapy failures**)
- ☐ **Orofacial Dyskinesia**
- ☐ **Overactive Bladder **AND** meets ALL the following**
 - ☐ Diagnosis of incontinence
 - ☐ Symptoms of urge incontinence, urgency, and frequency (experienced at least 3 urinary incontinence episodes and at least 24 micturitions in 3 days)
 - ☐ 8-12 week trial and failure of behavioral therapy (e.g., bladder training, control strategies, pelvic floor muscle training, fluid management)
 - ☐ Failed or inadequate response to anticholinergic therapy within the last 9 months (4-8 week trial per agent)
 - ☐ Trial and failure of ONE of the following medication regimens:
 - ☐ 2 anticholinergic agents and 1 β -3 adenosine receptor agonist (**requires PA**)
 - ☐ 1 anticholinergic agent and 1 alpha blocker and 1 β -3 adenosine receptor agonist (**requires PA**)Please indicate drugs used: _____
- ☐ **Strabismus** (injections done in lieu of coverage for surgery)
- ☐ **Synkinetic Eyelid Closure – VII Cranial Nerve**
- ☐ **Torticollis**

Medication being provided by: Please check applicable box below.

- ☐ **Physician's office** **OR** ☐ **Specialty Pharmacy**

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****
****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****