SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid)</u> on this request. All other information may be filled in by office staff; <u>fax to 1-800-750-9692</u>. No additional phone calls will be necessary if all information <u>(including phone and fax #s)</u> on this form is correct. <u>If the information provided is not complete, correct, or legible, the authorization process can be delayed.</u>

Hereditary Angioedema (HAE)

Drug Requested: (check box below that applies)

PREFERRED Medications (with Quantity Limits)					
□ Berinert ® - 4 vials per attack (plus 4 for emergency)	□ Cinryze [®] days	- 20 vials per 34	icatibant - 1 dose per attack (plus 1 for emergency)		
□ Kalbitor ® - 3 vials per attack (plus 3 for emergency) (see Black Box warning below)	· ·	- 1 dose per attack emergency)			
Non-Preferred Medications (with Quantity Limits)					
□ Firazyr ® - 1 dose per attack (plus 1 for emergency)		□ Haegarda ® - 2,000 IU SDV kit (16 kits per 28 days) & 3,000 IU SDV kit (8 kits per 28 days)			
□ Orladeyo [™] - 1 capsule per day		□ Ruconest® emergency)	- 2 vials per attack (plus 2 for		
□ Takhzyro [™] - 2 vials per 28 days					

Black Box Warning: Because of the risk of anaphylaxis, KALBITOR® should only be administered by a healthcare professional with appropriate medical support to manage anaphylaxis and hereditary angioedema.

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.			
Member Name:			
Member Sentara #:			
Prescriber Name:			
Prescriber Signature:			
Office Contact Name:			
Phone Number:	Fax Number:		
NPI #:			

(Continued on next page)

DRUG INFORMATION: Authorization may be delayed if incomplete.				
Drug Name/Form/Strength:				
Dosing Schedule:	Length of Therapy:			
Diagnosis:				
Weight (if applicable): Date				
CLINICAL CRITERIA: Check below all that apply support each line checked, all documentation, including la provided or request may be denied.				
 Has the recipient's diagnosis of HAE been confirme (type I or II HAE) as documented by one of the followallow C1 inhibitor (C1-INh) antigenic level below the performing the test 				
OR				
 C1-INh functional level below the lower limit of test 	normal as defined by the laboratory performing the			
 Was the medication prescribed by, or in consultation hematology, pulmonology, or medical genetics? If YES, document the physician's specialty: 	□ Yes □ No			
TREATMENT OF ACUTE HAE ATTACKS				
Berinert [®] (C1 esterase inhibitor), Firazyr [®] (icatibant), icatib inhibitor), Sajazir [™] (icatibant)	ant, Kalbitor® (ecallantide), Ruconest® (C1 esterase			
1. Will the requested medication be used as monothera	py to treat acute HAE attacks?			
	□ Yes □ No			
PROPHYLAXIS OF HAE ATTACKS				
Cinryze® (C1 esterase inhibitor), Haegarda® (C1 esterase in (ianadelumab-flyo)	hibitor), Orladeyo® (berotralstat), Takhzyro®			
1. Will the requested medication be used for prophylax	is of HAE attacks?			
(Continued on	next page)			

List pharmaceutical drugs attempted and outcome:	
Medical Necessity: Provide clinical evidence that the preferred drug(s) will not provide adequate benefined and/or provide clinical rationale for quantity exception requests:	ĭt

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. **

*Previous therapies will be verified through pharmacy paid claims or submitted chart notes. *