

# ASAM Level 2.1 Intensive Outpatient Services for Substance Abuse (Adolescent) Initial

### **Table of Content**

**Purpose** 

**Description & Definitions** 

Criteria

Coding

**Document History** 

References

Special Notes

**Keywords** 

6/2023

Next Review Date 6/2024

Coverage Policy BH 48

Version 1

All requests for authorization for the services described by this medical policy will be reviewed per Early and Periodic Screening, Diagnostic and Treatment (EPSDT) guidelines. These services may be authorized under individual consideration for Medicaid members under the age of 21-years if the services are judged to be medically necessary to correct or ameliorate the member's condition. Department of Medical Assistance Services (DMAS), Supplement B - EPSDT (Early and Periodic Screening, Diagnosis and Treatment) Manual.

# Purpose:

This policy addresses ASAM Level 2.1 Intensive Outpatient Services for Substance Abuse (Adolescent) Initial.

# **Description & Definitions:**

Intensive outpatient programs (IOPs) generally provide 9-19 hours of structured programming per week for adults and 6-19 hours for adolescents, consisting primarily of counseling and education about addiction-related and mental health problems. The patient's needs for psychiatric and medical services are addressed through consultation and referral arrangements if the patient is stable and requires only maintenance monitoring. (Services provided outside the primary program must be tightly coordinated.)

There are occasions when the individuals progress in the IOP no longer requires nine hours per week of treatment in such cases, less than nine hours per week for adults and six hours per week for adolescents as a transition step down in intensity should be considered as a continuation of the IOP program for one or two weeks. Such continuity allows for a smoother transition to Level 1 to avoid exacerbation and recurrence of signs and symptoms.

Intensive outpatient treatment differs from partial hospitalization (Level 2.5) programs in the intensity of clinical services that are directly available. Specifically, most intensive outpatient programs have less capacity to effectively treat patients who have substantial unstable medical and psychiatric problems than do partial hospitalization programs.

### Criteria:

Adolescent Admission to Intensive outpatient level of care for substance-related disorder is considered medically necessary when the following ASAM dimensions are met. The member meets ASAM criteria for intensive outpatient level of care when dimensions 1 and 2 and one of 3, 4,5 and 6 are met

- Individual is under the age of 18 years old and their mental and behavioral health status meets ALL of the following:
  - <u>Diagnosis</u>: The individual has at least one diagnosis from the most recent Diagnostic and Statistical Manual of Mental Disorders for Substance-Related and Addictive Disorders with the exception of tobaccorelated disorders, caffeine use disorder or dependence, and nonsubstance-related addictive disorders
  - o Dimension 1: Acute intoxication and/or withdrawal situation meets All of the following:
    - The individual is able to tolerate mild subacute withdrawal symptoms
    - The individual has made a commitment to sustain treatment and to follow treatment recommendations
    - The individual has external supports (family and/or court) that promote engagement in treatment
  - <u>Dimension 2</u>: The individual's biomedical conditions (i.e., hypertension, diabetes, asthma, etc.), if any, are stable or are being concurrently addressed or do not interfere with treatment at this level of care
- Individual is under the age of 18 years old and their mental and behavioral health status meets one of the following:
  - <u>Dimension 3</u>: The individual's current emotional, behavioral, or cognitive status meets 1 or more of the following:
    - There is mild risk of behaviors endangering self, others or property and requires frequent monitoring
    - Recovery efforts are affected by current status and require mild interference at this intensity
      of this level of care to support treatment engagement
    - Mild to moderate impairment in social functioning, but can sustain responsibilities
    - The adolescent is experiencing mild to moderate difficulties with activities of daily living, and requires frequent monitoring or interventions
    - The adolescent's history (combined with the present situation) predicts the need for frequent monitoring or interventions
  - <u>Dimension 4</u>: The individual's readiness to change requires close monitoring and support to promote change which is demonstrated by 1 or more of the following:
    - The individual requires structured therapy and a programmatic milieu to promote progress through the stages of change, as evidenced by behaviors including 1 or more of the following:
      - The individual is verbally compliant, but does not demonstrate consistent behaviors
      - The individual is only passively involved in treatment
      - The individual demonstrates variable adherence with attendance in outpatient sessions or support groups
    - The individual's perspective inhibits his/her ability to make progress through the stages of change. For example, he/she has unrealistic expectations that the program will resolve quickly and with little to no effort, or does not recognize the need for assistance.
  - Dimension 5: Significant risk of relapse or continued use potential is indicated by 1 or more of the following:
    - Although the individual has been an active participant at a less intensive level of care, he/she is experiencing an intensification of symptoms of substance-related disorder and his/her level of functioning is deteriorating despite modification of the treatment plan.
    - There is a high likelihood that the individual will continue to use or relapse to use of alcohol and/or other drugs or gambling without close outpatient monitoring and structured therapeutic

services, as indicated by his/her lack of awareness of relapse triggers, difficulty in coping, or in postponing immediate gratification or ambivalence toward treatment.

- Dimension 6: The individual's environment is impeding recovery as evidenced by 1 or more of the following:
  - Continued exposure to the individual's current school, work or living environment will render recovery unlikely. The individual lacks the resources or skills necessary to maintain an adequate level of functioning without intensive outpatient services.
  - The individual lacks social contacts, has unsupportive social contacts that jeopardize recovery, or has few friends or peers who do not use alcohol or other drugs. The individual lacks the resources or skills necessary to maintain an adequate level of functioning without intensive outpatient services.
  - Family conflicts and related family dysfunction impede the individual's ability to learn the skills necessary to achieve and maintain abstinence.

There is insufficient scientific evidence to support the medical necessity of partial hospitalization services for substance abuse for uses other than those listed in the clinical indications for procedure section.

#### **Service Units and Limitations:**

- Members shall be discharged from this service when other less intensive services may achieve stabilization, the
  member requests discharge, the member ceases to participate, or the member demonstrates a need for a higher
  level of care. Discharge planning shall document realistic plans for the continuity of MOUD services with an innetwork Medicaid provider.
- Intensive Outpatient Services may not be authorized concurrently with ASAM Level 2.1, 3.3, 3.5, 3.7 or 4.0;
   Mental Health Services including Mental Health Intensive Outpatient Services, Mental Health Partial
   Hospitalization Programs, Psychosocial Rehabilitation, Therapeutic Day Treatment, Intensive In-Home Services,
   Therapeutic Group Home, Community Stabilization, Residential Crisis Stabilization Unit (RCSU), Assertive
   Community Treatment, Multisystemic Therapy, Functional Family Therapy, Psychiatric Residential Treatment or
   inpatient admission. A seven day overlap with any outpatient or community based behavioral health service may
   be allowed for care coordination and continuity of care.
- One unit of service is one day with a minimum of 3 service hours per service day to achieve an average of 9 to 19 hours of services per week for adults and an average of 6 to 19 hours of services per week for children and adolescents, with regards to the first and last week of treatment. A maximum average of 19 hours shall be billed per week.
- ASAM Criteria allows for less than an average of 9 hours per week for adults and an average of 6 hours per week
  for adolescents as a transition step down in intensity for 1 to 2 weeks prior to transitioning to Level 1 to avoid
  relapse. The transition step down needs to be approved by the MCO or the BHSA (based on the member's
  benefit) and documented and supported by the member's ISP.
- Group substance use counseling by CATPs, CSACs and CSAC supervisees shall have a maximum limit of 12
  individuals in the group or less depending on the clinical model. Group size may exceed this limit based on the
  clinical determination of the CATP. Such counseling shall focus on the needs of the members served.
- CSACs and CSAC-supervisees, by scope of practice, are able to perform group substance use counseling, thus
  could provide counseling and psychoeducational services in this level of care.
- There are no maximum annual limits.

# <u>Discharge/Transfer Criteria It is appropriate to transfer or discharge the member from the present level of care if he or she meets the following criteria:</u>

- The member has achieved the goals articulated in the ISP, thus resolving the problem(s) that justified admission to the current level of care; or
- The member has been unable to resolve the problem(s) that justified admission to the present level of care, despite amendments to the ISP. Treatment at another level of care or type of service therefore is indicated; or

- The member has demonstrated a lack of capacity to resolve his or her problem(s). Treatment at another level of care or type of service therefore is indicated; or
- The member has experienced an intensification of his or her problem(s), or has developed a new problem(s), and can be treated effectively only at a more intensive level of care.

## Coding:

Medically necessary with criteria:

Coding	Description
H0015	Alcohol and/or drug services; intensive outpatient (treatment program that operates at least 3 hours/day and at least 3 days/week and is based on an individualized treatment plan), including assessment, counseling; crisis intervention, and activity therapies or education

Considered Not Medically Necessary:

Coding	Description
	None

U.S. Food and Drug Administration (FDA) - approved only products only.

# Document History:

**Revised Dates:** 

**Reviewed Dates:** 

Effective Date:

- June 2023

### References:

Including but not limited to: Specialty Association Guidelines; Government Regulations; Winifred S. Hayes, Inc; UpToDate; Literature Review; Specialty Advisors; National Coverage Determination (NCD); Local Coverage Determination (LCD).

- 1. DMAS Manual- Addiction and Recovery Treatment Services
- 2. DMAS Medallion 4.0 Contract: Section 8.2.A, 8.2.B
- 3. DMAS CCC Plus Contract: Section 4.2.4
- 4. Cardinal Care Contract: Section 5.5.6
- 5. MCG 26th Edition: https://careweb.carequidelines.com/ed26/index.html
- 6. American Society of Addiction Medicine (ASAM) Edition 3

# Special Notes: \*

This medical policy express Sentara Health Plan's determination of medically necessity of services, and they are based upon a review of currently available clinical information. These policies are used when no specific guidelines for coverage are provided by the Department of Medical Assistance Services of Virginia (DMAS). Medical Policies may be superseded by state Medicaid Plan guidelines. Medical policies are not a substitute for clinical judgment or for any prior authorization requirements of the health plan. These policies are not an explanation of benefits.

Medical policies can be highly technical and complex and are provided here for informational purposes. These medical policies are intended for use by health care professionals. The medical policies do not constitute medical advice or medical care. Treating health care professionals are solely responsible for diagnosis, treatment and medical advice. Sentara Health Plan members should discuss the information in the medical policies with their treating health care professionals. Medical technology is constantly evolving and these medical policies are subject to change without notice, although Sentara Health Plan will notify providers as required in advance of changes that could have a negative impact on benefits.

The Early and Periodic Screening, Diagnostic and Treatment (EPSDT) covers services, products, or procedures for children, if those items are determined to be medically necessary to "correct or ameliorate" (make better) a defect, physical or mental illness, or condition (health problem) identified through routine medical screening or examination, regardless of whether coverage for the same service or support is an optional or limited service under the state plan. Children enrolled in the FAMIS Program are not eligible for all EPSDT treatment services. All requests for authorization for the services described by this medical policy will be reviewed per EPSDT guidelines. These services may be authorized under individual consideration for Medicaid members under the age of 21-years if the services are judged to by medically necessary to correct or ameliorate the member's condition. Department of Medical Assistance Services (DMAS), Supplement B - EPSDT (Early and Periodic Screening, Diagnosis and Treatment) Manual.

## Keywords:

IOP, Intensive Outpatient Program, substance abuse, alcoholism, SHP behavioral health 48, withdrawal, intoxication, relapse, drug abuse, alcohol abuse, SHP Intensive Outpatient Program for Substance Abuse, ASAM Level 2.1, Adolescent, Initial Medicaid