SENTARA HEALTH PLANS

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions</u>: <u>The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request</u>. All other information may be filled in by office staff; fax to <u>1-800-750-9692</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization may be delayed.</u>

<u>Drug Requested</u>: Camzyos[®] (mavacamten)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.			
Member Name:			
	Date of Birth:		
Prescriber Name:			
Prescriber Signature:			
Office Contact Name:			
Phone Number:	Fax Number:		
DEA OR NPI #:			
DRUG INFORMATION: Authoriz	ation may be delayed if incomplete.		
Drug Form/Strength:			
Dosing Schedule:			
Diagnosis:	ICD Code:		
Weight:	Date:		
Quantity limit: 1 capsule per day			
	ow all that apply. All criteria must be met for approval. To support uding lab results, diagnostics, and/or chart notes, must be provided		
Initial Authorization: 8 months			
☐ Member is 18 years of age or older			
☐ Prescribed by or in consultation with	h a cardiologist specialist		
 Member has a diagnosis of sympton 	natic obstructive hypertrophic cardiomyopathy (HCM)		
☐ Member had an adequate echocardic	ogram or cardiovascular magnetic resonance imaging (CMR)		
☐ Member has New York Heart Associ	ciation (NYHA) class II-III symptoms		

(Continued on next page)

	Baseline peak oxygen consumption (pVO2) determined by cardiopulmonary exercise testing (CPET) has been submitted		
	Member has documented left ventricular ejection fraction (LVEF) $\geq 55\%$		
	Member has a left ventricular outflow track (LVOT) gradient of 50 mmHg or higher		
	Member remains symptomatic despite trial of, or intolerant to at least <u>TWO</u> of the following (verified be chart notes or pharmacy paid claims):		
		Beta-blocker (e.g., metoprolol, carvedilol)	
		Calcium channel blocker (e.g., verapamil, diltiazem)	
		disopyramide	
		Septal reduction therapy	
	Member will avoid concomitant use with moderate to strong CYP2C19 inhibitors/inducers, strong CYP3A4 inhibitors/inducers		
	Member will avoid concomitant dual therapy with beta-blockers and calcium channel blockers or monotherapy with disopyramide or ranolazine		
	Me	mber will NOT take disopyramide in conjunction with the requested medication	
ach 1	ine	orization: 12 months. All criteria that apply must be checked for approval. To support checked, all documentation (lab results, diagnostics, and/or chart notes) must be provided may be denied.	
	Me	mber has experienced continued clinical benefit as demonstrated by at least ONE of the following:	
		Improvement of at least 1.5 mL/kg/min in peak oxygen consumption (pVO2) as determined by cardiopulmonary exercise testing (CPET) AND a reduction of \geq 1 New York Heart Association (NYHA) functional classification (e.g., I, II, III, or IV)	
		Improvement of at least 3.0 mL/kg/min in pVO2 with no worsening in NYHA functional classification	
	Member has <u>NOT</u> experienced any treatment-restricting adverse effects (e.g., heart failure, LVEF <50% while taking requested medication		
		vider has submitted the results of member's most recent echocardiogram or cardiovascular magnetic onance imaging obtained after starting the requested medication	
1ed	icat	ion being provided by a Specialty Pharmacy - PropriumRx	

**Use of samples to initiate therapy does not meet step edit/ preauthorization criteria. ** *Previous therapies will be verified throu.gh pharmacy paid claims or submitted chart notes.*

REVISED/UPDATED: 8/7/2022; 10/4/2022; 6/15/2023

^{*}Approved by Pharmacy and Therapeutics Committee: 7/21/2022; 5/25/2023