

SENTARA COMMUNITY PLAN (MEDICAID)

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-750-9692. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. If the information provided is not complete, correct, or legible, the authorization process can be delayed.

Drug Requested: Orkambi[®] (ivacaftor/lumacaftor)

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.

Member Name: _____

Member Sentara #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

DRUG INFORMATION: Authorization may be delayed if incomplete.

Drug Form/Strength: _____

Dosing Schedule: _____ Length of Therapy: _____

Diagnosis: _____ ICD Code, if applicable: _____

Weight: _____ Date: _____

(Continued on next page)

Recommended Dosing:

Age	Weight	Dose	Administration
1 through 2 years	7 kg to < 9kg	1 packet of lumacaftor 75 mg/ivacaftor 94 mg granules	Mixed with one teaspoon (5 mL) of soft food or liquid and administered orally every 12 hours with fat-containing food
	9kg to < 14kg	1 packet of lumacaftor 100 mg/ivacaftor 125 mg granules	
	≥ 14 kg	1 packet of lumacaftor 150 mg/ivacaftor 188 mg granules	
2 through 5 years	< 14 kg	1 packet of lumacaftor 100 mg/ivacaftor 125 mg granules	
	≥ 14 kg	1 packet of lumacaftor 150 mg/ivacaftor 188 mg granules	
6 through 11 years		2 tablets of lumacaftor 100 mg/ivacaftor 125 mg (lumacaftor 200 mg/ivacaftor 250 mg per dose)	
12 years and older		2 tablets of lumacaftor 200 mg/ivacaftor 125 mg (lumacaftor 400 mg/ivacaftor 250 mg per dose)	

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

Initial Authorization: 6 months

- Member is **1 years of age or older** with a diagnosis of Cystic Fibrosis
- Member is confirmed to be homozygous for the F508del gene mutation of the CFTR protein in the cystic fibrosis transmembrane conductance regulator (CFTR) confirmed by an FDA-cleared test (**test results must be attached**)
- Prescribing physician is a pulmonologist or has consulted with a pulmonologist who specializes in the treatment of Cystic Fibrosis
- Baseline FEV1 completed within the last 30 days must be submitted (**test results must be attached**), unless the member is unable to perform a pulmonary function test (**documentation required**)
- Baseline LFTs have been completed prior to initiating therapy and will be completed annually (**labs must be attached**)
- Number of pulmonary exacerbations or hospitalizations in the preceding 6 months must be noted:

(Continued on next page)

- Baseline body mass index must be noted: _____
- Attestation that baseline ophthalmic examination to monitor lens opacities/cataracts has been completed for pediatric members
- Member will **NOT** take Orkambi[®], in combination with any other CFTR modulator therapy (i.e. Symdeko[®], Kalydeco[®], Trikafta[®]); concurrent therapy with these agents will not be approved
- Member will avoid concomitant use of strong CYP3A inducers (e.g., rifampin, carbamazepine, phenytoin, phenobarbital, St. John’s Wort) and strong or moderate CYP3A inhibitors (e.g., fluconazole, itraconazole)

Reauthorization: 12 months. Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

- Member continues to meet all initial authorization criteria
- Member has demonstrated disease response as indicated by **at least ONE** of the following (**select all that apply; must submit current labs and chart notes**):
 - Decreased pulmonary exacerbations or hospitalizations compared to pretreatment baseline
 - Stabilization of lung function as measured by FEV1 within the last year compared to baseline
 - Improvement in quality of life, weight gain, or growth
- Member has **NOT** received a lung transplant
- Member has experienced an absence of unacceptable toxicity from therapy (i.e. elevated transaminases (ALT or AST), development of cataracts or lens opacities)

Date of initiation of Orkambi [®] therapy: _____	Reauthorization Date: _____
Baseline FEV1 (last FEV1 prior to starting Orkambi [®]): _____	Current FEV1 (FEV1 AFTER last dose of Orkambi [®]): _____
Baseline Weight: _____	Current Weight: _____
BMI Baseline: _____	Current BMI: _____
Number of hospitalizations since last approval of Orkambi [®] must be noted: _____	

Medication being provided by Specialty Pharmacy - PropriumRx

*****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.*****

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****