SENTARA COMMUNITY PLAN (MEDICAID)

MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

<u>Directions:</u> The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; <u>fax to 1-844-305-2331</u>. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. <u>If information provided is not complete, correct, or legible, authorization can be delayed</u>.

Noninfectious Uveitis (NIU) Drugs (MEDICAL)

<u>Drug Requested</u>: (Check box below that applies)

□ Retisert® (fluocinolone acetonide intravitreal □ Yutiq[®] (fluocinolone acetonide intravitreal implant, 0.59 mg) (J7311) implant, 0.18 mg) (**J7314**) • 1 package: 0.59 mg implant = 59 billable • 1 package: 0.18 mg implant = 18 billable units every 30 months units every 36 months Quantity Limit: 2 implants every 30 months Quantity Limit: 2 implants every 36 months (1 implant per eye) (1 implant per eye) • Max Units (per dose): 118 billable units per • Max Units (per dose): 36 billable units every 36 months 30 months

MEMBER & PRESCRIBER INFORMATION: Authorization may be delayed if incomplete.				
Member Name:				
Member Sentara #:	Date of Birth:			
Prescriber Name:				
Prescriber Signature:	Date:			
Office Contact Name:				
	Fax Number:			
DEA OR NPI #:				
DRUG INFORMATIO	N: Authorization may be delayed if incomplete.			
Drug Form/Strength:				
	Length of Therapy:			
Diagnosis:	ICD Code, if applicable:			
Weight:	Date:			
	king this box, the timeframe does not jeopardize the life or health of the member regain maximum function and would not subject the member to severe pain.			
	□ Left Eye □ Right Eye □ Both Eyes			

(Continued on next page)

(Continued on next page)

Note: Sentara Health Plans considers the use of concomitant therapy with Dextenza[®], Ozurdex[®], Iluvien[®], Retisert[®], Xipere[™], or Yutiq[®], to be experimental and investigational. Safety and efficacy of these combinations have NOT been established and will NOT be permitted. In the event a member has an active Dextenza[®], Ozurdex[®], Iluvien[®], Retisert[®], Xipere[™], or Yutiq[®] authorization on file, all subsequent requests for an additional ophthalmic corticosteroid injection will NOT be approved.

CLINICAL CRITERIA: Check below all that apply. All criteria must be met for approval. To support each line checked, all documentation, including lab results, diagnostics, and/or chart notes, must be provided or request may be denied.

SECTION A.				
SECTION A:				
Slit lamp examination used	□ Yes	□ No		
Intraocular pressure (IOP) Baseline IOP results:	□ Yes	□ No		
Visual Acuity Test results:				
Labs and documentation to	□ Yes	□ No		
Anterior Chamber cells pr		□ Yes	□ No	
Initial Authorization: Please note member's diag	UVEITIS (NIU): (Please see 12 months gnosis. **NOTE: If member is tiology will be required for approximately see 12 months.	is only diagnos		
☐ Anterior Uveitis	☐ Intermediate Uveitis	□ Posterior	Uveitis	☐ Pan Uveitis
Is this member positive for HLA-B27 antigen? Please include other diagnosis that contributes to Anterior Uveitis ONLY diagnosis:				
□ Completed SECTIO	ON A			
☐ Diagnosis of chronic (1+ years) non-infectious uveitis affecting the posterior segment of the eye				
☐ Diagnosis and disease progression confirmed/determined by fluorescein angiography, Optical Coherence Tomography (OCT) or Scanning Computerized Ophthalmic Diagnostic Imaging (SCODI)				

(Continued on next page)

PAID	CLAIMS	MUST	MATCH	STATEN	MENT	BELOW:
------	---------------	-------------	--------------	---------------	------	---------------

Member must have inadequate response (i.e. recurrent uveitis despite use of traditional therapy), clinically significant adverse effects associated with high dose systemic steroid or immunosuppressive therapy, labeled contraindication, or clinical rationale supporting the inappropriateness of the following, include date(s) of failed therapy or clinical event(s). Documentation required.							
	Iember must meet at least <u>ONE</u> of the following:					east 4 weeks resulting in	
		Dexamethasone		☐ difluprednate (Durezol®) ☐ :		duoromethalone (FML®)	
□ loteprednol (Lotemax®) □ Oral pred			☐ Oral predn	isone or equivalent		orednisolone acetate (Pred Forte®)	
Name, dose and dates of the equivalent high does steroid trials:							
Tried and failed at least <u>ONE</u> immunosuppressive agent of 3 months due to toxicity OR failure to stabilize disease. (Submit supporting documentation of toxicities and progression, include label CBC, BUN, SCr, AST, ALT and albumin). Check ALL that apply:					•		
	☐ Adalimumab ☐ Azathioprin		thioprine	☐ Cyclosporine or tacrolimus		□ Infliximab	
	٥	Etanercept	□ Gol	imumab	☐ Methotrexate		□ Mycophenolate
		Rituximab	□ Toc	ilizumab			
☐ Member has received at least TWO administration or intra- or peri-ocular corticosteroid injecti					r corticosteroid injections		
	☐ Member has received at least <u>TWO</u> separate recurrences of uveitis requiring treatment with systemic corticosteroids or ocular injections of corticosteroids						
	Medication is being prescribed by a board-certified ophthalmologist or retinal specialist experienced in administration of intravitreal injections						
	Member will <u>NOT</u> be administered intravitreal implant simultaneously or with other intravitreal implants at the same time						
Me	mbe	r does <u>NOT</u> have a	any of th	e following con	ntraindications/exclus	ions to	either therapy:
	Hyp	ersensitivity to flu	ocinolor	ne or any of its	components		
	Ocu	lar or periocular in	fection				
	Adv	anced glaucoma					
	□ Concurrent intravitreal implants						

(Continued on next page)

IOI	N-INFECTIOUS UVEITIS (NIU): (Please submit supporting document to questions below)
lea ı	uthorization: 12 months
	Member continues to meet initial authorization criteria and has continued need for treatment (assessed and documented by provider)
	Reauthorization is being requested at least 30 months (Retisert®) or 36 months (Yutiq®) since previous implant
	Member has experienced a response to treatment as indicated by an improvement in uveitis and lack of recurrence within the preceding 30 months (Retisert®) or 36 months (Yutiq®)
	No unacceptable complications/toxicities due to implant have occurred (e.g., pain, hyperemia, decreased visual acuity, conjunctival hemorrhage)
	Therapy was NOT discontinued for any of the following reasons:
	☐ Loss of visual acuity from baseline
	☐ Severely increased intraocular pressure
	☐ Limited benefit of treatment
	☐ Unacceptable toxicities/complications to implant (eye pain, hyperemia, conjunctival hemorrhage)
	☐ Contraindications/exclusions:
	 Hypersensitivity to fluocinolone or its components
	Ocular or periocular infection
	Advanced glaucoma
	Concurrent intravitreal implant
Лed	lication being provided by (check box below that applies):
	Location/site of drug administration:
	NPI or DEA # of administering location:
	OR
	Specialty Pharmacy – PropriumRx

For urgent reviews: Practitioner should call Sentara Health Pre-Authorization Department if they believe a standard review would subject the member to adverse health consequences. Sentara Health's definition of urgent is a lack of treatment that could seriously jeopardize the life or health of the member or the member's ability to regain maximum function.

Use of samples to initiate therapy does not meet step edit/preauthorization criteria.

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.